

5
ENLARGED AND REVISED TO 1854.

THE
MEDICAL FORMULARY:

BEING

A COLLECTION OF PRESCRIPTIONS,

DERIVED FROM THE

WRITINGS AND PRACTICE OF MANY OF THE MOST EMINENT
PHYSICIANS IN AMERICA AND EUROPE.

TOGETHER WITH THE

USUAL DIETETIC PREPARATIONS AND ANTIDOTES FOR POISONS.

TO WHICH IS ADDED

AN APPENDIX,

ON THE

ENDERMIC USE OF MEDICINES, AND ON THE USE OF ETHER AND CHLOROFORM.

THE WHOLE ACCOMPANIED WITH A FEW BRIEF

PHARMACEUTICAL AND MEDICAL OBSERVATIONS.

BY BENJAMIN ELLIS, M. D.,

LATE PROFESSOR OF MATERIA MEDICA AND PHARMACY IN THE PHILADELPHIA COLLEGE OF PHARMACY.

TENTH EDITION,


REVISED AND MUCH EXTENDED.

BY ROBERT P. THOMAS, M. D.,

PROFESSOR OF MATERIA MEDICA IN THE PHILADELPHIA COLLEGE OF PHARMACY.

“Morbos autem, non eloquentiâ sed remediis curari.”—CELS. DE MED. LIB. I.

PHILADELPHIA:
BLANCHARD AND LEA.
1854.



QV
E47m
1854

Entered according to the Act of Congress, in the year 1853, by

BLANCHARD AND LEA,

in the Office of the Clerk of the District Court of the United States in and for
the Eastern District of Pennsylvania.

PHILADELPHIA:

T. K. AND P. G. COLLINS, PRINTERS.

TO
N. CHAPMAN, M. D.,

PROFESSOR OF THE INSTITUTES AND PRACTICE OF MEDICINE AND

CLINICAL PRACTICE

IN THE

University of Pennsylvania,

ETC. ETC.

WHOSE TALENTS AND URBANITY

HAVE RAISED HIM TO THE FIRST RANK IN THE PROFESSION, AND HAVE ACQUIRED FOR HIM
THE CONFIDENCE AND ESTEEM


OF A LARGE AND ENLIGHTENED COMMUNITY,

THIS WORK

IS VERY RESPECTFULLY DEDICATED

BY HIS OBLIGED FRIEND,

THE AUTHOR.



EDITOR'S PREFACE.

THE lamented decease of Dr. Samuel George Morton, since the appearance of the last edition, has rendered it necessary to place this work in other hands. During the lifetime of its author, Dr. Morton afforded a friendly revision of the first two editions, and after his death assumed the editorial charge of the seven succeeding editions. Being intimately acquainted with the author's views, he faithfully carried them out, by adding, from time to time, standard formulæ from the best writers.

Influenced by similar feelings, the present Editor has made a careful selection of a large number of prescriptions from the leading American and foreign authorities; in most instances, giving a preference to those which serve to introduce the new remedies announced in the last few years. Some of the old formulæ have been omitted, and others less complicated have been substituted. To all of the additions, the word "EDITOR" or the names of their authors have been scrupulously placed.

The arrangement of the original work, founded on that adopted by Dr. Chapman in his *Therapeutics*, has been retained for internal remedies, which constitute the subject-matter of Book I. But for convenience of refer-

ence, it has been thought advisable to bring together under a common head those intended for external use. Accordingly, BOOK II. has been devoted to external applications. This course appeared the more indispensable from the fact that the true therapeutic value of a large number of the articles used in the skin diseases has not yet been tested by time and experience.

The table of the doses of medicines has been entirely rewritten, and the present Editor is alone responsible for any errors therein contained.

The various changes in nomenclature that have occurred during twenty years, have caused much discrepancy in prescribing, and much difficulty in compounding. To obviate these, all the formulæ embracing officinal articles have been brought to the standard of our National Pharmacopœia, for the double purpose of extending a practical knowledge of its terminology among prescribers, and of facilitating as well as simplifying the duties of the apothecary.

ROBERT P. THOMAS.

PHILADELPHIA, October, 1853.

*Dec will be received before September
16th 1846.*

September 15th 1846

CONTENTS.

*John
Allen*

	PAGE
Preface of the Editor	v
Preface of the Author	xi
Introduction	xv
Table of Drops	xvii
Table of Abbreviations	xix
Table of Doses for Children	xx
Table of the Doses of Medicines	xxi
Tabular View of the Doses of the Principal Articles of the Materia Medica	xxiii

BOOK I.

OF INTERNAL REMEDIES.

CLASS I. Emetics	35
<i>Powders</i>	36
<i>Wines and Tinctures</i>	38
<i>Solutions and Mixtures</i>	39
<i>Various Emetic Preparations</i>	42
II. Cathartics	43
<i>Powders</i>	44
<i>Pills</i>	47
<i>Confections and Electuaries</i>	52
<i>Emulsions</i>	53
<i>Mixtures</i>	55
<i>Infusions and Decoctions</i>	58
<i>Wines and Tinctures</i>	60
<i>Enemata</i>	61
<i>Suppositories</i>	63
III. Diuretics	65
<i>Powders</i>	65
<i>Pills</i>	66
<i>Infusions and Decoctions</i>	68
<i>Solutions and Mixtures</i>	70

CLASS	PAGE
IV. Antilithics and Lithontriptics	76
<i>Powders</i>	76
<i>Pills</i>	77
<i>Solutions and Mixtures</i>	77
V. Diaphoretics	79
1. Ordinary Diaphoretics	80
<i>Powders</i>	80
<i>Mixtures</i>	82
<i>Infusions</i>	86
2. Alterative Diaphoretics	87
<i>Pills</i>	87
<i>Decoctions</i>	88
<i>Mixtures</i>	90
VI. Expectorants	92
<i>Powders, Pills, and Lozenges</i>	92
<i>Sirups</i>	96
<i>Mixtures</i>	98
<i>Inhalation</i>	107
VII. Emmenagogues	110
<i>Powders and Pills</i>	110
<i>Decoctions and Infusions</i>	113
<i>Tinctures</i>	114
VIII. Anthelmintics	117
<i>Powders and Pills</i>	117
<i>Infusions and Decoctions</i>	120
<i>Electuaries and Mixtures</i>	121
<i>Clysters</i>	123
IX. Stimulants	124
<i>Powders and Pills</i>	124
<i>Mixtures and Infusions</i>	126
X. Narcotics	133
<i>Pills</i>	133
<i>Mixtures and Tinctures</i>	137
<i>Clysters and Suppositories</i>	142
<i>External Use of Narcotics</i>	143
XI. Antispasmodics	147
XII. Sialagogues	151
XIII. Tonics	155
1. Anti-intermittents	155
<i>Powders and Pills</i>	155
<i>Decoctions and Infusions</i>	159
<i>Solutions and Mixtures</i>	161
2. Ordinary tonics	164
<i>Powders and Pills</i>	164
<i>Decoctions and Infusions</i>	170
<i>Mixtures and Tinctures</i>	172

CONTENTS.

ix

CLASS	PAGE
XIV. Alteratives	177
<i>Powders and Pills</i>	177
<i>Tinctures and Mixtures</i>	181
XV. Astringents	190
<i>Powders and Pills</i>	190
<i>Mixtures and Infusions</i>	194
XVI. Absorbents or Antacids	198
<i>Powders and Pills</i>	198
<i>Mixtures and Solutions</i>	199

BOOK II.

OF EXTERNAL APPLICATIONS.

XVII. Caustics or Escharotics	205
Issues and Setons	209
XVIII. Epispastics	211
XIX. Rubefacients	215
XX. Anomalous External Applications	222
A. Collyria	222
<i>Powders</i>	222
<i>Solutions and Mixtures</i>	223
B. Injections	228
a. <i>For the Ear</i>	228
b. <i>For the Urethra</i>	230
C. Gargles and Mouth Washes	233
D. Liniments, Lotions, &c.	237
E. Ointments	245
<i>Powders</i>	252
F. Cataplasms	253
G. Baths	255
H. Fomentations	257
I. Fumigations	258

BOOK III.

OF DIETETIC PREPARATIONS AND POISONS.

Dietetic Preparations	261
Poisons	269
APPENDIX I. Endermic Use of Medicines	278
II. Use of Ether and Chloroform	280
Index	285

P R E F A C E .

THE elegant and judicious formation of prescriptions is one of the difficulties which the young practitioner in medicine is obliged to encounter. While a student, he is compelled, from the circumstances under which he is placed, to confine his attention to the leading principles of the science. Consequently, the minor points (and the art of directing medicines is considered one of them) are postponed to that period when he shall have undertaken the practical duties of his profession.

To obviate, in some measure, the inconvenience which the graduate at first experiences, the volume now offered the public was undertaken and executed. It contains, in a condensed form, and we think advantageously arranged, many of the most important prescriptions employed in modern practice, and which, though most of them may be already recorded, are scattered through various medical works.

The arrangement framed by Professor Chapman for his *Therapeutics*, appearing to combine greater advantages than any other, we have taken the liberty of adopting it as the basis of this *Formulary*.

Each Class has been subdivided into Powders, Pills, and Liquids; without any attempt to preserve the different preparations of the same article together, but only with a view to convenience. A few concise rules will be found prefixed to each Class, and observations have been attached to those prescriptions which possess peculiarities either in their composition or effects.

The application of the remedies to diseases has been generally left to the judgment of the practitioner, and therapeutical detail as much as possible avoided, as it would have been inconsistent with the nature and design of the work.

The object is to furnish models for extemporaneous formulæ, the proportions of which may be either increased or diminished, according to the age, sex, constitution or idiosyncrasy of the patient, all of which are to be taken into consideration when a prescription is directed, as well as the climate and peculiar epidemic which may be prevailing.

The prescriptions have been couched in the Latin language, and the names of the several articles will be found regularly terminated. The design was to furnish something like a Medical Grammar, to those unacquainted with the dead language; and even those versed in the classics, may occasionally find words used in prescriptions which are new to them.

A series of Dietetic preparations and the principal poisons have been placed in Book III. The former are written in the English language, and it is hoped that they may be found useful in furnishing directions necessary to be given to the nurse.

The latter are those Poisons which are usually taken by accident or design, with their appropriate antidotes and some practical observations.

Appendix No. I. contains some concise remarks on the mode of using remedies on the "*Endermic plan*," as well as the quantities employed, and the effects produced by several of the most potent remedial agents.¹

The materials for the work have been collected from various sources, the writings of different authors, periodical journals, manuscripts, and the contributions of our friends. It is hoped, therefore, that the difficulties which have attended its progress will form some apology for the errors which it may possibly be found to contain. As a reason for not more frequently attaching the names of authors to the prescriptions, it may be observed that they are a kind of common medical property, which every physician feels himself privileged not only to employ, but modify as may best correspond with his own views. It is, therefore, difficult to say to whom many of them belong; they have been touched, and retouched, till they have lost their identity.

In conclusion, we beg leave to present our acknowledgments to some of our friends for the encouragement they have given to the undertaking. To Dr. Chapman we are particularly indebted for the kindness and interest he has manifested in its publication, and offer to him (the only return we can make), our warmest thanks.

¹ Appendix No. II. is a concise and able article on the use of Ether and Chloroform, drawn up by the late Dr. Morton, to which the attention of the reader is directed.

It is with particular pleasure we embrace this opportunity of expressing our obligations to our friend Dr. S. G. Morton, whose intimate acquaintance with the subject, and classical knowledge, are only equalled by the kindness and industry he has displayed in facilitating the work.

There are others whose names it would gratify us to mention, but who, we know, will be more pleased by our silence. The countenance they gave us on the present occasion has only increased the debt previously incurred.

B. E.

INTRODUCTION.

BEFORE entering upon the main object of this work, it is necessary to notice a few circumstances which pertain more especially to Pharmaceutic Chemistry, yet appear to be called for in this place ; and though they are familiar to every educated physician, may be useful to the student who has not yet turned his attention to prescriptions.

In the first place, it is requisite to be well acquainted with the *signs of quantity*, or those symbolic characters which are used in writing prescriptions. In the ancient practice of medicine, there were very many of these symbols, not only for the purpose of denoting the quantity of the ingredient, but the ingredient itself. Most of these, however, have become obsolete ; and a few only are retained as indispensable to the convenience of the practitioner. These will be found in the following *Table*, which is adopted in the Pharmacopœias for the regulation of *weights*.

English Name.	Latin Name.	Symbol.	Contains.
A pound,	Libra,	℔,	12 ounces.
An ounce,	Uncia,	℥,	8 drams.
A dram,	Drachma,	ʒ,	3 scruples.
A scruple,	Scrupulus,	ʒ,	20 grains.

In the manipulation of *liquid substances*, *measures* have in most countries superseded the use of *weights*. The *Table of Measures* recommended by the London College, gives particular names and symbols to the parts forming a gallon, in order to prevent their being confounded with those denoting weights; as in the following *Table* :—

English Name.	Latin Name.	Symbol.	Contains.
A gallon,	Congius,	none,	8 pints.
A pint,	Octarius,	O,	16 fluidounces.
A fluidounce,	Fluiduncia,	f 3,	8 fluidrams.
A fluidram,	Fluidrachma,	f 3,	60 minims.

The *minims* used in the Table, have been adopted in the United States Pharmacopœia in place of *drops*; and although we have not employed this measure in our prescriptions, yet we can readily perceive it to possess some important advantages over the ancient practice. The symbol (℥) is sometimes used by medical writers, though very improperly, to denote both minims and drops.¹ The size of a drop varies according to the greater or less fluidity and gravity of the liquid, and the shape of the vessel from which it is dropped; but, as a general rule, we cannot do better on these occasions than use a small phial with a thin edge. Dr. Dorsey observes that, whenever great precision is necessary, it is easy to dilute the active medicine, and give it in form of a mixture. If sixty grains be diffused in six ounces of water, for instance, each half ounce will contain five.

¹ Ten minims of most fluids, if measured by the graduated minim measure, are equal to about fifteen drops.

Mr. Durand, of this city, has made various experiments to ascertain *the number of drops of different liquids equivalent to a fluidram*; and his results, which are of practical importance, are given in the following table:—

	Drops.
Acetone,	110
Acid acetic (crystallizable),	120
Acid hydrocyanic (medicinal),	45
Acid muriatic,	54
Acid nitric,	84
Acid nitric, diluted (1 to 7),	91
Acid sulphuric,	90
Acid sulphuric aromatic,	120
Acid sulphuric, diluted (1 to 7),	54
Alcohol, rectified spirit,	138
Alcohol diluted, proof spirit,	120
Arsenite of potassa, solution of,	57
Chloroform,	260
Ether sulphuric,	150
Oil of aniseed, of cinnamon, of cloves, of peppermint, of sweet almond, of olives,	} 120
Tincture of assafetida, of foxglove, of guaiac, of opium,	
Tincture of muriate of iron,	132
Vinegar distilled,	78
Vinegar of colchicum,	78
Vinegar of opium (black drop),	78
Vinegar of squills,	78
Water distilled,	45
Water of ammonia (strong),	54
Water of ammonia (weak),	45
Wine Teneriffe,	78
Wine antimonial,	72
Wine of colchicum,	75
Wine of opium,	78

Now it is evident from the preceding facts, that in

giving 150 drops of sulphuric ether (and proximately of any ethereal tincture), we give but a fluidram; whereas, the same number of drops of medicinal hydrocyanic acid (one of the most poisonous preparations of the *materia medica*), would be equal to something more than three fluidrams. The same remark, it will be observed, is applicable to various other preparations; which suggests a caution in rapidly increasing the dose of such medicines by drops.

Whenever a *tablespoonful* of any liquid is ordered, it is considered to be equal to *half an ounce* by measure; and in the same way a *teaspoonful* is used for a *dram*: these measures are sufficiently accurate where no great precision is requisite.

The prescriptions contained in this work are given in the Latin language, without other abbreviations than those in common use to denote the *quantity* of an ingredient. The directions for the administration of medicines are expressed in English.

Prescriptions, as usually sent to the Apothecaries, are much abbreviated, and the same course may be pursued with regard to those contained in this collection. For the information of students who have not been in the practice of writing prescriptions, a *Table of Abbreviations* is subjoined.

TABLE OF ABBREVIATIONS.

Abbreviation.	Latin Word.	English Word.
āā.	¹ Ana,	Of each.
Ad saturand.	Ad saturandum,	Until saturated.
Ad lib.	Ad libitum,	At pleasure.
Aq. tepid.	Aqua tepida,	Warm water.
Aq. ferv.	Aqua fervens,	Hot water.
C.	Congius,	A gallon.
Chart.	Chartula,	A small paper.
Coch.	Cochlear,	A spoonful.
Coch. mag.	Cochlear magnum,	A tablespoonful.
Coch. parv.	Cochlear parvum,	A teaspoonful.
Colent.	Colentur,	Let them be strained.
Collyr.	Collyrium,	An eye-water.
Contus.	Contusus,	Bruised or broken.
F. vel Ft.	Fiat, vel fiant,	Let there be made.
Fol.	Folium, vel folia,	A leaf, or leaves.
Garg.	Gargarysma,	A gargle.
Gr.	Granum, vel grana,	A grain, or grains.
Gtt.	Gutta, vel guttæ,	A drop, or drops.
Haust.	Haustus,	A draught.
Infus.	Infusum,	An infusion.
M.	Misce,	Mix.
Mass.	Massa,	A mass.
Mist.	Mistura,	A mixture.
O.	Octarius,	A pint.
Pil.	Pilula, vel pilulæ,	A pill, or pills.
Pulv.	Pulvis, vel pulveres.	A powder, or powders.
Q. S.	Quantum sufficit,	A sufficient quantity.
R.	Recipe,	Take.
Rad.	Radix,	A root.
S.	Signa,	Write.
Ss.	Semi, vel dimidius,	The half.
Tinct.	Tinctura,	A tincture.

The *doses* throughout this work are applicable to adult age, unless the contrary is specified.

For the convenience of young practitioners, the rules

¹ This is not properly a Latin word, but the Greek preposition *ana*.

furnished by Gaubius and Dr. Young, for ascertaining the proper doses of medicines for children, are given below.

Gaubius takes the dose for an adult as unity, and for other ages, as follows :—

One year old,	. . .	$\frac{1}{13}$	Seven years old	. . .	$\frac{1}{3}$
Two years “	. . .	$\frac{1}{8}$	Fourteen “ “	. . .	$\frac{1}{2}$
Three “ “	. . .	$\frac{1}{6}$	Twenty “ “	. . .	$\frac{2}{3}$
Four “ “	. . .	$\frac{1}{4}$	From twenty to sixty years old	1	

Dr. Young says : “For children under twelve years, the doses of most medicines must be diminished in the proportion of the age, to the age increased by 12.” Thus, for a child of two years, $= 2 : 2 + 12 ::$ the adult dose, or 1 : to the child’s dose, or $\frac{2}{14}$. Or, to state it more simply, $\frac{2}{2+12} = \frac{1}{7}$. Hence,

$$\text{For one year } \frac{1}{1+12} = \frac{1}{13}$$

$$\text{“ two years } \frac{2}{2+12} = \frac{1}{7}$$

$$\text{“ three “ } \frac{3}{3+12} = \frac{1}{5}$$

$$\text{“ four “ } \frac{4}{4+12} = \frac{1}{4}$$

$$\text{“ six “ } \frac{6}{6+12} = \frac{1}{3}$$

&c. &c.

At twelve, the dose is one-half that of the adult.

TABLE

OF THE

DOSES OF MEDICINES ARRANGED IN ALPHABETICAL ORDER.

IN preparing the following Table, a great object has been to present, at a glance, the minimum and medium doses of each medicine ; or, as sometimes happens in respect to the more familiar articles, the ordinary dose, as in many of the infusions and tinctures. It is necessary to premise that these preparations are adapted to adult age ; nor can any precise rule be laid down which will be applicable to all the periods of life between that period and early infancy. Nothing but the cautious observation and judgment of the physician can possibly supply this deficiency. The best mode of giving active medicines to infants is in solution or suspension, so that the dose may be readily and safely graduated in reference to age, constitution, and disease. As the same medicine will sometimes be found in several different classes of the *Materia Medica*, its dose must be varied accordingly. Thus, ipecacuanha is both an emetic and diaphoretic, and the large dose we give for the former indication would not bear the necessary repetition for the second. Calomel is purgative and alterative ; but in the table we

only give the dose necessary to produce the first of these effects; and so also of many other medicines. Here again the formulæ, as given under the several Classes, provide the necessary information in detail.

We would also impress upon every practitioner, the vital importance of knowing the properties and strength of a medicine before prescribing it; and this caution is especially requisite in directing those preparations, which in an overdose, produce distressing and sometimes fatal effects. No prescription of this kind should ever be copied at random, and without first satisfying one's self, by a little calculation, whether the proportions directed in it are safe and consistent; and we would farther insist that the use of every poisonous medicine should be commenced in a *minimum* dose, and gradually increased according to its effects on the system. It is often necessary, also, in these instances, to put the patient on his guard in case any inordinate or unusual symptoms should arise, with a request that the prescription, in such contingency, may be taken in less quantity, at longer intervals, or discontinued. This precaution is all-important, for example, in the administration of alterative doses of the mercurial preparations; for the same quantity that would have little or no effect on one person, might, and often does produce a violent and deleterious effect on another. This remark is also equally applicable to the whole class of narcotic medicines.

In truth, nothing but *habitual circumspection* in prescribing medicines, can render this art beneficial to the patient, or honorable to the physician.

TABULAR VIEW

OF THE

DOSES OF THE PRINCIPAL ARTICLES OF THE MATERIA MEDICA.

Absinthium, ℥j. to ij.	Ætheris Nitrici, Spiritus, f℥j.
Acetinum, gtt. x. to xxx.	Allium, ℥ss. to ℥j.
Acetum, f℥j. to iv.	Allii, Succus, f℥ss.
Acidum Arseniosum, gr. $\frac{1}{10}$.	——, Syrupus, f℥j. to iv.
—— Benzoicum, gr. x. to xxx.	Aloe, gr. x. to xx.
—— Boracicum, gr. v. to xx.	Aloe et Canella, gr. v. to xx.
—— Gallicum, gr. v. to x.	Aloës, Tinctura, f℥ss.
—— Hydrocyanicum dilutum, gtt. j. to ij.	——, Vinum, f℥j. to f℥ss.
—— Muriaticum dilutum, gtt. xx. to lx.	Alumen, gr. x. to xx.
—— Nitricum dilutum, gtt. x. to xl.	—— Exsiccatum, gr. v. to x.
—— Nitro-muriaticum, gtt. iij. to iv.	Ammoniac Acetatis, Liquor, f℥ss. to j.
—— Sulphuricum dilutum, gtt. x. to xxv.	—— Carbonas, gr. v. to x.
—— Sulphuricum Aromaticum, gtt. x. to xxv.	—— Phosphas, gr. x. to xl.
—— Tannicum, gr. ij. to v.	—— Murias, gr. v. to xv.
—— Tartaricum, ℥j. to ij.	—— Aromaticus, Spiritus, gtt. xx. to f℥j.
Aconiti Folia, gr. j. to ij.	Ammoniacum, gr. x. to xx.
——, Extractum, gr. j.	Ammoniaci, Mistura, f℥ss.
——, Extractum Alcoholicum, gr. $\frac{1}{2}$.	Angustura, gr. x. to xxx.
—— Foliorum, Tinctura, gtt. xx.	Angusturæ, Infusum, f℥ij.
—— Radicis, Tinctura, gtt. v. to viij.	Anthemis, ℥ss. to j.
Æther, gtt. l. to f℥j.	Anthemidis, Infusum, f℥j. to ij.
——, Spiritus Compositus, f℥j.	Antimonii Sulphur. Præcipitatum, gr. j. to v.
	Antimonii, Oxidum, gr. iij. to v.
	—— et Potassæ Tartras, gr. $\frac{1}{2}$ to iij.
	——, Vinum, gtt. x. to f℥j.
	Antimonialis, Pulvis, gr. iij. to x.
	Apocynum, gr. x. to xxx.
	Argentii Nitras, gr. $\frac{1}{2}$.

Argenti Cyanuretum, gr. $\frac{1}{15}$.	Cantharis, gr. j. to ij.
—— Iodidum, gr. $\frac{1}{2}$ to ij.	Cantharidis, Tinctura, gtt. x. to xxx.
—— Oxidum, gr. $\frac{1}{2}$.	Capsicum, gr. v. to x.
Arnica, gr. v. to xx.	Capsici, Tinctura, f3ss.
Arseniosum Acidum, gr. $\frac{1}{10}$.	Cardamomum, gr. x. to xxx.
Arsenici Iodidum, gr. $\frac{1}{2}$.	Carum, gr. xx. to xl.
Arsenici et Hydrargyri Iodidi,	Caryophyllus, gr. v. to x.
Liquor, gtt. v. to x.	Caryophylli Oleum, gtt. ij.
—— Potassæ Arsenitis, Liquor,	Cascarilla, gr. xx. to xxx.
gtt. x.	Castoreum, gr. x. to xx.
Asclepias Tuberosa, gr. xx. to 3j.	Castorei Tinctura, gtt. xx. to l.
Assafoetida, gr. x.	Catechu, gr. x. to xxx.
Assafoetidæ, Mistura, f3ss.	——, Tinctura, f3j. to ij.
——, Tinctura, f3j.	Cerevisiæ Fermentum, f3ss. to ij.
Atropia, gr. $\frac{3}{10}$.	Cetrariæ, Decoctum, f3ij.
Auri Sales, gr. $\frac{1}{18}$ to $\frac{1}{10}$.	Chenopodium, ʒj. to ij.
Balsamum Peruvianum, f3ss.	Chenopodii, Oleum, gtt. v. to x.
—— Tolutanum, gr. x. to xxx.	Chimaphilæ, Decoctum, f3ij.
—— Tolutanus Syrupus,	Chloroformum, gtt. xx. to l.
f3ss.	Chondri, Decoctum, ad libitum.
—— Tolutana Tinctura, f3j.	Cimicifuga, ʒj. to 3j.
Barii Chloridi, Liquor, gtt. v.	Cinchona, gr. xx. to xxx.
Bebeerinæ Sulphas, gr. iij. to xx.	Cinchonæ, Extractum, gr. x. to xx.
Belladonna, Folia, gr. j.	——, Infusum, } f3j. to ij.
Belladonnæ, Extractum, gr. $\frac{1}{4}$.	vel } f3j. to ij.
——, Extractum Alcoholic.	——, Decoctum, } f3j. to iv.
gr. $\frac{1}{2}$.	——, Tinctura, f3j. to iv.
——, Tinctura, gtt. xv. to xx.	——, Tinctura Composita,
Bismuthi, Subnitrates, gr. v. to x.	f3j. to iv.
Brominium, gtt. $\frac{1}{4}$.	Cinnamomum, gr. x. to xx.
Brucia, gr. $\frac{1}{8}$ to $\frac{1}{4}$.	Cinnamomi, Aqua, f3ss.
Buchu, gr. xx. to xxx.	——, Oleum, gtt. j. to ij.
——, Infusum, f3ij.	——, Tinctura Composita,
Cahina, ʒj. to 3j.	f3j. to ij.
Calamus, ʒj. to 3j.	Coccus, gr. j. to x.
Calceis, Liquor, f3j. to ij.	Colchici Radix, gr. ij. to vj.
Calcei Chloridi, Liquor, gtt. xxx. to f3j.	—— Semen, gr. ij. to vj.
Calx Chlorinata, gr. iij. to vj.	—— Radicis, Vinum, gtt. x. to xx.
—— Creta præparata, gr. x. to xxx.	—— Seminis, Vinum, f3j.
—— Testa præparata, gr. x. to xxx.	—— Seminis, Tinctura, f3ss. to j.
Camphora, gr. v. to x.	Colchici, Acetum, f3ss. to j.
Camphoræ, Aqua, f3ss.	——, Extractum Aceticum, gr. j.
——, Tinctura, gtt. v. to xx.	Colocynthis, gr. v. to x.
Canella, gr. x. to xxx.	Colocynthidis, Extractum Compositum, gr. x. to xv.
	Colomba, gr. x. to xxx.
	Colombæ, Infusum, f3ij.

- Colombæ, Tinctura, f3j. to iv.
 Conii Folia, gr. iij.
 —, Extractum, gr. ij.
 —, Tinctura, grt. xx. to xxx.
 Copaiba, f3ss. to j.
 Copaibæ, Oleum, gtt. x. to xv.
 Coptis, gr. x. to xxx.
 Coriandrum, ʒj. to ʒj.
 Creasotum, gtt. j. to ij.
 Creta præparata, gr. xv. to ʒj.
 Crocus, gr. x. to xxx.
 Cubeba, ʒss. to iij.
 Cubebæ, Extractum Fluidum, ℥x.
 to xxx.
 —, Oleum, gtt. x. to xij.
 —, Tinctura, f3ss. to j.
 Cuprum Ammoniatum, gr. ¼ to ½.
 Cupri Sulphas, gr. ¼ to ij.

 Dippel's Animal Oil, gtt. v. to x.
 Digitalis, gr. j.
 —, Infusum, f3ij. to iv.
 —, Tinctura, gtt. x.
 Digitalin, gr. ⅙.
 Donovan's Solution, gtt. v. to xx.
 Dover's Powder, gr. x.
 Dracontium, gr. x. to xx.
 Dulcamara, ʒss. to j.
 Dulcamaræ Decoctum, f3ij.

 Elaterium, gr. ½.
 Elaterin, gr. ⅙.
 Ergota, gr. xx.
 Ergotæ, Vinum, f3j. to ij.
 Ether, gtt. l. to f3j.
 Eupatorii, Infusum, f3ij.
 Extractum sive Resina Cannabis,
 gr. ¼ to ½.

 Fel Bovinum, gr. v. to x.
 Ferri Pulvis, gr. iij. to v.
 — Chloridi, Tinctura, gtt. x. to
 xxv.
 — Citras, gr. v. to x.
 — Ammonio-Citras, gr. v.
 — et Potassæ Tartras, gr. x. to
 xx.
 — Ferrocyanturetum, gr. iij. to v.
 — Iodidi, Liquor, gtt. x. to l.
 — Nitratis, Liquor, gtt. viij. to x.
- Ferri Phosphas, gr. v. to x.
 — Subcarbonas, gr. x. to ʒij.
 — Carbonatis, Pilulæ, gr. x.
 — Sulphas exsiccatus, gr. ij.
 to v.
 — Valerianas, gr. j.
 — Vinum, f3ss.
 Ferrum Ammoniatum, gr. v. to x.
 Filix Mas, ʒj. to ij.
 Fœniculum, ʒj. to ʒss.
 Fœniculi, Aquæ, f3ss.
 Fowler's Solution, gtt. x.

 Galla, gr. x. to xx.
 Gambogia, gr. ij. to iij.
 Gaultheriæ Oleum, gtt. ij. to v.
 Gentiana, gr. x. to xxx.
 Gentianæ, Extractum, gr. x. to xx.
 — Composita, Tinctura, f3j.
 to ij.
 Gillenia, gr. xx. to xxx.
 Granati Fructus Cortex, gr. xx. to
 xxx.
 Guaiaci Resina, gr. x. to xx.
 — Ammoniata, Tinctura, f3j.

 Hæmatoxyli, Extractum, gr. x. to
 xx.
 Helleborus, gr. v. to xv.
 Hellebori, Extractum, gr. v. to x.
 —, Tinctura, gtt. xxx. to f3j.
 Hoffman's Anodyne, f3j.
 Hordei Decoctum, ad libitum.
 Humuli, Tinctura, f3ij.
 Lupulina, gr. v. to x.
 Lupulinæ, Tinctura, f3j.
 Hydrargyri Chloridum Corrosi-
 vum, gr. ⅓ to ½.
 — Chloridum mite, gr. ¼
 to x.
 — Cyanuretum, gr. ⅙ to ½.
 — Iodidum, gr. ½ to j.
 — Iodidum Rubrum, gr.
 ⅙.
 — Oxidum Nigrum, gr.
 ½ to iij.
 — Oxidum Rubrum, not
 used internally.
 — Sulphas Flavus, gr. ¼
 to v.

Hydrargyri Sulphuretum Nigrum, gr. v. to x.	Lugol's Solution, gtt. v. to x.
———— Sulphuretum Rubrum, gr. x.	Lupulina, gr. v. to x.
Hydrargyrum cum Creta, gr. v. to xv.	Lupulinæ, Tinctura, f3j.
———— cum Magnesia, gr. v. to xv.	Magnesia, 3ss. to j.
Hyoscyami Folia, gr. v. to x.	Magnesiae Carbonas, 3ss. to j.
————, Extractum, gr. ij.	———— Citratis Liquor, f3iv. to vj.
————, Tinctura, f3ss. to j.	———— Sulphas, 3j.
Inula, ʒj. to 3j.	Magnolia, 3ss. to j.
Iodinium, gr. ¼ to j.	Manganesii Oxidum gr. iij. to x.
Iodinii, Tinctura, gtt. v. to x.	———— Iodidi, Liquor, gtt. x. to xxx.
Iodinii Compositus, Liquor, gtt. v. to x.	———— Sulphas, ʒj. to 3ij.
———— Composita, Tinctura, gtt. xv. to xxx.	Manna, 3j.
Ipecacuanha, gr. ij. to xx.	Mannite, 3j.
Ipecacuanhæ, Syrupus, f3j. to ij.	Matico, 3ss. to ij.
————, Vinum, f3j. to iv.	Matricaria, 3ss. to j.
Ipecacuanhæ et Opii, Pulvis, gr. x.	Menthæ Piperitæ, Oleum, gtt. j. to ij.
Jalapa, gr. x. to xx.	————, Tinctura, gtt. x. to xx.
Jalapæ, Extractum, gr. x. to xv.	————, Aqua, f3ss.
————, Tinctura, f3j. to ij.	Menthæ Viridis, Aqua, f3ss.
Juglandis, Extractum, gr. xx. to xxx.	Mezerei Decoctum, f3ij. to iv.
Juniperi, Baccæ, 3j. to ij.	Mistura Cretæ, f3ss. to j.
————, Oleum, gtt. v. to xv.	Monardæ Oleum, gtt. ij. to iij.
———— Compositus, Spiritus, f3j. to iv.	Morphiæ Acetas, gr. ½.
Kino, gr. x. to xx.	———— Murias, gr. ½.
————, Tinctura, f3j. to ij.	———— Sulphas, gr. ½.
Kousso, Flores, 3ss.	———— Sulphatis, Liquor, f3j. to ij.
Krameria, gr. xx. to xxx.	Moschus, gr. x.
Krameria, Extractum, gr. x. to xx.	Mucunæ Electuarium, f3j. to f3ss.
————, Syrupus, f3j. to ij.	Myrrha, gr. x. to xx.
Lactucarium, gr. iij. to x.	Myrrhæ Tinctura, f3ss. to j.
Lappa, 3j.	Naphthaline, gr. ij. to xv.
Lauro-Cerasi, Aqua, f3ss. to j.	Nucis Vomica, Extractum, gr. ss. to j.
Lavandulæ, Oleum, gtt. ij. to v.	Nux Vomica, gr. iij. to v.
———— Compositus, Spiritus, f3ss. to j.	Oleum Amygdalæ Amaræ, gtt. ¼ to j.
Liriodendron, 3ss. to ij.	———— Anisi, gtt. v. to x.
Lobelia, gr. v. to x.	———— Cajuputi, gtt. ij. to v.
Lobeliæ Tinctura, gtt. xx. to f3j.	———— Limonis, gtt. ij. to iv.
	———— Morrhæ, f3ss.
	———— Olivæ, f3j.
	———— Pimentæ, gtt. ij. to v.
	———— Ricini, f3j.
	———— Rosmarini, gtt. ij. to iij.

Oleum Rutæ, gtt. ij. to iij.	Pruni Virginianæ, Infusum, f 3ij.
——— Sabinae, gtt. ij. to iij.	——, Syrupus, f 3ss.
——— Sassafras, gtt. iv. to vj.	Pulvis Aromaticus, gr. x. to xx.
——— Succini Rectificatum, gtt. v. to xv.	——— Doveri, gr. x.
——— Terebinthinæ, gtt. x. to f3ij.	Quassia, Extractum, gr. ij. to v.
——— Tiglii, gtt. ½ to ij.	——, Tinctura, f 3j. to ij.
Opium, gr. j.	Quercus Alba, et Q. Tinctoria, gr. xx. to 3j.
Opii, Acetum, gtt. x.	Quiniæ Sulphas, gr. j. to x.
——, Confectio, gr. x.	—— Valerianas, gr. ij. to vj.
——, Extractum, gr. ss.	—— et Ferri Citras, gr. iij. to x.
——, Tinctura, gtt. xx. to xxv.	Rhamni Syrupus, f 3ss.
—— Acetata, Tinctura, gtt. xx.	Rheum, ʒj. to 3ss.
—— Camphorata, Tinctura, f 3j. to ij.	Rhei Fluidum, Extractum, f 3ss.
——, Vinum, gtt. xx. to xxv.	—— Aromaticus, Syrupus, f 3j. to iv.
——, McMunn's Elixir, gtt. xx. to xxv.	——, Tinctura, f 3ss.
Pareira, 3ss. to j.	——, Vinum, f 3ij.
Petroleum f3ss. to 3j.	Rubia, ʒj. to 3ss.
Phloridzin, gr. x. to xij.	Rubus Villosus, et R. Trivialis, ʒj. to 3ss.
Piperis Fluidum, Extractum, gtt. j. to iij.	Sabbatia, 3ss. to j.
Pix Liquida, 3ss. to j.	Sabina, gr. v. to x.
Plumbi Acetas, gr. j. to iij.	Sabinae Oleum, gtt. ij. to iij.
—— Iodidum, gr. ss. to iij.	Salicin, gr. ij. to x.
—— Nitras, gr. ¼ to ½.	Sanguinariæ Tinctura, f 3ss. to j.
Podophylli, Extractum, gr. v. to xv.	Santonine, gr. iij. to iv.
Podophyllin, gr. ss. to ij.	Sarsaparillæ, Extractum, gr. x. to xx.
Potassæ Liquor, gtt. x. to xx.	—— Fluidum, Extractum, f 3j.
—— Arsenitis, Liquor, gtt. x.	—— Compositus, Syrupus, f 3ss.
—— Acetas, ʒj. to 3ij.	Scammonium, gr. v. to xv.
—— Bitartras, 3j. to ij.	Scilla, gr. j. to ij.
—— Carbonas Purus, gr. x.	Scillæ, Acetum, f 3ss. to j.
—— Bicarbonas, gr. x. to 3ss.	——, Syrupus, f 3j.
—— Chloras, gr. x. to xxx.	—— Compositus, Syrupus, f 3ss. to j.
—— Citras, gr. xx. to xxv.	——, Oxymel, f 3j. to ij.
—— Citratis, Liquor, f 3ss.	Senega, gr. x. to xx.
—— Nitras, gr. x. to xv.	Senegæ Syrupus, f 3j. to ij.
—— Sulphas, 3ss. to j.	Senna, 3ss. to 3ij.
—— Tartras, 3j. to 3ss.	Sennæ Confectio, 3j. to ij.
Potassii Bromidum, gr. iij. to x.	—— Fluidum, Extractum, f 3ss.
—— Cyanuretum, gr. ¼ to ½.	Serpentaria, gr. x. to xxx.
—— Ferrocyanuretum, gr. x. to xv.	Serpentariæ Tinctura, f 3j.
—— Iodidum, gr. iij. to x.	
—— Sulphuretum, gr. ij. to vj.	
Prunus Virginiana, 3ss. to j.	

Simaruba, ʒj. to ʒj.	Tabacum, gr. j. to v.
Sinapis (as an emetic), ʒij.	Tabaci, Vinum, gtt. x. to xx.
Sodæ Acetas, ʒj. to ʒj.	Taraxaci, Extractum, ʒj. to ʒj.
—— Boras, ʒj. to ij.	Terebinthina, ʒj. to ij.
—— Carbonas, gr. v. to xx.	Testa præparata, gr. x. to l.
—— Bicarbonas, gr. x. to xxx.	Ulmi, Infusum, ad libitum.
—— Phosphas, ʒss. to j.	Uva Ursi, ʒss. to j.
—— Sulphas, ʒss.	
—— et Potassæ Tartras, ʒij. to ʒss.	Valeriana, ʒss. to j.
Sodii Chloridum, gr. x. to ʒj.	Valerianæ Fluidum, Extractum, fʒj.
Spigelia, gr. x. to ʒj.	——, Oleum, gtt. v.
Spigeliæ et Sennæ Fluidum, Extract. fʒj. to fʒss.	——, Tinctura, fʒj. to ij.
Spiritus Ætheris Nitrici, fʒj.	Veratrum Album, gr. j. to iij.
Spiritus Ætheris Compositus, fʒj.	Veratri Albi Vinum, gtt. x. to xx.
Spiritus Pyroxilicus, gtt. x. to xl.	Veratrum Viride, gr. j. to iij.
Stanni Pulvis, ʒss.	Veratria, and its salts, gr. $\frac{1}{12}$ to $\frac{1}{8}$.
Stramonium, gr. j. to ij.	
Stramonii Foliorum, Extractum, gr. j.	Zinci Acetas, gr. j. to ij.
—— Seminis, Extractum, gr. $\frac{1}{4}$ to $\frac{1}{2}$.	—— Chloridum, gr. $\frac{1}{4}$ to ij.
—— Tinctura, gtt. xx. to xxx.	—— Cyanuretum, gr. $\frac{1}{12}$ to $\frac{1}{4}$.
Strychnia, and its salts, gr. $\frac{1}{12}$ to $\frac{1}{8}$.	—— Oxidum, gr. ij. to v.
Styrax, gr. x. to xx.	—— Sulphas, gr. j. to x.
Sulphur, ʒj. to ʒj.	—— Valerianas, gr. j. to ij.
Sulphuris Iodidum, gr. j.	Zingiber, gr. x. to xx.
	Zingiberis Tinctura, fʒj.

BOOK I.
OF
INTERNAL REMEDIES.

CLASS I.

EMETICS.

EMETICS are medicines which excite vomiting, independently of any effect arising from the mere quantity of matter introduced into the stomach.

In the administration of Emetics a few simple rules are worthy of attention. Whatever may be the apparent necessity for emptying the stomach, we must take into consideration all the circumstances of the case, and more particularly if there be much arterial excitement with determination to the head; for where these symptoms are present, bloodletting should be sometimes premised.

When it is required, as in cases of poisoning, a full dose of the emetic substance is to be administered; but in the generality of instances, it is better to give it in divided doses, frequently repeated, until the desired effect is produced. In cases of torpor or congestion, it is sometimes requisite to arouse the system by retching or vomiting, which may be done by administering the emetic with only a small quantity of fluid; but when the object is to empty the stomach and duodenum merely, free draughts of tepid water or weak chamomile tea ought to be given as soon as nausea takes place.

When *poisons* have been taken, our object is to evacuate the stomach as speedily as possible, and for this purpose we select an emetic which is prompt in its action. But where the excitability of the stomach is greatly diminished by a *Narcotic*, it is necessary to assist the emetic by the addition of some excitant. When such an effect is caused by opium, the system may be generally rendered susceptible by the use of vinegar, mustard, or ammonia. In these cases, however, the stomach-tube is our best resource.

The evacuation of the stomach may be greatly facilitated by tickling the fauces with a feather, or introducing the point of the finger as far back as possible and titillating the parts. If

this fails, a resort to another article of the same class will often accomplish the object.

There are a few cases in which emetics are obviously inadmissible, as in apoplexy, or where there is great determination of blood to the head, depending upon constitutional causes: pregnancy; hernia; prolapsus uteri; active hemorrhage from the lungs and uterus; acute gastritis, &c. If emetic medicines are ever allowable in these diseases it is for the most part in nauseating doses only.

By the frequent use of emetics the stomach acquires an increased susceptibility to their action; hence persons of delicate habits should use them cautiously.

When excessive vomiting occurs, it may be checked by resorting to some one or more of the following resources:—

Ice, broken into small pieces and allowed to dissolve in the mouth.

Lime-water and milk, a tablespoonful of each, mixed cold and given at intervals of fifteen or twenty minutes.

A drop or even half a drop of creosote, mixed in a wine-glass of cold water, of which give a teaspoonful every 15 or 20 minutes.

The infusion of camphor made with boiling water and given *cold*, a teaspoonful frequently repeated.

Brandy and water, and clove and green tea, conduce to the same end; and where other internal means fail in irritable constitutions, an anodyne injection will sometimes accomplish the object.

Among external appliances, mustard to the epigastrium is one of the best; also, spicc-plasters and hot fomentations of brandy and cloves.

EMETIC POWDERS.

Powder of Ipecacuanha.

R. Pulveris Ipecacuanhæ, ℥ij.

Divide it into two doses. *Signa.*—One to be taken in a little molasses or sugar and water, and followed by a draught of tepid water. If one powder does not produce the desired effect, the second to be taken in like manner.

Powder of Ipecacuanha with Tartarized Antimony.

R. Pulveris Ipecacuanhæ, ℥ss.

Antimonii et Potassæ Tartratis, gr. ij.

Misce, et divide in pulveres ij.—To be taken in the same manner as the former.

Tartar emetic increases the efficacy of ipecacuanha, producing a more prompt and certain effect than the latter drug. This combination is an eligible one in most cases where emetics are indicated. When tartar emetic is given alone, it is generally in solution, a formula for which will be found below.

Powder of Alum.

R. Aluminis, ʒj.
Fiat pulvis.

This is highly recommended by Dr. Meigs in croup. "It is best given in the state of fine powder, mixed with honey, sirup of any kind, or molasses."

The dose is a teaspoonful, to be repeated every fifteen minutes until free emesis is produced. MEIGS.

Turpeth Mineral.

R. Hydrargyri Sulphatis flavi, gr. xij.
Divide in pulveres iv.

Dose, one powder for a child three years old, to be repeated, in croup, every fifteen minutes, until emesis is produced. "If the first dose fails, the second usually acts as soon as it touches the stomach." HUBBARD.

In cases of narcotic poisoning it is necessary to resort to the most prompt and certain means of evacuating the stomach. Ipecacuanha was formerly thought to be a specific for poisoning by opium, but experience has shown that mustard, salt, alum, turpeth mineral, and the sulphates of zinc and copper are more reliable.

Powder of Mustard.

R. Pulveris Sinapis, ʒj.

Signa.—Take a teaspoonful in a cupful of tepid water every ten minutes.

This is a very prompt and safe emetic.

Emetic of Common Salt.

R. Sodii Chloridi, ʒj.

Signa.—Administer one-half in a cupful of warm water, and repeat the dose in ten minutes.

Salt frequently proves a prompt and efficient emetic, invigorating rather than depressing the powers of the system.

Powder of Sulphate of Copper.

R. Cupri Sulphatis, gr. ij.

Signa.—To be given in a little sirup and water.

This medicine is seldom employed as an emetic except in those cases of poisoning where other articles have failed, or are not at hand. It should be followed by a large draught of warm water. All vegetable astringent infusions are incompatible substances.

Powder of Sulphate of Zinc.

R. Zinci Sulphatis, gr. x. vel ʒss.

Signa.—To be taken in molasses, and followed by a draught of warm water.

This medicine is very prompt in its operation and very generally employed in cases of poisoning. It is also used as a nauseating remedy in phthisis; and both as an emetic and tonic in dyspepsia. The stomach is said to be invigorated, rather than exhausted by its operation.

Powder of Ipecacuanha and Calomel.

R. Pulveris Ipecacuanhæ, gr. viij.

Hydrargyri Chloridi Mitis, gr. v.

Misce.

This powder generally proves both emetic and purgative in a child five years old.

Powder of Ipecacuanha with Rhubarb.

R. Pulveris Ipecacuanhæ,

———— Rhei, āā ʒj.

Misce. *Signa.*—To be taken in sirup.

More or less purging will generally follow the emetic operation of this dose.

EMETIC WINES AND TINCTURES.

Wine of Ipecacuanha with Antimony.

R. Vini Ipecacuanhæ, f ʒj.

Antimonii et Potassæ Tartratis, gr. j.

Fiat solutio. *Signa.*—A teaspoonful to be taken every ten or fifteen minutes, until vomiting is induced, which may be assisted with tepid drinks.

Antimonial Wine.

R. Vini Antimonii, f℥j.

Signa.—A teaspoonful to be taken every ten or fifteen minutes, until free emesis is effected.

If antimonial wine is given for the purpose of producing speedy vomiting, we may safely give half an ounce of it at once; and in the croup of infants, twenty drops every fifteen minutes, until the medicine operates, is a safe dose.

Tincture of American Hellebore.

R. Radicis Veratri Viridis, ʒviij.

Alcoholis, Oj.

Macera per dies decem, vel quatuordecim, et exprime.

Signa.—Dose for an adult, six to eight drops every three hours, and to be increased one or two drops each dose, until nausea, vomiting, or a reduction of the frequency of the pulse takes place. Then reduce to one-half in all cases. NORWOOD.

Dr. Norwood describes this article as a good diaphoretic, a prompt and efficient emetic, and an arterial sedative of the first rank. He recommends its use in hooping-cough, croup, inflammation of the lungs, and typhoid fever.—*Southern Med. and Surg. Journal*, Jan. 1853.

Tincture of Lobelia.

R. Tincturæ Lobeliæ, f℥j.

Signa.—Give a teaspoonful every two hours in asthma.

This quantity will produce distressing nausea in other diseases, and sometimes will even cause vomiting.

EMETIC SOLUTIONS AND MIXTURES.

Solution of Tartrate of Antimony.

R. Antimonii et Potassæ Tartratis, gr. iv.

Sacchari, ʒj.

Aquæ destillatæ, f℥iv.

Misce, et fiat solutio.—A tablespoonful to be given every ten or fifteen minutes, until it operates freely.

Solution of Sulphate of Zinc and Alum.

R. Zinci Sulphatis, ʒiss.

Aluminis, ʒss.

Aquæ ferventis, Oss.

Misce.—Let the solution be filtered through blotting-paper.—A

tablespoonful for adults, and a teaspoonful for a child of six months old (and so on in proportion to the strength and age of the patient), every morning upon an empty stomach.

This is the celebrated *solutio vitriolica* of Dr. Mosely, who principally employed it in dysentery and phthisis pulmonalis.

Solution of Emetia.

R. Emetiæ fuscae, gr. iv.

Aquæ destillatæ, f℥ij.

Syrupi, f℥ss.

Fiat solutio. *Signa.*—Give two teaspoonsful every half hour. Emetia is destitute of the peculiar odor and nauseous taste of ipecacuanha, and may therefore be preferable in some cases.

It will be observed that we have introduced the impure or colored emetia, which is much less energetic than the perfectly white alkali. The best antidote for an overdose, is a strong infusion of nutgalls.

Mixture of Ipecacuanha, &c.

R. Pulveris Ipecacuanhæ, ℥ss.

Antimonii et Potassæ Tartratis, gr. j.

Tincturæ Scillæ, f℥j.

Aquæ destillatæ, f℥vij.

Fiat mistura. *Signa.*—Four tablespoonsful at first; afterwards two tablespoonsful every ten minutes until vomiting is induced.

This mixture is recommended in dropsies previously to giving digitalis.

Infusion of Ipecacuanha.

R. Pulveris Ipecacuanhæ, ℥ij.

Aquæ bullientis, f℥vj.

Macerate for one hour and filter.—Of this from one to two tablespoonsful may be given every half hour, until it excites vomiting.

Infusion of Lobelia Inflata, or Indian Tobacco.

R. Lobeliæ, ℥j.

Aquæ bullientis, Oj.

Fiat infusio. *Signa.*—A small wineglassful may be taken every half hour, until vomiting succeeds.

This remedy is too hazardous and distressing for ordinary use; but in asthma, and some spasmodic affections, much bene-

fit is often experienced. It should, however, be administered with extreme caution.

Infusion of Tobacco.

R. Tabaci, ʒj.

Aquæ bullientis, Oj.

Macerate per horam unam in vase leviter clauso, et cola.

This infusion is characterized by the nausea and depression following its administration. It is only used for strangulated hernia, to reduce dislocations, for obstinate colic, and in spasm of the urethra. One-half of the above may be employed at one time as an enema, to be followed by the remaining portion in thirty minutes, if sufficient relaxation is not produced. One or two tablespoofuls by the mouth generally prove emetic.

Sirup of Seneka and Squills.

R. Syrupi Scillæ Compositi, fʒj.

Signa.—Give from ten drops to a teaspoonful, every quarter or half hour, or hour, according to the age of the patient and violence of the disease.

This is the *hive sirup* of Dr. Coxe; it has acquired much confidence from its efficacy in croup; and should it not be sufficiently active in this violent disease, a grain of tartrate of antimony may be added to each ounce.

Sirup of Ipecacuanha.

R. Syrupi Ipecacuanhæ, fʒj.

Signa.—Give a teaspoonful to a child two years old, every fifteen minutes until it proves emetic.

As one fluidounce only represents fifteen grains of the powdered root, this must be considered a weak preparation. The following formula employed in New York, I consider better: When made it affords three pints of sirup, which keeps well without crystallization. A fluidounce of this sirup is equivalent to one dram of powdered ipecacuanha.

Sirup of Ipecacuanha.

“R. Take of Ipecacuanha, bruised, *six ounces* (troy),
Alcohol, *one pint and a half*,
Water, *one pint*.

Mix, to form a tincture. Digest for ten days, filter and add one pint of water by way of displacement, evaporate in a water

bath to two pints, add immediately, of refined sugar, *three pounds and a quarter* (troy) and bring to the boiling point."

COGGESHALL.

The dose as an emetic, and modes of using, are the same as above.

EDITOR.

VARIOUS EMETIC PREPARATIONS.

Tartar Emetic and Quinia.

R. Antimonii et Potassæ Tartratis, gr. iij.

Quiniæ Sulphatis, gr. x.

Misce, et divide in partes vj. equales.—This preparation is recommended by Dr. Gola as successful in intermittent fever, when quinia alone produced no favorable result. He directs a powder to be given every two hours during the apyrexia. It causes both emesis and alvine evacuations, and prevents the recurrence of the paroxysm.

Emetic Lozenges of Emetia.

R. Sacchari, ℥ij.

Emetiæ Impuræ, gr. xxxij.

Misce, et fiant trochisci singuli gr. xvij.

One of these lozenges, taken fasting, will generally produce vomiting in children. Three or four will speedily excite it in adults.

MAGENDIE.

Tobacco Poultice.

R. Tabaci, ℥j.

To be beat up with sufficient water to form a cataplasm, and applied to the region of the stomach, when emetics cannot be administered by the mouth. A dram or two of snuff mixed with lard, or a single tobacco-leaf bruised and softened with hot water and applied over the sternum of a child in croup, usually provokes vomiting. The utmost care should be taken to guard against the violent effects of tobacco, even when thus externally applied.

CLASS II.

CATHARTICS.

As a general rule it may be observed of cathartics that they are capable of fulfilling three different indications: 1. Simply to evacuate the bowels; 2. To excite an increased discharge from the mucous coat of the intestines; and 3. To stimulate the neighboring viscera, and cause them to secrete a greater quantity of their peculiar fluids. It seldom or perhaps never happens that the same article will produce these three different effects; and in order to accomplish them, we must combine different substances in one formula.

There are some, as magnesia, that produce their effect by absorbing the acid generated in the alimentary canal, and thus become converted into a purgative salt. There are others, as aloes, that expend their principal force upon the rectum and great intestines, and for this reason are very advantageously employed in diseases of the lower bowels and of the uterus and its appendages.

These remedies should always be given on an empty stomach; if administered immediately after a full meal, they arrest the digestive process, are liable to produce nausea, if not vomiting, and do not open the bowels with the same certainty or effect. When simple costiveness is to be removed, the evening may be selected for the exhibition of a cathartic.

If this should not prove sufficient, we can repeat the dose in the morning and at regular intervals through the day until the effect is produced. The principle, says Dr. Paris, which should govern our conduct in the administration of purgatives, may be briefly stated: *The excitement is to be repeated till the requisite action is induced, yet no single excitement being such as may prove an irritant to the organ.*

But in the treatment of many diseases, it is preferable to purge through the day, in order that the sleep of the patient may not be disturbed at night.

The operation of these remedies may be very much accelerated by the free exhibition of diluent drinks, such as gruel, barley water, &c. &c.

By combination we are enabled to modify and control the action of many remedies, and some of the most drastic purgatives may be made to operate mildly and yet certainly by uniting several of them in the same dose.

The addition of an emetic substance, such as tartar emetic, or ipecacuanha, in small quantities, gives activity to the combination, while it modifies the harshness of the drastic purgatives.

Finally, the operation of a cathartic may often be promoted by judicious venesection; and if spasm of the intestines be present, opium may be advantageously joined with the purgative.

PURGATIVE POWDERS.

Powder with Calomel and Jalap.

R. Hydrargyri Chloridi Mitis, gr. v.

Pulveris Jalapæ, gr. x.

Misce. *Signa.*—To be given in molasses or sirup.

Powder with Calomel and Rhubarb.

R. Hydrargyri Chloridi Mitis,

Pulveris Rhei, āā gr. v.

Olei Cinnamomi, gtt. j.

Misce. *Signa.*—To be given in molasses.

Powder with Calomel and Soda.

R. Hydrargyri Chloridi Mitis, gr. viij.

Sodæ Bicarbonatis, gr. xij.

Misce. *Signa.*—To be given in molasses.

This combination forms a safe and mild purgative. Nausea seldom follows its administration.

Powder with Calomel, Jalap, and Rhubarb.

R. Hydrargyri Chloridi Mitis, gr. v.

Pulveris Jalapæ,

———— Rhei, āā gr. v.

Olei Cinnamomi, gtt. j.

Misce. *Signa.*—To be given as the former.

Powder with Rhubarb and Magnesia.

R. Pulveris Rhei, ℥j.

Magnesiæ, ℥ss.

Olei Cinnamomi, gtt. j.

Misce. *Signa*.—To be given in sirup, or sugar and water.*Calcined Magnesia.*

R. Magnesiæ, ℥j.

R. Magnesiæ (HENRY), ℥j.

R. Magnesiæ (HUSBAND), ℥j.

R. Magnesiæ (ELLIS), ℥j.

Magnesia is an antacid, and laxative, much used in the diseases of women and children. The lightness and consequent great bulk of the officinal powder, constitute serious objections to its use. To obviate these, Henry of England, and Thos. J. Husband, and Chas. Ellis and Co. of this city, have severally prepared a heavy magnesia as a substitute. Henry's magnesia has long been favorably known; Husband's and Ellis's are entitled to equal consideration with the imported article, since the experiments of Prof. Procter have shown that, for forming liquid mixtures, and for rapidity of antacid action, they are equal if not preferable to it.

Dose of either, from thirty grains to a dram, in milk or water. The operation of magnesia, when it meets with no acid in the bowels, is promoted by giving the patient lemonade.

Powder of Jalap and Cream of Tartar.

R. Pulveris Jalapæ, ℥j.

Potassæ Bitartratis, ℥vj.

Misce, et divide in chartulas vj.

Signa.—Give one powder every three hours in molasses. This powder is a hydragogue cathartic. "It constitutes one of the best medicines which I have ever tried, as well in dropsical as in all other cases, where long-continued purging is demanded."

CHAPMAN.

Compound Powder of Jalap and Gamboge.

R. Pulveris Jalapæ Compositi, ℥iij.

——— Gambogiæ, gr. vj.

Misce, et divide in chartulas vj.

Signa.—Give one powder every two hours in molasses or sirup.

The compound powder of jalap has long been celebrated in dropsical affections; but it also forms a very convenient purge in many inflammatory cases. In order to increase its certainty and activity, we may conveniently add the Gamboge, as in the above formula.

Powder of Sulphur and Cream of Tartar.

R. Sulphuris loti, ʒss.

Potassæ Bitartratis, ʒj.

Misce. *Signa.*—Mix in a sufficient quantity of molasses, and give a teaspoonful three or four times a day to children. In some diseases of the skin.

It should be always recollected that sulphur promotes insensible perspiration in a very marked degree, and hence renders the patient more liable to be affected by vicissitudes of climate.

Powder with Milk of Sulphur and Calcined Magnesia.

R. Sulphuris præcipitati,

Magnesiae, āā ʒss.

Misce. *Signa.*—A teaspoonful to be taken three or four times a day.

This preparation is better adapted to delicate stomachs than the former. Either of them forms a good aperient during an alterative course of medicine.

CHAPMAN.

Seidlitz Powders.

R. Sodæ et Potassæ Tartratis, ʒij.

Sodæ Bicarbonatis, ʒij.

Fiat pulvis. *Signa.* To be dissolved in half a tumbler of cold water.

R. Acidi Tartarici pulverizati, gr. xxv.

Signa.—Dissolve in an equal quantity of water with the above, and mix them both together. To be taken while in a state of effervescence.

This is a very popular medicine, and justly so; it is a gentle aperient, and may be taken by persons whose stomachs would reject any of the ordinary saline preparations.

Soda Powders.

R. Sodæ Bicarbonatis, ʒss.

Signa.—To be dissolved in a wineglassful of cold water.

R. Acidi Tartarici, gr. xxv.

Signa.—Dissolve in the same quantity of sweetened water, flavored with two drops of essence of lemon.

Mix the two solutions, and drink in a state of effervescence. These effervescing drinks, in addition to their laxative effects, allay thirst, check sickness of stomach, and diminish febrile heat.

Stevens's Saline Powders.

R. Sodæ Bicarbonatis, ℥ss.

Sodii Chloridi, ʒj.

Potassæ Chloratis, gr. vij.

Misce.

This powder was administered by Dr. Stevens in half a tumbler of tepid water, or a little thin beef tea, in cholera, and repeated every half hour until the patient recovered from the collapse.

PURGATIVE PILLS.

Griffitts's Cathartic Pills.

R. Pulveris Jalapæ,

———— Rhei,

Saponis, āā ℥ss.

Hydrargyri Chloridi Mitis, gr. xxv.

Antimonii et Potassæ Tartratis, gr. jss.

Aquæ destillatæ, quantum sufficit ut fiat massa. Divide in pilulas xxv.

Signa.—Two of these pills may be taken at once, and repeated in two hours, if not sufficient.

This combination of purgatives with the tartrate of antimony, forms an excellent antibilious remedy; and, in cases of fever, prepares the alimentary canal for the reception of other remedies.

Pills of Blue Mass, Aloes, &c.

R. Pilulæ Hydrargyri,

Pulveris Jalapæ,

———— Aloës, āā gr. xv.

Syrupi, q. s. ut fiat massa. Divide in pilulas xij.

Signa.—Three may be taken at night, and the same dose repeated in the morning, if the first is ineffectual.

Pills of Blue Mass, Soda, &c.

R. Pilulæ Hydrargyri, gr. ix.

Pulveris Rhei,

Sodæ Bicarbonatis, āā gr. xij.

Syrupi Rhei Aromatici, q. s. ut fiat massa, in pilulas xij. dividenda.

Signa.—One, twice or three times a day, as an alterative, in derangement of the liver. DR. HARTSHORNE.

Pills of Blue Mass, and Colocynth.

R. Pilulæ Hydrargyri,

Extracti Colocynthidis Compositi, āā gr. v.

Olei Cari, gtt. ij.

Misce, et fiant pilulæ ij.

These constitute an active cathartic.

Pills of Calomel and Colocynth.

R. Extracti Colocynthidis Compositi, gr. xlvij.

Hydrargyri Chloridi Mitis, ʒj.

Misce, et divide in pilulas xx.

Signa.—Two or three of these pills will generally act as a cathartic.

Pills of Extract of May-Apple and Podophyllin.

R. Extracti Podophylli, ʒss.

Podophyllin (*Merrill, Parrish*), gr. x.

Misce, et fiat massa in pilulas x. dividenda.

Signa.—Give one pill every three hours.

Podophyllin is a composite principle, lately reintroduced into notice by W. S. Merrill, of Cincinnati, and Edward Parrish, of Philadelphia. It is spoken of as a chologogue cathartic. It certainly does operate promptly, and sometimes harshly on the bowels, in the dose of one or two grains. EDITOR.

Anti-Gout Pill.

R. Extracti Colchici Acetici,

———— Colocynthidis Compositi, āā gr. x.

Morphiæ Muriatis, gr. ijss.

Misce, et fiat massa in pilulas x. dividenda.

This is a useful anti-gout pill; one adapted to mitigate an actual paroxysm. One pill should be given every two hours, until the bowels and kidneys are acted on. ROBERT DICK.

Pills with Extract of Butternut and Jalap.

R. Extracti Juglandis, ʒss.

Pulveris Jalapæ, ʒj.

Saponis, gr. x.

Misce, et fiant pilulæ xv.

Signa.—Three or four of these may be taken for a dose, and if they do not operate, two or more may be safely administered.

The extract of butternut, either alone or in combination, is highly recommended as a cathartic in fevers, dysentery, &c.

Pills of Compound Extract of Colocynth, &c., commonly called Fothergill's Pills.

R. Extracti Colocynthidis Compositi, ʒiss.

Antimonii Oxidi, *Ed.* ʒss.

Misce, et divide in pilulas xxx.—One of these pills will generally produce a free evacuation; but two or more may be taken when required.

Pills of Compound Extract of Colocynth, &c.

R. Extracti Colocynthidis Compositi, ʒj.

Resinæ Jalapæ, *Ed.* gr. vj.

Pulveris Scammonii Compositi, *Lond.* gr. x.

Hydrargyri Chloridi Mitis, gr. x.

Antimonii et Potassæ Tartratis, gr. j.

Saponis, gr. v.

Olei Cinnamomi, gtt. iv.

Misce, et divide in pilulas xv.

Signa.—One, two, or three to be taken at bedtime.

DR. MEIGS.

Mitchell's Pills.

R. Pulveris Rhei, ʒiv.

—— Aloës, ʒij.

Hydrargyri Chloridi Mitis, gr. iv.

Antimonii et Potassæ Tartratis, gr. ij.

Fiat pulvis, et adde—

Saponis, in pulverem, ʒij.

Aquæ, q. s. ut fiat massa. Divide in pilulas xl.

Signa.—Take one pill every night as a laxative, or three pills as a purgative.

J. K. MITCHELL.

Pills of Elaterium.

R. Elaterii, gr. ij.

Extracti Gentianæ, ʒss.

Misce, et divide in pilulas xvj.

Signa.—Give one pill every three hours, until free purging takes place. Elaterin, the active principle of this drug, can now be procured in a state of purity; and as it is more certain in its effects, a formula is subjoined under the head of purgative wines and tinctures.

Pills of Croton Oil.

R. Olei Tiglii, gtt. ij.

Micæ Panis, q. s. ut fiant pilulæ iv.

Signa.—One every hour until they operate.

This powerful remedy has lately been restored to the materia medica. One drop is a full dose, and frequently less will answer every purpose.

The following is a more certain and less drastic mode of exhibiting Croton Oil.

Pills of Croton Oil and Colocynth.

R. Olei Tiglii, gtt. ij.

Extracti Colocynthidis Compositi, gr. xij.

Misce, et divide in pilulas iv.

Signa.—Give one pill every two hours.

Another Form.

R. Extracti Colocynthidis Compositi, gr. l.

Saponis, gr. x.

Olei Tiglii, gtt. ij.

Misce, fiat massa in pilulas, xij. dividenda.

Signa.—Take one pill occasionally.

BRODIE.

Antibilious Pills.

R. Pilulæ Catharticæ Compositæ, No. iij.

One pill often acts as a laxative, and three as a purgative. Each pill contains one grain of calomel.

Peristaltic Persuaders.

R. Pulveris Rhei, ʒj.

———— Ipecacuanhæ, gr. x.

Olei Cari, gtt. x.

Pulveris Acaciæ, q. s.

Fiat massa, et divide in pilulas xx.

Signa.—Take two pills at bedtime, in obstinate constipation.

Dr. Chapman also highly recommends the use of ten drops of the wine of colchicum root, every four hours, in the same complaint.

CHAPMAN.

VARIOUS ALOETIC PILLS.

Pills of Aloes with Soap.

R. Pilulæ Aloës, No. v.

Each pill contains two grains of aloes.

Pills of Aloes and Assafetida.

R. Pilulæ Aloës et Assafœtidæ, No. iij.

Each pill contains one and a third grains of the respective ingredients.

Rufus's Pills.

R. Pilulæ Aloës et Myrrhæ, No. iij.

Each pill contains two grains of aloes, one of myrrh, and half a grain of saffron.

Pills of Rhubarb, Aloes, and Myrrh.

R. Pilulæ Rhei Compositæ, No. iij.

Each pill contains two grains of rhubarb, one and a half grains of aloes, and one grain of myrrh.

Aperient Pills.

R. Pulveris Aloës,

———— Rhei, āā ʒj.

———— Ipecacuanhæ, gr. vj.

Saponis, ʒj.

Aquæ, q. s.

Fiant pilulæ lx.

As a simple aperient, one may be taken every night at bed-time; but where a full purgative effect is desirable, the dose is two or three.

Pills with Aloes, Gentian, &c.

R. Pulveris Aloës, ʒj.

Extracti Gentianæ, ʒss.

Olei Cari, gtt. x.

Syrupi, q. s. ut fiat massa, in pilulas singulas grana quatuor pendentes, dividenda.

Signa.—Take two or three pills for a dose.

Pills with Aloes, Guaiacum, &c.

R. Pulveris Guaiaci, ʒj.

———— Aloës, gr. xxxvj.

———— Rhei, ʒij.

Terebinthinæ Canadensis, quantum sufficit ut fiat massa in pilulas xlvij. dividenda.

Signa.—One to be taken every three hours.

This combination forms a highly stimulating cathartic.

Hooper's, Anderson's, Webster's, and Morrison's Pills, which enjoy so high a reputation with the public, depend in a great measure for their efficacy on the presence of aloes.

Pills of Rhubarb and Sulphate of Iron.

R. Pulveris Rhei, ʒiss.

Ferri Sulphatis, ʒss.

Saponis, ʒij.

Aquæ destillatæ, q. s. ut fiat massa in pilulas xl. dividenda.

The intention of these pills is obviously to remove costiveness, and impart tone to the bowels. They were a favorite medicine with Dr. Griffiths, late of this city. Three or four may be taken for a dose at bedtime.

Pills of Ox Gall.

R. Fellis Bovini inspissati, ʒij.

Olei Cari, gtt. x.

Magnesiae Carbonatis, q. s. ut fiat massa, in pilulas xxxvj. dividenda.

Signa.—Take two pills three times daily.

CLAY.

Ox gall has again been brought forward as a remedy in dyspepsia and obstinate constipation. Its advocates claim for it certainty of action, attended with little pain or griping.

PURGATIVE CONFECTIONS AND ELECTUARIES.

Lenitive Electuary.

R. Confectionis Sennæ, ʒj.

Signa.—Take a piece the size of a marble, night and morning, in costiveness.

Confection of Sulphur.

- R. Sulphuris loti, ʒiss.
 Confectionis Sennæ, ʒij.
 Potassæ Nitratis, ʒj.
 Syrupi Aurantii Corticis, q. s. ut fiat confectio.

Signa.—One or two drams may be taken once or twice a day.
 In hemorrhoidal affections.

Electuary of Senna and Figs.

- R. Pulveris Sennæ, ʒss.
 Pulpæ Ficûs, quantum sufficit ut fiat electuarium.

Signa.—A piece the size of a marble, night and morning, in costiveness.

Electuary with Jalap, Nitrate of Potash, &c.

- R. Pulveris Jalapæ,
 Potassæ Bitartratis,
 ——— Nitratis, āā ʒss.
 Confectionis Sennæ, ʒj.
 Syrupi, q. s. ut fiat electuarium. Of which a bolus the size of a hazel-nut may be taken three or four times a day.

This preparation once gained considerable celebrity as a popular remedy for hemorrhoids.

Confection of Sulphur, Guaiac, &c.

- R. Sulphuris, ʒij.
 Potassæ Bitartratis, ʒj.
 Pulveris Guaiaci, ʒj.
 ——— Rhei, ʒij.
 Spiritûs Myristicæ, fʒij.
 Mellis, quantum sufficit, ut fiat electuarium.

Signa.—Take one or two teaspoonfuls morning and evening.

This preparation has long been used successfully in England in the treatment of chronic rheumatism.

PURGATIVE EMULSIONS.

Oleaginous Mixture.

- R. Pulveris Acaciæ, ʒij.
 Aquæ, fʒij.
 Olei Ricini, ʒj.
 Aquæ Cinnamomi, vel
 ——— Menthæ Viridis, fʒj.

Misce secundum artem.

Signa.—Take a tablespoonful every two hours.

Sugar is generally prescribed, but the mixture remains more uniform without it. The compounder should mix the gum arabic with sufficient water to form a thick mucilage, and then rub in small portions of the oil and water until the whole is suspended. Finally, add the cinnamon or mint-water, and shake well together.

Another Form for Children.

R. Olei Ricini, ʒij. vel ʒss.

Vitellum Unius Ovi,

Aquæ Fœniculi,

——— Menthæ Viridis, āā fʒj.

Fiat emulsio. *Signa*.—Give a teaspoonful morning and night to a child three months old.

Another Form.

R. Mucilaginis Acaciæ, fʒj.

Olei Ricini, ʒj.

Aquæ Menthæ Viridis, fʒij.

Mix the oil and mucilage, add the mint-water, and shake well together.

Signa.—Take a tablespoonful every two hours.

Mixture of Oil of Turpentine.

R. Olei Terebinthinæ, fʒj.

Pulveris Acaciæ, ʒiij.

Aquæ Menthæ Viridis, fʒiv.

Misce. *Signa*.—A tablespoonful every two hours until it operates.

This preparation must be made in the same manner as the oleaginous mixture.

Mixture of Oil of Turpentine.

R. Olei Terebinthinæ, fʒj.

Vitellum Unius Ovi,

Tere simul, et adde gradatim—

Misturæ Amygdalæ, fʒiv.

Syrupi Aurantii Corticis, fʒij.

Spiritûs Lavandulæ Compositi, fʒjss.

Olei Cinnamomi, gtt. iv.

Misce, et fiat emulsio. *Signa*.—Take a tablespoonful every two hours.

Oleaginous Mixture with Oil of Turpentine.

- R. Olei Ricini, f̄iss.
 Vitellum Ovi, j.
 Sacchari, ʒij.
 Olei Terebinthinæ, f̄ss.
 Aquæ Menthæ Viridis, f̄iv.

Misce optime. *Signa.*—A large tablespoonful may be taken every hour or two, according to circumstances.

This preparation is much used by some of our best practitioners in cases requiring a brisk and stimulating cathartic. It has also been recommended as a purgative after the effects of an overdose of opium have been partially subdued by preliminary remedies.

Mixture of Croton Oil.

- R. Olei Tiglii, gtt. j. vel ij.
 Mucilaginis Acaciæ,
 Aquæ destillatæ, āā f̄ʒj.

Misce. *Signa.*—To take a teaspoonful every two hours until it operates.

Saponaceous Solution of Croton Oil.

- R. Olei Tiglii, gtt. viij.
 Potassæ, gr. vj.
 Aquæ destillatæ, f̄ʒij.

Fiat solutio. *Signa.*—From three to six drops may be given for a dose.

This is a substitute for the soap of croton oil proposed by the French, which is more troublesome to make.

Mr. Conwel states that four drops of this oil rubbed on and round the umbilicus, will produce a cathartic effect; even the odor of it has been known to excite the alimentary canal, either as an emetic or purgative.

P U R G A T I V E M I X T U R E S.

Solution of Sulphate of Magnesia, and Tartar Emetic.

- R. Magnesiæ Sulphatis, ʒj.
 Antimonii et Potassæ Tartratis, gr. ss.
 Aquæ destillatæ, f̄ʒiij.
 Succī Limonis recentis, f̄ʒj.

Misce, et fiat solutio.—A tablespoonful to be given every two hours until it operates. Or one-half may be taken as a purgative dose.

Solution of Epsom Salts with Nitric Acid.

R. Magnesiae Sulphatis, ʒss.

Tincturae Jalapae, fʒj.

Acidi Nitrici, gtt. ij.

Aquae Menthae Viridis, fʒij.

Misce. *Signa.*—To be taken for a draught. The nitric acid is said to destroy in a great measure the bitter taste of the salt.

Mixture of Carbonate of Magnesia, &c.

R. Magnesiae Carbonatis, ʒss.

———— Sulphatis, ʒij.

Spiritûs Ammoniae Aromatici, fʒj.

Tincturae Rhei, fʒss.

———— Ilyocyami, fʒss.

Aquae Menthae Viridis, fʒiv.

Fiat mistura. *Signa.*—A large tablespoonful to be taken two or three times a day. DR. MEIGS.

Mixture of Magnesia and Rhubarb.

R. Magnesiae, ʒss.

Pulveris Rhei, gr. ij.

Sacchari, ʒj.

Tincturae Olei Menthae Piperitae, gtt. vj.

Aquae destillatae, fʒiss.

Fiat mistura. *Signa.*—A teaspoonful to be given every two hours to children, until it operates.

This preparation is designed for the bowel complaints of children, in which it is very serviceable. Laudanum can be added according to the age and condition of the patient.

Magnesia and Blue Pill.

R. Magnesiae, ʒiss.

Acaciae,

Sacchari, āā q. s.

Aquae, fʒiv.

Pilulae Hydrargyri, gr. iv. vel viij.

Olei Anisi, gtt. iv.

Misce. *Signa.*—To take a tablespoonful every two hours until it operates freely. An alterative cathartic.

Magnesia and Colchicum.

R. Magnesiae, ʒj.

Sacchari,

Acaciae, āā q. s.

Aquæ destillatæ, f ʒiv.

Vini Colchici Radicis, gtt. xl.

Fiat mistura. *Signa.*—To take a tablespoonful every two hours until it operates.

This preparation is much used in gouty and rheumatic affections. Laudanum can be added if requisite.

Scudamore's Mixture of Colchicum and Epsom Salts.

R. Magnesiae, ʒiv.

Magnesiae Sulphatis, ʒj.

Aquæ Menthæ Viridis, f ʒv.

Aceti Colchici, f ʒss.

Syrupi, f ʒj.

Fiat mistura.—Dose, one, two, or three tablespoonfuls, repeated every two hours, until four or six evacuations are produced in twenty-four hours. Chiefly used in gouty paroxysms.

Solution of Sulphate of Veratria.

R. Veratriæ Sulphatis, gr. j.

Aquæ destillatæ, f ʒij.

Fiat solutio. Dose, a dessertspoonful. This solution is said to be a good substitute for the Eau Medicinale of Husson, in gouty affections. MAGENDIE.

Solution of Citrate of Magnesia.

R. Liquoris Magnesiae Citratis, f ʒxij.

Signa.—Take one-half at a draught, and repeat in two hours, if necessary.

From its resemblance to lemonade in taste, this constitutes one of the most pleasant laxatives now in use. A child five years old, may take a wineglassful as a dose.

The country practitioner can make a good approximation by neutralizing carbonate of magnesia with citric acid, in the proportion of fourteen parts of the former to twenty parts of the latter, and flavoring the solution with lemon sirup.

Half an ounce of the neutral salt of citrate of magnesia constitutes a dose.

Fluid Extract of Senna.

R. Extracti Sennæ Fluidi, f ʒj.

Signa.—Give a tablespoonful.

CHARLES ELLIS.

The fluid extracts, of which a few are officinal, are intended to combine strength, smallness of dose, facility of administration, and permanency, in one preparation. They have become quite popular with the profession in most of the large cities, and hence there is an obvious propriety in making them officinal.

The fluid extract of senna is a safe, and tolerably certain purgative.

Fluid Extract of Rhubarb.

R. Extracti Rhei Fluidi, f ʒj.

Signa.—Give half a teaspoonful night and morning.

One fluidram of the extract represents one dram of rhubarb.

Sulphate of Potash and Jalap Mixture.

R. Potassæ Sulphatis, ʒij.

Aquæ, f ʒvss.

Tincturæ Jalapæ, f ʒss.

Fiat solutio. Signa.—Take a tablespoonful every two hours; a mild purgative, and deobstruent.

PURGATIVE INFUSIONS AND DECOCTIONS.

Infusion of Senna, Sulphate of Magnesia, &c.

R. Sennæ, ʒss.

Mannæ optimæ,

Magnesiæ Sulphatis, āā ʒj.

Cardamomi, ʒij.

Misce. Signa.—To be infused in a pint of boiling water until cold, and half a teacupful given every hour until it operates.

Infusion and Tincture of Senna, Salts, &c.

R. Infusi Sennæ, f ʒvj.

Tincturæ Sennæ, f ʒij.

Mannæ, ʒij.

Magnesiæ Sulphatis, ʒss.

Aquæ Menthæ Viridis,

— destillatæ, āā f ʒiss.

Misce. Signa.—One-half to be taken for a dose, and repeated if it does not operate. This was a very common prescription of the late Mr. Abernethy, of London.

Infusion of Senna and Manna.

R. Sennæ, ʒvj.
 Mannæ, ʒj.
 Cardamomi, ʒij.

Misce. *Signa*.—To be infused in a pint of boiling water until cold; one-half to be taken at once; and if insufficient, the remainder in two or three hours.

This infusion will keep longer if a little tincture of senna be added to it.

Decoction of Prunes and Senna.

R. Pruni, ʒiij.
 Infusi Sennæ, Oij.
 Decoque ad libras duas.

In making this decoction, the coriander seed directed in the officinal infusion of senna, had better be omitted until near the close of the process. This forms a good laxative for pregnant women.

Infusion of Senna with Rhubarb, &c.

R. Sennæ, ʒvj.
 Mannæ, ʒj.
 Rhei contusi, ʒij.
 Cardamomi, ʒij.

Misce. *Signa*.—To be infused in a pint of boiling water for one hour. Half a teacupful given every hour till it operates.

Decoction of Aloes, &c.

R. Extracti Glycyrrhizæ, ʒss.
 Potassæ Carbonatis, ʒij.
 Aloës,
 Myrrhæ contritæ, } āā ʒj.
 Croci,
 Tincturæ Cardamomi Compositæ, f ʒiv.
 Aquæ destillatæ, Oj.

Boil down all the ingredients, except the last, with the water to twelve fluidounces, and strain; then add the compound tincture of cardamom. The dose of this decoction is from half an ounce to an ounce and a half. It is a mild, warm cathartic.

PURGATIVE WINES AND TINCTURES.

Wine of Colchicum Root.

R. Vini Colchici Radicis, f ʒss.

Signa.—Ten or twenty drops of this may be taken in sugar and water, morning, noon, and night. The preparations of colchicum not unfrequently produce violent vomiting, and require to be continued with great caution. Used in rheumatism and gout.

There are two vinous preparations of colchicum, the wine of the root or corm, which is a saturated one given as above, and the wine of the seeds, which is only about half as strong, and may be given in the dose of a teaspoonful. The two should be carefully designated in prescriptions by their respective terminations, "Radicis," and "Seminis."

Tincture of Veratria.

R. Veratriæ, gr. iv.

Alcoholis, f ʒj.

Fiat tinctura.—This may be given in doses of ten, fifteen, or twenty drops, in a wineglass of water. It is designed as a substitute for the Vinous Tincture of Colchicum, in dropsies, gout, rheumatism, &c.

The Tincture of Veratria is recommended as an external application in the diseases above mentioned. For this purpose also the ointment is proposed.

Veratria is extremely poisonous, and, therefore, to be taken with great caution.

Tincture of Elaterin.

R. Elaterin, gr. j.

Alcoholis, f ʒj.

Solve, dein adde—

Acidi Nitrici, gtt. iv.

Misce. *Signa.*—Dose, twenty to forty drops in cinnamon water.
MORRIS.

Tincture of Croton Oil.

R. Olei Tiglii, gtt. iv.

Tincturæ Myrrhæ, f ʒj.

Misce, digere, et cola.

Signa.—Dose, one teaspoonful.

BATEMAN.

Tincture of Aloes, &c.

R. Pulveris Aloës,

———— Anisi, āā ʒij.

Spiritus Vini Gallici, Oij.

Fiat tinctura.—Dose, from one-third to half a wineglassful, three times a day.

The above is highly spoken of by Dr. R. Pindall in hemorrhoidal affections. He considers it almost a specific, notwithstanding the prejudice against aloes in these affections of the rectum.

ENEMATA.

Enemata, used either as auxiliaries to, or substitutes for cathartics, very properly succeed that class of remedies.

They are employed to fulfil other indications, and the different preparations will be noticed under their appropriate heads. It is usual to direct three times the quantity of the medicine to be given for an enema that is required for a dose by the mouth. For the purpose of procuring a discharge from the bowels, this rule may generally be observed; but in some of the powerful remedies occasionally administered, as opium and tobacco, it might be attended with serious consequences. When using these active drugs, it is better to give only double the quantity.

The bulk of the enema also varies according to the object we have in view. If purgation is to be effected, the subjoined proportions afford a good rule; but if the remedial agent is to be retained for some time, a small amount is obviously indicated, lest the mere bulk of the injection cause an immediate discharge.

The following are the proportions recommended by Professor Chapman in the ordinary enemata for persons of different ages: A child between the ages of one and five years, from fʒiv. to fʒvj. A youth of ten or fifteen years, half a pint; and an adult, from a pint to a quart, according to circumstances.

In all cases, a warm folded towel should be firmly pressed against the anus for fifteen minutes after the injection is thrown up.

Common Enema.

R. Olei Ricini, vel Olivæ, f ʒij.

Sacchari Fæcis, f ʒij.

Sodii Chloridi, ʒi. vel ʒij.

¹ Infusi Lini tepidi, Oj.

Fiat enema.—One-half to be administered at once, and the remainder in half an hour or an hour if the first produces no evacuation. The chloride of sodium is an active ingredient. In fact, there is no better enema than simple salt and water at the ordinary temperature; and sea-water fresh from the ocean answers the same purpose.

Soap Enema.

R. Saponis Vulgaris incisi, vel

——— Mollis, Lond. ʒj.

Infusi Lini tepidi, Oj.

Misce pro enemate.

This is an effectual clyster in cases of flatulence and impacted feces.

Ox Gall Enema.

R. Fellis Bovini spissati, ʒj.—ij.

Vel

Fellis Bovini recentis, f ʒj.

Aquæ tepidæ, Oj.

Misce. *Signa.*—Use for an injection, in cases of obstinate constipation and hardened feces.

Enema with Sulphate of Magnesia, &c.

R. Magnesiae vel Sodæ Sulphatis, ʒij.

Olei Olivæ, f ʒj.

Aquæ tepidæ, Oj.

Fiat enema. *Signa.*—To be used as an injection.

Enema of Colocynth and Manna.

R. Colocynthis Medullæ incisæ, ʒj.

Mannæ, ʒj.

Aquæ bullientis, f ʒx.

Sodii Chloridi, ʒss.

Boil the Colocynth and Manna in the water for ten or fifteen

¹ The officinal infusion of flaxseed is a compound one, containing liquorice root, and is chiefly intended for affections of the lungs and urinary passages. But the article alluded to in this formula, and elsewhere throughout the book, is the common flaxseed tea, made by infusing an ounce of the seeds in a pint of boiling water.

minutes and strain; then add the salt. Employed in obstinate constipation.

Enema of Oil of Turpentine.

- R. Olei Terebinthinæ, fʒjss.
 Vitellum Unius Ovi,
 Infusi Lini tepidi, Oj.

Misce.

Venice turpentine may be substituted for the oil directed in this prescription; and either of them, if required, may be given in larger quantities.

Enema of Infusion of Senna and Epsom Salts.

- R. Infusi Sennæ, Oj.
 Magnesiæ Sulphatis, ʒij.

Misce.—One-half to be exhibited as an enema, and the remainder if necessary.

Enema with Tartrate of Antimony.

- R. Antimonii et Potassæ Tartratis, ʒj.
 Mucilaginis Acaciæ, fʒiv.
 Aquæ tepidæ, fʒxij.

Misce.—Exhibit as the former.

Dr. Chapman recommends this preparation as one admirably calculated to overcome constipation which resists the ordinary remedies. It is very powerful, and requires great care in its management.

SUPPOSITORIES.

(From *Supponere*, to put under.)

Suppositories constitute a class of remedies that has hitherto been little used by American practitioners. They are solid, round, cylindrical, or conical in form, and intended for introduction into the anus. They are prepared by mingling sedative, astringent, or purgative medicines with cocoa-butter, suet, soap, or thick honey.

When made with suet, or cocoa-butter, these substances should be melted at a gentle heat, with an eighth or a twelfth of white wax, according to the season of the year. The medicinal agent is added, and the fluid mass poured into paper cones, to cool and become solid. A suppository should weigh one or two drams, and be as large as the end of the finger or thumb.

Purgative suppositories are often much larger. By the heat of the body, the suet or cocoa-butter is softened, and the remedial agent is thus permitted to develop its effects.

Soap suppositories are formed by cutting soap into convenient shapes. Suppositories with honey are prepared by boiling down this substance, till it becomes sufficiently thick to retain its shape when cool.

Laxative Suppository.

R. Saponis, \mathfrak{z} ij.
 Sodii Chloridi, \mathfrak{z} j.
 Mellis spissati, q. s.
 Misce.

Form into conical suppositories, weighing \mathfrak{z} ss., and oil them on the surface.

Suppository of Rhubarb.

R. Saponis, \mathfrak{z} ijj.
 Extracti Rhei, \mathfrak{z} ss.
 Pulveris Rhei, q. s.
 Misce, et fiant suppositoria iij.
Signa.—Use one suppository at bedtime.

Purgative Suppository.

R. Pulveris Colocynthis, \mathfrak{z} ss.
 Sodii Chloridi, \mathfrak{z} j.
 Mellis spissati, \mathfrak{z} j. vel q. s.
 Fiat suppositorium.

A large suppository of this kind, should be of a cylindrical form like a candle, to facilitate its introduction.

Formulæ, without number, of a purgative character, might be given, but the above are sufficient to show the usual modes of preparing them in France. Astringent, anthelmintic, and anodyne suppositories, are mentioned in other parts of this work, under their respective heads.

CLASS III.

DIURETICS.

THE rules for the administration of this class of remedies are few and simple.

Where there is much arterial excitement, a judicious use of the lancet should precede the employment of diuretics, and the patient be kept cool to avoid perspiration.

In order to insure the full effect of diuretics, diluent drinks should be freely given, and especially those containing some diuretic medicine.

DIURETIC POWDERS.

Powder of Squill and Nitre.

R. Pulveris Scillæ, gr. vj.

—— Potassæ Nitratis, ʒj.

Fiant pulveres vj. *Signa.*—One to be given thrice a day in sugar and water.

Some aromatic, such as cinnamon, may be added to this preparation if the stomach is irritable.

Powder of Uva Ursi and Soda.

R. Pulveris Uvæ Ursi, ʒiss.

Sodæ Bicarbonatis, ʒj.

Misce, divide in chartulas xij. *Signa.*—One three times a day in sugar and water.

In nephritic complaints and affections of the bladder.

Powder of Ergot and Cubebs.

- R. Pulveris Ergotæ, ℥ij.
 ———— Cubebæ, ʒj.
 ———— Cinnamomi, ʒss.
 Sacchari, ʒj.

Divide into eight powders, of which one may be given three or four times a day. In leucorrhœa and gleet. DR. RYAN.

Powder of Squill, Cream of Tartar, &c.

- R. Antimonii et Potassæ Tartratis, gr. ij.
 Pulveris Scillæ, ʒij.
 Potassæ Bitartratis, ʒjss.
 ———— Sulphatis, ʒss.

Fiat pulvis, et divide in partes xx. equales.

Signa.—Take one powder four or five times daily.

This combination produces copious watery stools, together with a considerable flow of urine, properties that render it valuable in cases of general dropsy. EBERLE.

On theoretical grounds, the quantity of squill in this formula appears too large to begin with, each dose containing six grains.

EDITOR.

DIURETIC PILLS.

Pills of Squill and Calomel.

- R. Hydrargyri Chloridi Mitis,
 Pulveris Scillæ, āā gr. xij.

Confectionis Rosæ, q. s. ut fiant pilulæ xij. *Signa.*—One night and morning, in dropsies depending upon visceral derangement.

This preparation is one of the most celebrated of the diuretics, especially when disease of the liver is the cause of effusion. When the calomel affects the mouth it should be omitted, and the squill combined with some other diuretic.

Pills of Digitalis, Calomel, and Opium.

- R. Pulveris Digitalis, gr. xij.
 Hydrargyri Chloridi Mitis, gr. vj.
 Pulveris Opii, gr. iv.
 Confectionis Rosæ, q. s.

Ut fiant pilulæ xij. *Signa.*—One to be taken every eight hours.

In hydrothorax and ascites.

Pills of Calomel, Digitalis, &c.

R. Pulveris Scillæ exsiccatae, gr. iv.

———— Myrrhæ, ʒj.

———— Digitalis, gr. x.

Hydrargyri Chloridi Mitis, gr. vj.

Simul tere et adde—

Assafoetidæ, ʒss.

Extracti Gentianæ, q. s.

Ut fiat massa, in pilulas xv. dividenda. *Signa.*—One night and morning. DR. PARIS.

Pills of Copaiba.

R. Pilulæ Copaibæ, No. xx.

Signa.—Take two pills for a dose, four times daily.

Each pill contains about five grains of copaiba.

Turpentine Pills.

R. Terebinthinæ, ʒj.

Divide in pilulas xv. *Signa.*—One four or five times a day.

Pills of Extract of Dandelion and Blue Mass.

R. Extracti Taraxaci, ʒss.

Pilulæ Hydrargyri, gr. x.

Pulveris Uvæ Ursi, q. s.

Ut fiant pilulæ x. *Signa.*—One to be taken morning, noon, and night. In dropsy complicated with disease of the liver.

Pills of Digitalis, Squill, and Juniper.

R. Pulveris Digitalis, ʒj.

———— Scillæ, ʒss.

Extracti Gentianæ, ʒj.

Olei Juniperi, gtt. viij.

Syrupi, q. s.

Ut fiat massa, in pilulas xij. dividenda.

Signa.—Take one pill three times daily.

Used in dropsy.

PEARSON.

Pills of Digitalis, Squill, and Blue Mass.

R. Pulveris Digitalis, gr. v.

———— Scillæ, gr. x.

Pilulæ Hydrargyri, ʒss.

Syrupi Acaciæ, q. s.

Fiat massa, et divide in pilulas x.

Signa.—Take one pill three times daily.

This combination was highly recommended by the late Dr. Baillie of London, as an alterative and diuretic in anasarca.

BAILLIE.

DIURETIC INFUSIONS AND DECOCTIONS.

Infusion of Digitalis.

R. Digitalis, ʒj.
Aquæ bullientis, Oss.

Macerate for three hours, then strain.—Dose, for an adult, half a tablespoonful two or three times a day.

The prostrating effect of digitalis should always be kept in mind by the practitioner; and the rule that when it acts upon the *stomach*, the *bowels*, the *pulse* or the *kidneys*, it should be discontinued, cannot, when we consider the power of the article, be too scrupulously observed.

Infusion of Juniper Berries, and Cream of Tartar.

R. Juniperi contusi, ʒjss.
Aquæ bullientis, Oj.

Macera per horas duas, in vase leviter clauso, et cola. Dein adde—

Potassæ Bitartratis, ʒj.

Misce. *Signa*.—Be careful to shake the bottle well before using. Of this, a teacupful may be taken four or six times a day; adding to each 40 drops of the sweet spirits of nitre. In general and local dropsy.

Infusion of Parsley Root.

R. Petroselini, ʒj.
Aquæ bullientis, Oj.

Fiat infusum et cola. A teacupful may be taken with a teaspoonful of sweet spirits of nitre, every three or four hours.

A popular and very useful remedy for strangury, arising from blisters.

Infusion of Scabious or Fleabane.

R. Erigerontis Canadensis, ʒj.
Aquæ bullientis, Oj.

Misce.—Pour on the water boiling hot, and infuse all night. A pint of the tea should be taken daily. Nitre or any other diuretic may be added. Prescribed in calculous and nephritic affections, in strangury, irritable bladder, &c.

Infusion of Uva Ursi.

R. Uvæ Ursi, ʒj.

Aquæ bullientis, Oj.

Pour the water boiling hot on the leaves, and let them macerate for twelve hours. To be taken cold, from half a pint to a pint, daily. Chiefly used in gonorrhœa and irritable bladder.

Infusion of Dandelion, &c.

R. Infusi Taraxaci, fʒiv.

Extracti Taraxaci, ʒij.

Sodæ Carbonatis, ʒss.

Potassæ Tartratis, ʒiij.

Tincturæ Rhei, fʒiij.

—————Hyoscyami, gtt. xx.

Fiat mistura. *Signa.*—One third part to be taken three times a day. In dropsical and visceral affections. DR. MEIGS.

Decoction of Pipsissewa.

R. Chimaphilæ, ʒj.

Aquæ, Oij.

Macerate for twelve hours, and then gradually boil down to one-half. Dose, a wineglassful three or four times a day. In dropsy.

Spirituuous Decoction of Pipsissewa.

R. Chimaphilæ, ʒij.

Aquæ, fʒxij.

Evaporate by heat to ʒvj., and then add two ounces of gin. Dose, a dessertspoonful every two or three hours, with diluent drinks. In dropsy accompanied by debility.

Decoction of Cainca.

R. Radicis Caincæ, ʒj.

Aquæ, Oij.

To be gradually reduced by heat to one-half. Dose, a table-spoonful three times a day. In dropsy.

This root is derived from the *Chiococca Anguifuga*, a Brazilian plant, which has of late been extensively and efficaciously employed in Europe. It occasions watery evacuations and diuresis, whence its use in dropsy.—DUNGLISON, *New Remedies*, p. 148.

Decoction of Indian Hemp.

R. Apocyni Cannabini, ʒss.

Aquæ, Ojss.

Coque ad Oj. et cola.

GRISCOM.

Signa.—Take a wineglassful three times a day. This has been found useful in dropsy, acting as a hydragogue cathartic.

The young practitioner must carefully distinguish between preparations of the native drug under consideration (*Apocynum Cannabinum*), and the powerful, imported extract of *Cannabis Sativa*; as, unfortunately, they both bear the same trivial name of *Indian Hemp*.

Decoction of Pareira Brava Root.

R. Pareiræ contusæ, ʒss.

Aquæ bullientis, Oijj.

Boil by gentle simmering down to a pint.

Signa.—Take a wineglassful every two hours.

It has a great influence over chronic inflammation of the bladder, lessening very materially the secretion of the ropy mucus, which is itself a very great evil, and, I believe, diminishing the inflammation and irritability of the bladder also.

SIR B. BRODIE.

DIURETIC SOLUTIONS AND MIXTURES.

Solution of Digitalin.

R. Digitalinæ, gr. ij.

Alcoholis diluti, fʒj.

Fiat solutio. *Signa.*—Take ten drops in sweetened water, three times daily.

Ten drops of this solution, are equivalent to one-fiftieth of a grain of digitalin, and consequently, the dose may be gradually increased until the peculiar effects of digitalis are developed.

Cider Mixture.

R. Juniperi contusi,

Sinapis,

Zingiberis, āā ʒss.

Armoraciæ contusæ,

Petroselini, āā ʒj.

Succi fermenti pomorum, Oij.

Fiat infusum.—Of which a wineglassful may be taken three or four times a day.

The above compound was recommended by the late Dr. Parrish, in cases of general dropsy which admit of stimulation, and in which the urinary secretion is deficient. The cider should be old and sound.

Mixture with Oil of Juniper.

- R. Olei Juniperi, gtt. xxv.
 Sacchari,
 Acaciæ, āā ʒij.
 Aquæ destillatæ, fʒiv.

Misce. *Signa*.—A tablespoonful to be taken every two hours.

Mixture with Juniper, Squill, &c.

- R. Potassæ Acetatis, ʒv.
 ——— Bicarbonatis, ʒj.
 Aquæ, fʒxij.
 Tincturæ Scillæ, fʒj.
 Spiritûs Juniperi Compositi, fʒj.

Fiat mistura. *Signa*.—Take two tablespoonfuls three times a day. BROWN.

Mixture of Digitalis, Acetate of Potash, &c.

- R. Infusi Digitalis, fʒiv.
 Potassæ Acetatis, ʒij.
 Spiritûs Ætheris Nitrici, fʒij.
 Aquæ Cinnamomi, fʒjss.

Fiat mistura. *Signa*.—Take a tablespoonful every four hours.

Recommended in the treatment of cardiac disease.

KILGOUR.

Mixture of Digitalis, Potash, &c.

- R. Infusi Digitalis, fʒiv.
 Tincturæ Digitalis, fʒi.
 Potassæ Acetatis, ʒj.
 Tincturæ Opii, gtt. x.

Fiat mistura. *Signa*.—Half a tablespoonful three or four times a day.

Mixture with Tincture of Cantharides, &c.

- R. Tincturæ Cantharidis,
 Spiritûs Ætheris Nitrici, āā gtt. lx.
 Sacchari, ʒj.
 Aquæ Ment hæ Viridis, fʒij.

Fiat mistura. *Signa*.—A teaspoonful may be taken every three or four hours, desisting if strangury should supervene.

Mixture of Spirits of Turpentine.

R. Olei Terebinthinæ, gtt. c.

Pulveris Acaciæ,

———— Sacchari, āā ʒj.

Aquæ Menthæ Viridis, fʒiv.

Fiat mistura. *Signa*.—A tablespoonful four or five times a day, aided by some diuretic drink.

This remedy is used with great benefit in nephritic complaints, and especially in the passage of calculi through the ureters.

Mixture of Elaterium, Colchicum, &c.

R. Elaterii, gr. j.

Spiritûs Ætheris Nitrici, fʒij.

Tincturæ Scillæ, fʒss.

Aceti Colchici, fʒij.

Syrupi, fʒj.

Fiat mistura. *Signa*.—To take a teaspoonful three or four times a day in a little water.

Dr. Ferriar recommends the above in the several forms of dropsy, and more especially in hydrothorax.

Mixture of Colchicum, Squill, and Tobacco.

R. Oxymellis Colchici, *Dub.*

———— Scillæ,

Vini Tabaci, āā fʒss.

Misce. Take a small teaspoonful in a little water four times a day.

If requisite, the *spiritus ætheris nitrici* may be added in the same quantity. This forms the *diuretic drops* of Dr. Ferriar, and requires to be given with great circumspection.

Hydragogue Draught of Dr. Ferriar.

R. Gambogiæ, gr. iv.

Spiritûs Ætheris Nitrici, fʒj.

Tincturæ Sennæ, fʒij.

Syrupi Rhamni, *Ed.*

Aquæ Menthæ Viridis, āā fʒss.

Misce et fiat haustus.—Dr. Ferriar observes that, in some cases of hydrothorax attended with imminent danger of suffocation, he has afforded immediate relief by the above draught.

Tincture of Colchicum and Digitalis.

R. Tincturæ Colchici Seminis,
 ——— Digitalis, āā fʒj.

Spiritûs Ætheris Nitrici, fʒj.

Misce. Dose, twenty drops on a lump of sugar. Given as a diuretic in dropsy and hydrothorax. HILDENBRAND.

Mixture of Carbonate of Potash, Squill, &c.

R. Potassæ Carbonatis, ʒj.

Succi Limonis, q. s. ad saturandum,

Tincturæ Scillæ, fʒss.

——— Opii, gtt. xxx.

Aquæ Cinnamomi, fʒiv.

Misce. *Signa.*—A tablespoonful every two or three hours throughout the day.

The vinegar or oxymel of squill, if preferred, may be substituted in this preparation. We use the tincture because it is more uniform, and less liable to spontaneous changes.

Mixture of the Acetates of Potash and Ammonia.

R. Liquoris Ammoniæ Acetatis, fʒij.

Potassæ Acetatis, ʒij.

Misce. *Signa.*—A tablespoonful every three hours for a dose.

Mixture of Buchu and Acetate of Potash.

R. Infusi Buchu, fʒviij.

Potassæ Acetatis, ʒij.

Misce. *Signa.*—Take a wineglassful three or four times a day.

This is an excellent diuretic combination, and its value may be farther enhanced by the addition of sweet spirits of nitre.

Mixture with Urea.

R. Ureæ, ʒj.

Aquæ, fʒijss.

Syrupi Aurantii Corticis, fʒss.

Fiat mistura. *Signa.*—Take a tablespoonful every six hours.

Its action as a diuretic should be aided by the free administration of diluents, as well as by keeping the skin moderately cool. The dose may be augmented to a scruple or more.

T. H. TANNER.

Hufeland's Diuretic Drops.

R. Olei Juniperi, fʒss.

Spiritûs Ætheris Nitrici,

Tincturæ Digitalis, āā fʒiij.

Misce. *Signa.*—Give twenty or thirty drops every three hours.

Use. This is one of the best diuretics, and may be employed for some time without offending the stomach.

HUFELAND.

Mixture of Buchu and Uva Ursi.

R. Buchu,

Uvæ Ursi, āā ʒss.

Aquæ ferventis, Oss.

Digere leni calore in vase clauso per dimidiam horam.

Cola, et adde—

Syrupi Senegæ, fʒss.

Signa.—Take one or two tablespoonfuls every two hours.

Used in blennorrhœa, atony, and paralysis of the bladder.

CLARUS.

Mixture of Balsam Copaiba.

R. Copaibæ,

Spiritûs Ætheris Nitrici, āā fʒss.

Pulveris Acaciæ, ʒij.

——— Sacchari, ʒj.

Aquæ destillatæ, fʒiv.

Spiritûs Lavandulæ Compositi, fʒij.

Tincturæ Opii, fʒj.

Fiat mistura secundum artem. *Signa.*—A tablespoonful three times a day.

CHAPMAN.

This remedy is more especially used in gonorrhœa, and is considered by Professor Chapman as a specific in that disease. A preferable mode of exhibiting balsam copaiva, consists in dropping the quantity prescribed for a dose in a wineglass of water, and then adding to it a small quantity of the *tinct. gentiænæ comp.* The balsam collects into a globule and is surrounded by the tincture, which, leaving its taste in the mouth, covers that of the nauseous balsam.

Mixture of Balsam Copaiba, Cubebs, &c.

R. Copaibæ,
 Pulveris Cubebæ, āā ʒss.
 Acaciæ, ʒij.
 Sacchari, ʒij.
 Aquæ, fʒviij.
 Tincturæ Opii Camphoratæ, fʒss.

Misce secundum artem. *Signa.*—A tablespoonful every three hours.

We have found this remedy, in some obstinate cases of gonorrhœa, more efficacious than the preceding formula. The tincture of cubebs may be substituted for the powder—and the tinct. opii acetat. fʒj. for the paregoric, if desirable. Balsam copaiva is sometimes given by the French physicians in the form of enema, for the cure of gonorrhœa. Capsules containing it are also in common use.

S. G. MORTON.

Another Form.

R. Copaibæ, fʒjss.
 Mucilaginis Acaciæ, fʒv.
 Olei Limonis, gtt. vj.
 Tincturæ Cubebæ, fʒj.
 Spiritûs Ætheris Nitrici, fʒss.

Fiat mistura. *Signa.*—Take a tablespoonful three times a day, in gonorrhœa.

RANDOLPH.

Compound Mixture of Copaiba, &c.

R. Copaibæ,	} āā fʒss.
Alcoholis,	
Aquæ Menthæ Piperitæ	
Syrupi,	
Spiritûs Ætheris Nitrici, fʒss.	

Fiat mistura.—To be taken during the day in four doses, followed by demulcent drinks. In chronic gonorrhœa.

Mixture of Fluid Extract of Cubebs.

R. Extracti Cubebæ Fluidi, fʒij.
 Mucilaginis Acaciæ,
 Syrupi Tolutani, āā fʒss.
 Spiritûs Lavandulæ Compositi, fʒij.
 Aquæ, fʒijss.

Fiat mistura. *Signa.*—Give a tablespoonful every three hours in gonorrhœa.

EDITOR.

CLASS IV.

ANTILITHICS AND LITHONTRIPTICS.

ANTILITHICS are those substances which arrest or prevent the formation of calculi in the kidneys or bladder. Lithontriptics are supposed to possess the power of dissolving calculous concretions, by acting on the substance which serves as the bond of union to the saline or acid particles composing the stone.

To administer these remedies with any degree of success, requires a chemical knowledge of the ingredients which enter into the formation of the different calculi; attention must also be given to the properties of the urine.

As derangement of the digestive organs almost always co-exists with these foreign bodies in the urinary apparatus, we shall gain a very considerable advantage by combining the remedies of this class with those which have a tendency to restore the healthy action of the stomach and bowels.

When lithic or uric acid abounds in the urine, the alkalies will be found to produce the greatest benefit; but where the salts are evidently of the calcareous or magnesian character, the muriatic, or some one of the mineral acids, will constitute the most appropriate remedy.

POWDERS.

Powders of Uva Ursi, Bark, and Opium.

R. Pulveris Uvæ Ursi,
——— Cinchonæ, āā ʒj. vel ʒij.
——— Opii, gr. iij.

Misce, et divide in chartulas vj.—Of which one may be taken three or four times a day.

The above is the original prescription of Dr. Ferriar, who advises the patient to drink two ounces of lime-water after each dose.

PILLS.

Soap Pills.

R. Saponis, ʒj.

Divide in pilulas xv. *Signa.*—One every two hours during the day.

Pills of dried Carbonate of Soda.

R. Sodæ Carbonatis Exsiccatae,

Saponis, āā ʒj.

Extracti Glycyrrhizæ,

Aquæ, āā q. s.

Fiat massa in pilulas xxx. dividenda.

Signa.—Take three pills, three times a day, as an antilithic.

BEDDOES.

SOLUTIONS, MIXTURES, &c.

Solution of Bicarbonate of Soda, &c.

R. Sodæ Bicarbonatis, ʒj.

Infusi Quassia, fʒiv.

Tincturæ Colombæ, fʒj.

Misce. *Signa.*—A tablespoonful for a dose four times a day.

Solution of Salaëratu.

R. Potassæ Bicarbonatis, gr. x. vel xv.

Aquæ Seltzer, fʒvj.

Misce.—Draught, to be taken three or four times a day.

Mixture of Bicarbonate of Soda, &c.

R. Sodæ Bicarbonatis, ʒj.

Misturæ Amygdalæ, fʒiv.

Copaibæ, fʒij.

Tincturæ Opii, gtt. lx.

Fiat mistura secundum artem. *Signa.*—A tablespoonful *pro re nata*. In acute pain of the bladder.

In combination with each of these alkaline preparations, demulcent drinks may with benefit be freely employed.

Carbonate of Potash and Lime-water.

R. Potassæ Carbonatis, ʒij.

Aquæ Calcis, Oij.

Fiat solutio.—Of which a wineglassful may be taken every two hours, mixed with an equal quantity of new milk.

Mixture of Magnesia, &c.

R. Magnesiæ Carbonatis, ʒj.

Infusi Gentianæ Compositi, fʒvj.

Fiat mistura.—A wineglassful to be taken three times a day.

This mixture is recommended by Mr. Brande, of London, as superior to the alkalies, in cases where uric acid abounds.

Mixture of Muriatic acid in Barley-water.

R. Acidi Muriatici, gtt. x. vel xx.

Decocti Hordei, fʒviij.

Misce. Dose, a tablespoonful largely diluted, three or four times a day.—In stone, when the calcareous properties are most apparent.

Nitric acid has been given with the same intention, but appears to be less useful than muriatic.

Oil of Turpentine.

R. Olei Terebinthinæ, fʒss.

Signa.—From thirty drops to a large teaspoonful may be taken on sugar, and repeated *pro re nata*. See also the formula given under Diuretics.

This medicine has been frequently given of late in the passage of urinary calculi down the ureters. It is stated by Dr. Physick that this remedy, proportioned in its dose to the degree of suffering, affords more prompt relief than any other.

Urethral Injection of Carbonate of Soda.

R. Sodæ Carbonatis, ʒj.

Saponis, ʒij.

Aquæ destillatæ, fʒxij.

Fiat solutio et cola.

Signa.—Use as an injection, for softening uric acid deposits, and calculi.

HÔPITAL DE MONTPELLIER.

CLASS V.

DIAPHORETICS.

DURING the exhibition of this class of medicines the patient must be confined to his bed; and when diaphoresis is once excited, every precaution should be taken lest it be suddenly checked.

Venesection should generally be resorted to when there is much arterial excitement with great heat of the skin, before we attempt to give diaphoretics. When the skin continues steadily hot and dry, all external warmth must be removed, and cold water freely applied to the temples, hands, feet, &c., while the room is to be kept cool by a free current of fresh air. Under these circumstances, cold water may be frequently and plentifully drunk by the patient. Diaphoretics given in conjunction with these general means, will act with much more certainty than if great heat of the surface is maintained by covering the patient with heavy bedclothes. If it be desirable to produce perspiration when the heat of the skin is not above the natural standard, we shall obtain our object most readily by placing the patient in a warm bed, giving him warm drinks freely, placing his feet in warm water, or applying hot bottles, or bags of sand, to the different parts of his body. The bags of sand accommodate themselves to the inequalities of the surface, and therefore form an excellent mode of applying heat externally. Warm bricks, covered with cloths, and wetted with vinegar, may also be used advantageously. Employed in this way they form a steam-bath. The same effect may be produced by conducting the vapor of alcohol through a tube, under the bedclothes of the patient.

In a few words, then, we have described the manner of producing diaphoresis under opposite states of the system. In the first instance we solicit, in the second we rather extort it.

It may be proper to observe, that in the febrile affections the frequent change of linen about the patient is not only warrant-

able, but absolutely necessary, whilst employing this class of remedies.

Where, however, we are desirous of producing what is emphatically termed "*a sweat*," we should envelop the patient in a flannel shirt or gown, and avoid, as was before observed, every cause which may suddenly check the cutaneous discharge, and more especially purging and diuresis.

In the treatment of ordinary fevers, we usually resort to articles recognized by the profession as being capable of producing perspiration, or sweating, soon after their exhibition; while in chronic diseases, as syphilis, scrofula, and the cachectic state following the use or abuse of mercury, others are employed which combine an alterative property with their tendency to the skin. It is therefore deemed advisable to make two sub-classes, viz. 1. Ordinary Diaphoretics, and 2. Alterative Diaphoretics.

SUB-CLASS. I.—ORDINARY DIAPHORETICS.

POWDERS.

Dover's Powder.

R. Pulveris Ipecacuanhæ et Opii, ʒss. vel ʒj.

Divide in pulveres vj.

Signa.—Take one powder three times a day.

The most material difference between the officinal powder and that of Dr. Dover, made public in 1733, is in the substitution of the sulphate of potash for saltpetre.

The principal use of the former salt in this preparation, appears to be to facilitate the minute division of the other two ingredients. Its crystals are extremely hard, and the prolonged trituration necessary to reduce them, renders the opium and ipecacuanha impalpable. It is one of the happiest and most useful combinations, having stood the test of time and experience with undiminished reputation. To promote its operation, *warm drinks* may be freely taken, but not immediately after the dose, lest vomiting be induced.

Antimonial Powder.

R. Pulvis Antimonialis, *Ed.* ʒj.

Divide in chartulas vj. *Signa.*—One every two or three hours in sirup or molasses. In fevers.

This powder, which has been introduced as the succedaneum

of the celebrated Fever Powder of Dr. James has lately been pronounced inert by Mr. Richard Phillips, chemist of London. Learning that it had produced no effect when largely exhibited, he determined to examine it *chemically*, and found it to consist of peroxide of antimony (which he says is inert), and the phosphate of lime in mixture. But, as Dr. Paris observes, "it will be difficult for the chemist to persuade the physician he can never have derived any benefit from the exhibition of antimonial powder."

Nitrous Powders.

R. Pulveris Potassæ Nitratis, ʒj.

Antimonii et Potassæ Tartratis, gr. ss.

Hydrargyri Chloridi Mitis, gr. vj.

Fiant pulveres vj. *Signa*.—One to be taken every two hours in sirup or molasses. In febrile affections.

The calomel may be omitted or employed, at the discretion of the practitioner.

Powders with Opium, Ipecacuanha, Calomel, and Nitre.

R. Pulveris Opii, gr. iij.

———— Ipecacuanhæ, gr. vj.

Hydrargyri Chloridi Mitis, gr. jss.

Potassæ Nitratis, ʒss.

Fiant pulveres vj. *Signa*.—One to be taken every three or four hours.

Powders with Calomel, Opium, and Ipecacuanha.

R. Pulveris Opii, gr. iv.

Hydrargyri Chloridi Mitis, gr. xvj.

Pulveris Ipecacuanhæ, gr. viij.

Divide in pulveres viij. *Signa*.—One to be taken every hour or two hours.

This combination is recommended by Professor Chapman in dysentery. From its evident diaphoretic tendency we have been induced to place it under this head.

Powders of Precipitated Sulphuret of Antimony, Calomel, &c.

R. Antimonii Sulphureti Præcipitati,

Hydrargyri Chloridi Mitis, āā gr. vj.

Pulveris Guaiaci, ʒj.

Fiant chartulæ vj. *Signa*.—One to be taken morning and evening, with double or even triple proportion of the ingredients, if necessary. In cutaneous diseases.

Soda Powder of Guy's Hospital.

- R. Sodæ Carbonatis exsiccatæ, ʒj.
 Hydrargyri Chloridi Mitis, gr. xij.
 Pulveris Cretæ Compositi, *Lond.* ʒij.

Misce. *Signa*.—Give three to eight grains at night for a week, in infantile gastric fever; followed in the morning by five grains of rhubarb mixed with ten grains of the sulphate of potash. Dr. Fordyce says of the above: "Had I been more ambitious of dying a rich man than of living a useful member of society, the powers of our antihectic powder in curing, as if by miracle, the hectic fever and the swelled bellies of children in London, would have remained a secret while I lived."

GOLDING BIRD.

Powders with Guaiacum, &c.

- R. Pulveris Guaiaci,
 ——— Potassæ Nitratis, āā ʒj.
 ——— Ipecacuanhæ, gr. iij.
 ——— Opii, gr. ij.

Fiant pulveres vj. *Signa*.—One to be taken every three hours in a little jelly or molasses.

A stimulating diaphoretic, and particularly serviceable in acute rheumatism after the reduction of inflammatory symptoms.

DIAPHORETIC MIXTURES.

Neutral or Saline Mixture.

- R. Succī Limonis recentis, fʒiss.
 Potassæ Carbonatis, q. s. ad saturandum,
 Sacchari, ʒij.
 Antimonii et Potassæ Tartratis, gr. ss.
 Aquæ destillatæ, fʒiij.

Misce.—A tablespoonful to be taken every two hours. In fevers.

This preparation is one of our most agreeable diaphoretics. Its powers are decidedly increased by the addition of tartar emetic; which, however, may be omitted, and the sweet spirits of nitre substituted, if preferred.

Solution of Citrate of Potash.

- R. Potassæ Citratis, ʒiij.
 Olei Limonis, gtt. ij.
 Sacchari, ʒij.
 Aquæ destillatæ, fʒiv.

Misce.—A tablespoonful to be taken every two hours. This

preparation is a good substitute for the preceding one, and is in several respects to be preferred. Spirits of nitre may be added.

Effervescing Draught.

R. Potassæ Carbonatis, ʒij.

Aquæ destillatæ, fʒiv.

Misce, et fiat solutio.

R. Succi Limonis recentis,

Aquæ destillatæ, āā fʒij.

Misce. *Signa.*—Mix two tablespoonfuls of the lemonade with one of the alkaline solution, and let it be drunk while effervescing, to be repeated *pro re nata*.

This preparation differs from the neutral mixture more in the mode of exhibition than in any other important particular. It is of great value when irritability of the stomach precludes the use of ordinary diaphoretics.

Spirits of Mindereri.

R. Aceti destillati, vel Acidi Acetici diluti, fʒvj.

Ammoniæ Carbonatis, q. s. ad saturandum.

Add the carbonate of ammonia gradually to the acid or vinegar until it is saturated.

Signa.—Take a tablespoonful every two hours.

This excellent diaphoretic in inflammatory diseases, appears to have been first used by Boerhaave, in 1732. With us it is generally prescribed in combination, under one of the following forms:—

Spirits of Mindereri and Antimonial Wine.

R. Liquoris Ammoniæ Acetatis, fʒij.

Aquæ Cinnamomi, fʒj.

Vini Antimonii, fʒj.

Aquæ destillatæ, fʒij.

Fiat mistura. *Signa.*—A tablespoonful every two or three hours.

Spirits of Mindereri, Sweet Spirits of Nitre, &c.

R. Liquoris Ammoniæ Acetatis, fʒvj.

Syrupi Limonis, fʒij.

Vini Antimonii, fʒjss.

Spiritûs Ætheris Nitrici, fʒij.—fʒss.

Tincturæ Opii, gtt. xlv.

Fiat mistura. *Signa.*—Take a tablespoonful every two hours.

EBERLE.

Solution of Citrate of Ammonia, &c.

R. Succî Limonis recentis, f̄3j.

Ammonia Carbonatis, q. s. ad saturandum.

Sacchari, 5j.

Aquæ destillatæ, f̄3iv.

Fiat mistura. *Signa*.—A tablespoonful may be taken every two or three hours, *pro re nata*.

This preparation is seldom employed in the practice of this city. It may, however, be beneficially given as a pleasant stimulating diaphoretic.

Sweet Spirits of Nitre, and Antimonial Wine.

R. Spiritûs Ætheris Nitrici, f̄3j.

Vini Antimonii, f̄3ij.

Misce. *Signa*.—A teaspoonful may be taken every hour or two hours, in barley-water, during the hot stage of intermittent and other fevers.

Sweet Spirits of Nitre, with Antimonial Wine and Laudnum.

R. Tincturæ Opii, gtt. xxv.

Spiritûs Ætheris Nitrici, f̄3j.

Vini Antimonii, gtt. xx.

Aquæ destillatæ, f̄3ss.

Fiat mistura.—To be given at a dose, and repeated *pro re nata*.

Dr. Chapman observes, in his Therapeutics, that "this combination very frequently acts as a diuretic in dropsy, and has more than an once, within my knowledge, removed the effusion in this disease, under circumstances the least to be expected."

Mixture of Saltpetre and Tartar Emetic.

R. Potassæ Nitratiss, 3ij.

Antimonii et Potassæ Tartratis, gr. j.

Pulveris Extracti Glycyrrhizæ,

———— Acaciæ, āā 3ij.

Aquæ Fœniculi,

———— destillatæ, āā f̄3iijss.

Fiat mistura. *Signa*.—Give a tablespoonful every two hours.

This is proposed as a substitute for the nitrous powders, as being less apt to offend the stomach than the same substances in a pulverulent form.

EBERLE.

Mixture with Guaiacum, &c.

- R. Pulveris Guaiaci Resinæ, ʒij.
 ——— Potassæ Nitratis, ʒiss.
 Antimonii et Potassæ Tartratis, gr. j.
 Pulveris Acaciæ, ʒj.
 ——— Extracti Glycyrrhizæ, ʒj.
 Aquæ destillatæ, fʒviij.
 Fiat mistura. *Signa*.—A tablespoonful three or four times a day.

This preparation, if made without the extract of liquorice, is disposed to turn green, which gives it an unpleasant appearance, and sometimes induces the patient to refrain from it until the physician arrives. It is employed principally in rheumatic cases.

Solution of Nitrate of Potash.

- R. Decocti Hordei, Oj.
 Potassæ Nitratis, ʒij.
 Succī Limonis, fʒj.
 Fiat potus.—To be used warm as a common drink.

Tincture of Tolu with Wine of Antimony.

- R. Tincturæ Tolutanæ, fʒv.
 Pulveris Acaciæ, ʒiij.
 Sacchari, ʒj.
 Vini Antimonii, fʒj.
 Aquæ Cinnamomi, fʒij.
 Fiat mistura. *Signa*.—A tablespoonful three or four times a day.

This is a stimulating diaphoretic, and will be found serviceable in pectoral diseases accompanied with an excessive mucous discharge depending on debility of the bronchial membrane.

Mixture with Aconite and Wine of Colchicum.

- R. Extracti Aconiti Alcoholici, gr. xij.
 Syrupi, fʒj.
 Vini Colchici Seminis, fʒss.
 Fiat mistura. *Signa*.—Take fifteen drops every three hours.

Used in acute and chronic rheumatism, gradually increasing the dose.

WEBER.

Mixture with Aconite and Valerian.

R. Pulveris Valerianæ, ʒss.

Aquæ bullientis, f ʒvj.

Macula per horas duas et cola. Dein adde—

Extracti Aconiti Alcoholici, gr. vj.

Syrupi, f ʒss.

Liquoris Ammoniacæ Acetatis, f ʒj.

Misce. *Signa.*—Take a tablespoonful every two hours.

Used in nervous gout.

RICHTER.

Mixture with Arnica.

R. Arnicæ Florum, ʒij.

Aquæ bullientis, f ʒvj.

Macula per horas duas, et cola. Dein adde—

Pulveris Acaciæ, ʒij.

Syrupi Aurantii Corticis, f ʒj.

Misce. *Signa.*—Take a tablespoonful every two hours in low nervous fever, as a stimulant diaphoretic.

BERENDS.

I have selected the above, out of thirty German prescriptions for arnica, as being one of the most simple, and perhaps one of the best forms, for administering an article, which has hitherto attracted little attention among us, although it is frequently prescribed by men of the highest standing in Germany.

EDITOR.

INFUSIONS.

Infusion of Boneset.

R. Eupatorii, ʒj.

Aquæ bullientis, Oj.

Infuse for thirty minutes in a covered vessel; then decant and take a wineglassful every hour, till perspiration is freely induced. The infusion must be taken as hot as it can be drunk; otherwise it may produce vomiting.

Infusion of Pipsissewa.

R. Chimaphilæ, ʒj.

Aquæ bullientis, Oj.

Fiat infusum.—A teacupful every two hours.

Infusion of the Asclepius Tuberosa, or Pleurisy Root.

R. Radicis Asclepiadis Tuberosæ, contritæ, ʒj.

Aquæ bullientis, Oiss.

Fiat infusum. *Signa*.—A teacupful to be taken warm, as frequently as the stomach will bear it. CHAPMAN.

Professor Chapman observes of this plant: "As a diaphoretic, I think it is distinguished by great certainty and permanency of operation, and has this estimable property, that it produces its effects without increasing much the force of the circulation, raising the temperature of the surface, or creating inquietude and restlessness."—*Therapeutics*, vol. i. p. 347.

I have found the virtues of the pleurisy root to be better developed, by boiling one ounce in a quart of water down to a pint, and straining. EDITOR.

SUB-CLASS II.—ALTERATIVE DIAPHORETICS.

PILLS.

Pills of Red Sulphuret of Mercury, &c.

R. Hydrargyri Sulphureti Rubri,

Pulveris Serpentariæ, āā ʒss.

Syrupi, q. s.

Fiant pilulæ xij. *Signa*.—Two of these to be taken three times a day.

They are alterative as well as diaphoretic, and are useful in herpetic and other obstinate cutaneous complaints.

Pills of Sulphuret of Antimony, &c.—Plummer's Pills.

R. Hydrargyri Chloridi Mitis,

Antimonii Sulphureti præcipitati, āā ʒss.

Pulveris Guaiaci Resinæ, ʒj.

Sacchari Fæcis, q. s. ut fiant pilulæ lx.

Three pills to be taken night and morning.

These pills were recommended by Professor Plummer, in leprosy, secondary syphilis affecting the skin, gleans, old ulcers, &c. &c. The decoction of guaiacum generally accompanies their use.

DECOCTIONS.

Sirup or Rob Anti-syphilitique of L'afsecteur.

- R. Sarsaparillæ,
 Arundinis Phragmitis, āā 3xxx.
 Florum Boraginis Officinalis, 3viij.
 Sennæ,
 Rosæ Albæ, āā 5ij.
 Sacchari,
 Mellis, āā 1bvj.
 Aquæ, q. s.

Boil the sarsaparilla and marsh reed grass in nine pints of water for one hour; strain off the decoction, and pour the same quantity of water on the residuum, which is to be boiled for two hours; towards the end of the boiling add the borage flower, senna, and rose leaves, and then strain off, and to both decoctions add the sugar and honey, and boil the whole to the consistence of a sirup.

The dose for a man is six tablespoonfuls, and for a woman four, taken without addition, at six in the morning.

The sarsaparilla ptisan is then to be commenced, and is made as follows:—

Boil 5ij. of Sarsaparilla in three quarts of water, to two quarts, suffer it to infuse during the night, and strain it off in the morning.

This is to be drunk by the tumblerful, and several are to be taken during the day.

The above preparation has been asserted, by the New York Medical Society, to be nearly identical with the noted Panacea of Swain. It is believed that corrosive sublimate is often introduced into these nostrums at the discretion of the vender. That both the French and American have frequently salivated, is well ascertained.

Decoction of Burdock.

- R. Radicis Lappæ, 3iij.
 Aquæ, Oijj.

Reduce the fluid by boiling to two pints, and strain. A teacupful may be taken for a dose, five or six times a day.

In syphilis, scurvy, arthritic affections, &c.

Decoction of Mezereon.

R. Mezerei, ℥vj.

Aqua, Ovj.

Reduce by boiling to ℔iv., and near the end of the decoction, add Radicis Glycyrrhizæ, ℥j. Strain, and give from one to four pounds per diem, as the stomach will bear it. In scrofula complicated with syphilis, &c.

Compound Decoction of Sarsaparilla.

R. Decocti Sarsaparillæ Compositi, Oj.

Give a teacupful four or five times a day.

This preparation, a formula for which is to be found in almost every Pharmacopœia, has long been celebrated in the sequelæ of syphilis, and the disease produced by the improper exhibition of mercury. It is also much employed in rheumatism. The rules laid down in the Dispensatories for the preparation of the decoction should be closely observed.

Decoction of Zittmann.

This preparation has long been used successfully in Germany and England, and is now coming into notice in this country, as a valuable remedy in the treatment of secondary syphilis, complicated with a scrofulous diathesis. The formulæ given below I have translated from a work published by Prof. Walther, of Leipzig, in 1847, and which he says are copied from Zittmann's original prescription (nach der Originalvorschrift.) It will be seen that the strength of the decoction varies materially from the formulæ heretofore published.

EDITOR.

R. Sarsaparilla, in chips, ℥xij.

Digest in a tinned vessel for 24 hours, with spring water, ℔xlviij., then add, inclosed in a small bag—

Of Saccharine Alum,¹ ℥jss.

" Calomel, ℥ss.

" Cinnabar, ℥j.

Boil down to ℔xvj. When the boiling is nearly completed add—

Of Senna, ℥iij.

" bruised Aniseed,

" Fennelseed each, ℥ss.

Remove from the fire and add—

Of bruised Liquorice root, ℥jss.

And when cold, strain, and mark: *The Strong Decoction.*

¹ Saccharine Alum consists of Alum four parts, and Kino one part.

- R. To the dregs of the strong decoction, add,
 Sarsaparilla, in chips, ℥vj.
 Digest for twenty-four-hours in spring water, ℔xlviij.
 Then boil down to ℔xxvi. When the boiling is nearly completed, add—
 Of Lemon Peel, sliced, }
 “ Cinnamon, bruised, } each ℥ijj.
 “ Cardamoms, “ }
 Remove from the fire; add—
 Of bruised Liquorice root, ℥vj.
 When cold, strain, and mark: *The Weak Decoction.*

Patients with constitutional, or secondary syphilis, or inveterate skin diseases complicated with a scrofulous habit, are treated in the following manner: At night, the patient takes a purgative dose of calomel and jalap, or some other active cathartic, “and the next morning begins the regular course, by drinking one-half of a quart bottle of the strong decoction, taking a glass of it warm every half hour, in bed to bring out the perspiration. In the afternoon, he takes a whole bottle of the weak decoction, cold, in divided doses; and in the evening he drinks (cold) the remaining half of his bottle of strong decoction. This is repeated for four days. On the sixth, a purgative is again administered; and for the following four days the decoctions are used. On the eleventh, pills are again given, and then all treatment is suspended for a week. If the cure is not completed, the same process is again resorted to. The diet should consist of light broths, bread, and lean roasted meats.”

Although the above prescription cannot be defended on strict chemical and pharmaceutical grounds, yet the remarkable success attending its use in many of the hospitals of Europe, entitle it to notice.

MIXTURES.

Mixture of Muriate of Baryta and Iron.

- R. Barii Chloridi, ℥ss.
 Aquæ Menthæ Piperitæ, f ℥ijss.
 Tincturæ Ferri Chloridi, f ℥ss.
 Syrupi Aurantii Corticis, f ℥j.
 Fiat mistura. *Signa.*—Take a teaspoonful every three hours.

Used in obstinate scrofulous diseases. Each dose contains about one grain of the chloride of barium, and seventeen drops of the muriated tincture of iron. This corresponds in strength with a prescription by

FISCHER.

Mixture with Buchu and Iodide of Potassium.

- R. Infusi Buchu, f ʒviij.
 Liquoris Potassæ, f ʒj.
 Potassii Iodidi, ʒss.
 Spiritûs Ætheris Nitrici, f ʒiij.
 Fiat mistura. *Signa*.—Take two tablespoonfuls three times a day.

Used in chronic rheumatism.

S. WRIGHT.

Corrosive Sublimate Solution.

- R. Hydrargyri Chloridi Corrosivi, gr. j.
 Solve in Aquam, f ʒss.
 Dein adde—
 Syrupi Sarsaparillæ Compositi, f ʒviijss.
 Misce bene. *Signa*.—Take a tablespoonful three times daily.

Used in chronic rheumatism, secondary syphilis, and various skin diseases.

EDITOR.

Sarsaparilla Beer.

- R. Sarsaparillæ contusæ, ℥ij.
 Pulveris Guaiaci Corticis, ʒviij.
 Ligni Guaiaci concisi,
 Anisi,
 Radicis Glycyrrhizæ contusæ, āā ʒiv.
 Mezerei, ʒj.
 Sacchari Fæcis, ℥ij.
 Caryophylli contusi, ʒss.
 Aquæ bullientis, congios iv.

Mix these ingredients, and agitate the vessel thrice a day. When a fermentation has well begun it is fit for use, and may be taken in the dose of a small tumblerful twice or thrice a day.

“This is the Spanish recipe for the Jarave or diet drink, somewhat modified. If intended for old and obstinate complaints, as leprous affections, elephantiasis, various anomalous ulcerations and foul disorders of the skin, there may be added to the jug, *after fermentation has well begun*, tartrate of antimony twelve grains, corrosive sublimate eight or ten grains, muriate of ammonia one dram, dissolved previously in a little water. These remedies greatly increase the alterative powers of the vegetable infusion, and so effectually prevent its decomposition that it may be kept for a long time unaltered, *even in a hot climate*.”—See *Hancock on Sarsaparilla. Journal of the Philadelphia College of Pharmacy*, vol. i. No. 4.

CLASS VI.

EXPECTORANTS.

DURING the administration of Expectorants the patient should be kept warm, and, if the complaint be violent, in bed. The dose of all Expectorants should be repeated sufficiently often to allay the irritation as much as possible; for it is obvious that the act of coughing increases that very irritation which originally gave rise to it.

POWDERS, PILLS, AND LOZENGES.

Powders of Myrrh, Ipecacuanha, &c.

R. Pulveris Myrrhæ, gr. xij.
——— Ipecacuanhæ, gr. vj.
——— Potassæ Nitratis, ʒss.

Misce, et divide in partes iv. One to be taken every fourth hour. A stimulating expectorant.—PARIS'S *Pharmacologia*.

Compound Powder of Ipecacuanha.

R. Pulveris Ipecacuanhæ, et Opii, ʒss.
Divide in chartulas iij. *Signa*.—One, two or three times in 24 hours, mixed in sugar and water.

This, the Dover's powder, has been already introduced under the head of Diaphoretics, where, perhaps, it most properly belongs; but it often proves of such decided advantage as an expectorant, as to deserve an insertion here.

Powders with Nitrate of Potash, Calomel, Opium, &c.

- R. Potassæ Nitratis, ʒj.
 Hydrargyri Chloridi Mitis, gr. iij.
 Pulveris Opii, gr. iij.
 ——— Ipecacuanhæ, gr. iij. vel vj.
 Misce, et divide in chartulas vj. *Signa*.—One every two or three hours in sirup or molasses.

This preparation closely resembles the former, with the exception that it contains *calomel*.

All powders containing calomel, or other metallic preparations, should be given in sirup, as they sink when water is employed as the vehicle.

Pills of Sulphate of Zinc and Myrrh.

- R. Zinci Sulphatis, gr. x.
 Pulveris Myrrhæ, ʒiss.
 Confectionis Rosæ, q. s. ut fiant pilulæ xx.

This prescription is taken from the Pharmacologia of Dr. Paris, who recommends two of the pills to be taken twice a day.

It is principally used in pertussis; but a more efficient direction would be to reduce the myrrh to a dram, on account of the size of the pills, and to give a pill every two hours. Dr. Paris remarks, that the principal nostrums for whooping-cough are formed either of opium or sulphate of zinc.

Pills of Conium, Ipecacuanha, &c.

- R. Extracti Conii,
 Pulveris Ipecacuanhæ et Opii, āā gr. x.
 Misce, et fiant pilulæ v.—Of which two may be taken at bedtime. In pulmonary irritation, attended with rheumatic and other local pains.

Pills of Tartar Emetic, &c.

- R. Antimonii et Potassæ Tartratis,
 Pulveris Opii, āā gr. jss.
 ——— Tragacanthæ, gr. x.
 Confectionis Rosæ, q. s. ut fiant pilulæ xxv. Two to be taken night and morning. In chronic pulmonary catarrh.
 DR. PARISET.

Pills of Squill, Ammoniac, &c.

R. Pulveris Scillæ, gr. xxx.

Ammoniaci, ʒiss.

Extracti Conii, gr. xxx.

Divide into xxx. pills; of which two may be taken twice a day.

In asthma and chronic catarrh.

Pills of Squill, Calomel, &c.

R. Hydrargyri Chloridi Mitis, gr. iij.

Scillæ recentis,

Ammoniaci, āā ʒj.

Pulveris Ipecacuanhæ et Opii, ʒss.

Confectionis Rosæ, q. s. ut fiat massa, et divide in pilulas xxx.

One, three or four times a day. In troublesome cough.

DR. LATHAM.

The recent squill is greatly preferred by some physicians. It is generally given in double the dose of the dried root.

Pills of Calomel, Squill, &c.

R. Hydrargyri Chloridi Mitis, gr. xxiv.

Pulveris Scillæ, ʒss.

Antimonii et Potassæ Tartratis, gr. vj.

Pulveris Opii, gr. xvij.

Syrupi, q. s.

Fiant pilulæ xxiv. *Signa.*—One every night. In catarrhal affections.

Pills of Sulphate of Zinc.

R. Zinci Sulphatis, ʒss.

Confectionis Rosæ, q. s. ut fiant pilulæ xx.—One of these pills may be given three times a day. In phthisis and pertussis.

Tar Pills.

R. Picis liquidæ, ʒj.

Pulveris Glycyrrhizæ, q. s. ut fiant pilulæ xxx.—One to be taken every three or four hours.

Pills of Tar, Gentian, and Quassia.

R. Picis liquidæ, ʒj.

Extracti Quassia, gr. xx.

Pulveris Gentianæ, q. s. ut fiant pilulæ xx. *Signa.*—To take a pill morning, noon, and night.

Tar may be advantageously made into pills with powdered gentian alone, and thus combine the tonic and expectorant effect.

Pills of Myrrh, &c.

- R. Myrrhæ, ʒiss.
 Pulveris Scillæ, ʒss.
 Extracti Hyoscyami, ʒij.
 Aquæ destillatæ, q. s. ut fiant pilulæ xxx.—Two to be taken night and morning.

These pills are recommended by Dr. Paris, in phthisis and catarrh attended with great debility of the pulmonary organs.

Pills of Cyanuret of Potassium.

- R. Potassii Cyanureti, gr. j.
 Amyli, gr. iv.
 Syrupi, q. s. ut fiant pilulæ viij. Dose, a pill night and morning, watching its effect.

See remarks on Prussic Acid, page 105.

Wistar's Cough Lozenges.

- R. Pulveris Acaciæ,
 ——— Extracti Glycyrrhizæ,
 ——— Sacchari, aa ʒij.
 ——— Opii, gr. vj.
 Olei Anisi, gtt. iv.
 Misce, et adde—
 Aquæ destillatæ, q. s. ut fiant trochisci lx. *Signa.*—One of these may be dissolved in the mouth three or four times a day.

Emetia Lozenges.

- R. Emetiæ Fuscæ, gr. xxxij.
 Sacchari, ʒij.
 Mucilaginis Acaciæ, q. s.
 Divide into lozenges, each weighing ten grains. One of these will produce more or less nausea in a child; but if this is not desirable, they can be given in divided portions. Adults may take several at intervals of two hours, without the risk of vomiting.

Cubeb Lozenges.

- R. Pulveris Cubebæ, ʒij.
 Balsami Peruviani, gr. vj.
 Syrupi Tolutani, fʒj.
 Extracti Glycyrrhizæ, ʒj.
 Acaciæ, q. s.

Mix, and divide into lozenges of ten grains each. These are called Spitta's lozenges, and are celebrated for the relief they afford in coryza.

Jackson's Pectoral Lozenges.

- R. Pulveris Ipecacuanhæ, gr. x.
 Antimonii Sulphureti præcipitati, gr. v.
 Morphiæ Muriatis, gr. vj.
 Pulveris Acaciæ,
 ——— Sacchari,
 ——— Extracti Glycyrrhizæ } āā ʒxj.
 Olei Sassafras, gtt. iv.
 Tincturæ Tolutanæ, fʒiv.
 Syrupi, q. s.

Fiat massa, in trochiscos cc. dividenda.

Each lozenge, weighing ten grains, contains of ipecacuanha gr. $\frac{1}{6}$, of precipitated sulphuret gr. $\frac{1}{6}$, and of the muriate of morphia gr. $\frac{1}{3}$.

Signa.—One of these to be dissolved in the mouth, when the cough is troublesome. S. JACKSON.

EXPECTORANT SIRUPS.

Sirup of Squills.

- R. Syrupi Scillæ, fʒj.

Signa.—Take a teaspoonful every three hours.

Sirup of Garlic.

- R. Syrupi Allii, fʒj.

Signa.—Give a teaspoonful to a child every three hours.

This, like the sirup of squills, is an excellent expectorant, being somewhat stimulant in its character.

Sirup of Seneka Snake-root.

- R. Syrupi Senegæ, fʒj.

Signa.—Take a teaspoonful every three hours as a stimulant expectorant.

Core's Hirc Sirup.

R. Syrupi Scillæ Compositi, f 5j.

A small teaspoonful may be given to a child of one year old with croup every quarter, half, or one hour, according to the violence of the disease, and until vomiting ensues; then discontinue it.

Sirups of Squills, Seneka, and Ipecacuanha.

R. Syrupi Scillæ,

——— Senegæ, āā f 3ss.

——— Ipecacuanhæ, f 5j.

Misce. *Signa.*—A teaspoonful every three hours.

This is a favorite combination with some physicians, as it combines the expectorant qualities of the squill and seneka, with the relaxing effect of the ipecacuanha.

Sirup of Assafetida.

R. Assafœtidæ, 3j.

Aquæ bullientis, Oj.

Sacchari, lbij.

Triturate the assafetida in a mortar, with a portion of the boiling water, until a uniform paste is formed; then gradually add the remainder of the water, strain, and add the sugar, applying a gentle heat to dissolve it. RICHARD PELTZ.

Dose, a teaspoonful every three hours, for a child five years old.

Each fluidounce contains fifteen grains of assafetida. This preparation will keep a month in the hottest weather. Its taste is less nauseous than the milk of assafetida, and altogether, it is one of the very best modes of administering the fetid drug to children in whooping-cough, I am acquainted with.

EDITOR.

Sirup of Wild Cherry Bark.

R. Syrupi Pruni Virginianæ, f 3ij.

Signa.—Take a tablespoonful every three hours.

When well prepared, this represents the tonic and sedative properties of the wild cherry, which render this bark so peculiarly applicable to the treatment of phthisis.

Jackson's Pectoral Sirup.

R. Sassafras Medullæ, ʒj.

Acaciæ, ʒj.

Aquæ, Oj.

Macera per horas decem vel duodecim, dein adde—

Sacchari, ʒxxj.

Et solve sine calore, cola, et adde—

Morphiæ Muriatis, gr. viij.

Signa.—Take from a teaspoonful to a tablespoonful, according to circumstances, every three hours.

It is intended that this preparation, when completed, shall measure two pints, and if it does not, the operator is to add the complement of water, so that each fluidounce shall contain one-fourth of a grain of the muriate of morphia. It is much employed in Philadelphia.

SAMUEL JACKSON.

EXPECTORANT MIXTURES.

Brown Mixture.

R. Pulveris Extracti Glycyrrhizæ,

———— Acaciæ, āā ʒij.

Aquæ ferventis, fʒiv.

Misce, fiat solutio, et adde—

Spiritûs Ætheris Nitrici, fʒj.

Vini Antimonii, fʒij.

Tincturæ Opii, gtt. xl. vel lx.

Signa.—A tablespoonful for a dose, *pro re nata*. In catarrh.

A preparation very similar to this is now officinal under the name of *Mistura Glycyrrhizæ Composita*.

Mixture with Sirup of Squill, &c.

R. Syrupi Scillæ, fʒss.

Tincturæ Opii Camphoratae, fʒij.

Vini Antimonii, fʒj.

Pulveris Acaciæ, ʒss.

Aquæ destillatæ, fʒiij.

Misce. *Signa.*—A teaspoonful every two or three hours.

This is more particularly designed for children.

Mixture with Seneka, Squill, Tartar Emetic, &c.

- R. Decocti Senegæ, f ̄iv.
 Syrupi Ipecacuanhæ, f ̄j.
 Oxymellis Scillæ, f ̄iij.
 Antimonii et Potassæ Tartratis, gr. j.
 Fiat mistura. *Signa*.—A teaspoonful, *pro re nata*, for children of two years of age and upwards. In croup.

Mixture with Seneka, Ammoniac, and Tolu.

- R. Senegæ contusæ, ʒij.
 Aquæ bullientis, f ̄viiij.
 Coque ad f ̄vj. Cola, et adde—
 Syrupi Tolutani, f ̄j.
 Ammoniaci, ʒss.
 Misc. —A tablespoonful every two hours. In catarrh.

Mixture of Extract of Conium, &c.

- R. Extracti Conii, ʒj.
 Tincturæ Opii Camphoratæ, f ̄ss.
 Syrupi Tolutani, f ̄ss.
 Aquæ Rosæ, f ̄iv.
 Misc. *Signa*.—From half to a whole teaspoonful may be given once in four hours to a child a year old, watching the effect.

Dr. Pearson, in his work on pertussis, recommends the above as a valuable formula.

Mixture of Almond Emulsion, &c.

- R. Misturæ Amygdalæ, f ̄v.
 Vini Ipecacuanhæ,
 Tincturæ Scillæ, āā f ̄j.
 Syrupi Tolutani, f ̄vj.
 Misc. *Signa*.—Take a tablespoonful whenever the cough is urgent.

White Linctus.

- R. Amygdalæ dulcis, No. xv.
 Tragacanthæ, ʒj.
 Syrupi, f ̄j.
 Olei Amygdalæ, f ̄ij.
 Aurantii Floris Aquæ, *Lond.* f ̄ij.
 Aquæ destillatæ, f ̄iv.
 Misc. *Signa*.—A tablespoonful for a dose, frequently through the day. In catarrh.

Mixture with Tartar Emetic and Laudanum.

R. Antimonii et Potassæ Tartratis, gr. j.

Aquæ destillatæ, f 3j.

Tincturæ Opii, ℥xx.

Misce. *Signa*.—Give a teaspoonful, or a dessertspoonful according to the age of the child, in hooping-cough.

SIR. WM. WATSON.

Mixture with Sal Ammoniac, Squills, etc.

R. Ammoniæ Murialis, 3ss.

Pulveris Acaciæ, ʒij.

———— Extracti Glycyrrhizæ, ʒiij.

Aquæ, f 3vjss.

Spiritus Ætheris Nitrici, f 3jss.

Aceti Scillæ, f 3iij.

Fiat mistura. *Signa*.—A tablespoonful every two hours.

A stimulating expectorant and alterative.

GOEDEN.

Mixture with Naphthaline.

R. Naphthalinæ, gr. xvj.

Alcoholis, q. s.

Ut fiat solutio. Dein adde—

Syrupi, f 3iv.

Misce. *Signa*.—Take a tablespoonful every two hours.

This is a stimulating expectorant, useful in chronic catarrh, humoral asthma, etc.

DUPASQUIER.

Mixture of Oil of Amber and Tolu.

R. Olei Succini rectificati, gtt. lxxx.

Acaciæ,

Sacchari, āā 3ss.

Aquæ destillatæ, f 3iv.

Tincturæ Tolutanæ, f 3ij.

Fiat mistura secundum artem. *Signa*.—A tablespoonful every two or three hours. In spasmodic cough.

Mixture with Carbonate of Potash, &c.

R. Potassæ Carbonatis, ʒij.

Vini Antimonii, f 3j.

Tincturæ Opii, gtt. xl.

Spiritus Lavandulæ Compositi, f 3ij.

Aquæ destillatæ, f 3iv.

Fiat mistura.—And let a tablespoonful be taken every hour or two hours.

Mixture with Bicarbonate of Soda, &c.

- R. Sodæ Bicarbonatis, gr. xij.
 Vini Ipecacuanhæ, gtt. xx.
 Tincturæ Opii, gtt. iv.
 Aquæ destillatæ, f ʒj.

Misce. *Signa*.—Give a teaspoonful every two or three hours.

This preparation is designed for children, and is chiefly used in pertussis.

Cochineal Mixture.

- R. Potassæ Carbonatis, ʒj.
 Pulveris Cocci, ʒss.
 Sacchari, ʒj.
 Aquæ destillatæ, f ʒiv.

Fiat solutio.—Dose for children, a teaspoonful every two or three hours.

A popular remedy in pertussis.

Mixture with Tincture of Digitalis, &c.

- R. Tincturæ Digitalis, f ʒj.
 ——— Opii, gtt. lx.
 Aquæ destillatæ, f ʒij.

Misce.—A small teaspoonful may be taken three or four times a day. In hæmoptysis and in incipient phthisis.

Mixture of Assafetida, &c.

- R. Assafœtidæ, ʒj.
 Aquæ, f ʒiv.
 Fiat mistura, et adde—
 Tincturæ Tolutanæ, f ʒss.
 ——— Opii, gtt. xx. vel xl.

A teaspoonful to a child every two hours, and a dessertspoonful (or more) as frequently to an adult.

This preparation is of great value in pertussis and other spasmodic diseases of the chest.

Tincture of Bloodroot.

- R. Tincturæ Sanguinariæ, f ʒj.
 Direct from 30 to 50 drops every three hours.

This native product is one of the most powerful medicinal agents. It is violently emetic in large doses, but in the quantity

recommended is said to produce diaphoresis and effects analogous to those of digitalis.

Its active principle is called *sanguinarina*.

Mixture of Gum Ammoniac, &c.

R. Ammoniaci, ʒj.

Aquæ destillatæ, f ʒiv.

Fiat mistura, et adde—

Oxymellis Scillæ, f ʒij.

Tincturæ Opii Camphoratæ, f ʒss.

A tablespoonful may be given every two hours.

A stimulating expectorant.

Mixture with Milk of Assafetida and Acetate of Ammonia.

R. Assafoetidæ, ʒss.

Liquoris Ammoniacæ Acetatis,

Aquæ Menthæ Viridis, āā f ʒij.

Fiat mistura.—A teaspoonful to children, and a tablespoonful to adults, in coughs, *pro re nata*.

This simple solution of assafetida is frequently employed, and is found to answer every purpose of more complicated prescriptions.

Mixture of Copaiba and Balsam Tolu, &c.

R. Copaibæ,

Balsami Tolutani,

Pulveris Acaciæ, āā ʒss.

Aquæ destillatæ, f ʒvj.

Acidi Sulphurici Aromatici, gtt. xx.

Fiat mistura.—Dose, a tablespoonful, *pro re nata*. In chronic hooping-cough. When there is hectic excitement, digitalis may be added.

Tincture of Tolu with Digitalis, &c.

R. Tincturæ Tolutanæ, f ʒiss.

Acidi Sulphurici Aromatici, f ʒss.

Tincturæ Digitalis, f ʒj.

Vini Antimonii, f ʒij.

Mellis despumati, f ʒiss.

Pulveris Extracti Glycyrrhizæ, ʒss.

Aquæ destillatæ, f ʒvj.

Fiat mistura. *Signa*.—A tablespoonful to be given according to circumstances.

Mixture with Balsam Peru, &c.

R. Balsami Peruviani, ʒss.

Acaciæ, q. s.

Aquæ Cinnamomi,

——— destillatæ, āā fʒss.

Misce. *Signa*.—To be taken at once, and repeated three or four times a day.

The different *balsams* were at one time almost proscribed from the class of expectorants, on account of their stimulating qualities. Copaiba, however, and some others of this class, are now much employed in diseases of the pulmonary mucous membrane; and the author has himself prescribed it with great advantage in the peculiar hoarseness that follows measles. ELLIS.

Tur Beer.

R. Strong Beer, one gallon.

Tar, sufficient to saturate.

Mix, and allow them to macerate for three days, with occasional agitation. Dose, one or two tablespoonfuls several times a day.

ROBINSON.

Tar Water.

R. Picis liquidæ, Oij.

Aquæ, Cong. j.

Mix them, and stir them together with a wooden rod for an hour; then let the tar subside, strain the water and keep it in well-corked bottles.—A pint, or even two pints, may be taken daily; but in some cases the dose must be greatly diminished.

Decoction of Tur and Bran, sometimes called Jew's Beer, or Wine of Tar.

Water, three quarts,

Wheat bran, one quart,

Tar, one pint,

Honey, half pint.

To be simmered together for three hours, and when cool, add a pint of brewer's yeast, and let it stand for thirty-six hours. Then bottle it. Dose, a tablespoonful morning, noon, and night.

Mixture of Gum Ammoniac and Nitric Acid, called Nitric Lac Ammoniac.

R. Ammoniaci ʒijss.

Acidi Nitrici, fʒij.

Aquæ, fʒviiij.

Pour the nitric acid and water together, then add the compound

gradually to the gum, incorporating them well in a composition mortar. A tablespoonful may be taken three or four times a day.

This preparation is strongly recommended in protracted catarrh and phthisis pulmonalis, by Dr. J. Stuart, author of a treatise on *Colds*.

Mixture of Spermaceti, &c.

- R. Cetacei, ℥ij.
 Sacchari, ℥iij.
 Tincturæ Opii Camphoratae, f℥ss.
 Aquæ, f℥viij.

Rub the spermaceti and sugar together with the yolk of an egg, then gradually mix the water.—A tablespoonful for a dose, four or five times a day. In catarrh.

Sirup of Tolu with Belladonna, &c.

- R. Extracti Belladonnæ, gr. viij. vel xij.
 Vini Ipecacuanhæ, f℥j.
 Syrupi Senegæ, f℥ss.
 ——— Tolutani, f℥iijss.

Misc. *Signa*.—To take a teaspoonful three or four times a day, freely using gum water acidulated with lemon-juice. In coughs.

Infusion of Flaxseed, &c.

- R. Lini, ℥j.
 Radicis Glycyrrhizæ contusæ, ℥ss.
 Aquæ bullientis, Oij.

Macerate for two or three hours near the fire, in a covered vessel; strain, and add lemon-juice sufficient to make it agreeable.

It may be given as a common drink in catarrh.

Decoction of Iceland Moss.

- R. Cetrariæ, ℥j.
 Aquæ, Ojss.

Boil to one pint, and strain.

This article once acquired great celebrity in coughs; but was overrated, and lost its reputation. It contains a bitter quality, which is sometimes an advantage in dyspepsia and phthisis. If freed from this, by macerating before boiling, and throwing the first liquor away, as is sometimes done, the pre-

paration is reduced to a mere demulcent, and has no advantages over other members of that class.

Prussic or Hydrocyanic Acid.

As prussic acid rises to the top of an aqueous mixture, thereby rendering the dose very uncertain, we prefer directing the patient to drop the dose each time into a solution of gum Arabic or simple sirup. It should be kept covered from the light, and not allowed to be accessible to those unacquainted with its power, as even a strong inhalation will sometimes prostrate an adult.

ELLIS & MORTON.

The physician should also know after what process the acid is made that he prescribes, and that it corresponds in strength with the officinal standard. In other words, that he is employing the *Acidum Hydrocyanicum dilutum* of the U. S. Pharmacopœia, the dose of which is one or two drops, three times a day, in mucilage, or sugar and water, and augmented one drop daily until its action on the system is developed.

EDITOR.

Pectoral Mixture.

R. Acidi Hydrocyanici diluti (one fluidram), fʒj.

Aquæ destillatæ, Oj.

Sacchari, ʒiiss.

Fiat mistura. *Signa.*—A dessertspoonful may be taken morning and evening, and gradually increased (if necessary) to six or eight spoonfuls in the twenty-four hours—shaking the vial each time.

MAGENDIE.

Mixture with Prussic Acid.

R. Acaciæ, ʒss.

Aquæ, fʒviijss.

Fiat solutio, et adde—

Syrupi Tolutani, fʒss.

Acidi Hydrocyanici diluti, gtt. xij.

Fiat mistura. *Signa.*—Take a tablespoonful every three hours.

This is “a medium dose,” which I prefer in commencing the use of so powerful a medicine.—S. G. MORTON, *Illustrations of Pulmonary Consumption*, p. 132.

In the former editions of this work, two formulæ were given for the preparation of “Magendie’s medicated solution of hydrocyanate of potassa.” The following is more simple, and corresponds with them precisely in strength and dose.

EDITOR.

Mixture of Hydrocyanate of Potash.

R. Potassii Cyanureti (seven grains), gr. viij.

Aquæ destillatæ, Oj.

Fiat solutio, et adde—

Sacchari, ʒjss.

Misce. Dose, a tablespoonful night and morning, watching its effect. Each dose contains one-fifth of a grain.

The cyanuret of potassium is said to exercise the same power over the system as prussic acid, being equally active, but with the advantage of being a more uniform and permanent medicine.

Mixture with Cyanuret of Potassium.

R. Potassii Cyanureti (half a grain), gr. ss.

Aquæ Lactucæ, fʒij.

Syrupi Altheæ, fʒj.

Fiat mistura. Dose, a tablespoonful every two hours.

MAGENDIE.

We cannot enjoin too much caution in the use of all the preparations of prussic acid, some of which are extremely variable in themselves, and all produce very unequal effects on different constitutions. It is questionable whether the practice of medicine has gained anything by their adoption into the *Materia Medica*.

Mixture with Tincture of Lobelia and Prussic Acid.

R. Tincturæ Lobeliæ, fʒj.

Acidi Hydrocyanici diluti, gtt. j. vel ij.

Misce. *Signa.*—To be taken at a dose, three or four times a day.

Highly serviceable in asthma and (in smaller doses) in whooping-cough.

LIVEZEY.

Mixture of Gum Ammoniac, &c.

R. Misturæ Ammoniaci, fʒij.

Tincturæ Castorei, fʒss.

Syrupi Tolutani, fʒss.

Tincturæ Opii, gtt. xx. vel xxx.

Aquæ Cinnamomi, fʒj.

Fiat mistura. *Signa.*—A small tablespoonful every three or four hours, in troublesome cough.

DR. MEIGS.

Mixture of Morphia and Tolu.

R. Oxy mellis Scillæ, f̄ss.

Syrupi Tolutani, f̄ss.

Morphiæ Acetatis, gr. ss. vel j.

Misce. *Signa*.—A teaspoonful may be taken every two or three hours, *pro re nata*. In phthisis, and other diseases attended with cough.

Tolu with Morphia and Antimony.

R. Mucilaginis Acaciæ, Oj.

Syrupi Tolutani, f̄ss.

Morphiæ Sulphatis, gr. i. vel ij.

Antimonii et Potassæ Tartratis, gr. i. vel ij.

Misce. *Signa*.—A tablespoonful to be given every two hours.

MORTON on Consumption, p. 123.

Decoction of Seneka Snakeroot.

R. Radicis Senegæ contusæ, ʒj.

—— Glycyrrhiæ, ʒss.

Aquæ destillatæ, Ojss.

Boil down to a pint, and strain. Dose, a tablespoonful.

This decoction is prescribed in pectoral diseases, and especially in croup. It is sometimes too pungent, and may require to be much diluted.

Infusion of Tar and Hops.

R. Picis liquidæ, ʒj.

Aquæ bullientis, Oij.

Humuli, ʒss.

Misce et cola.

Dose, a tablespoonful morning, noon, and night.

INHALATION.

Inhalation is a means of applying remedies directly to the lungs. A good instrument for this purpose is the *Inhaler* of Mudge; but as a substitute the common teapot may be used, or a funnel inverted over a bowl. The substances usually selected for inhalation are, the steam of warm water; the steam of vinegar, produced by pouring it into boiling water; and various other articles next to be enumerated.

Inhalation of Balsam of Tolu.

R. Balsami Tolutani, ʒj.

Aquæ bullientis, Oj.

Misce.—To be inhaled.

Ether is also used alone, or impregnated. When saturated with cicuta, and inhaled from a wineglass, it is adapted to protracted catarrh, pneumonia, consumption, &c. Dr. Morton advises Dr. Pearson's formula, which consists in keeping a dram of powdered cicuta in an ounce of ether for a week. A teaspoonful or two of this impregnated fluid may be inhaled at a time, and repeated thrice per diem.—MORTON *on Consumption*, p. 136. The latter practitioner has more recently recommended the following formula:—

Ethereal Tincture of Iodine and Cicuta for Inhalation.

R. Iodinii, gr. viij.

Pulveris Conii, gr. xvj.

Ætheris, fʒij.

Macerate the ingredients together for 24 hours or more. Half a teaspoonful at a time is to be breathed from a wineglass grasped in the warm hand, and repeated twice or thrice a day. If it does not affect the head, the quantity inhaled may be gradually increased to a teaspoonful. S. G. MORTON.

The following is Dr. Scudamore's recipe for Iodine inhalation:—

R. Iodinii,

Potassii Iodidi, āā gr. vj.

Aquæ destillatæ, fʒv. ʒvj.

Alcoholis, fʒj.

Mix, and use for inhalation.

I usually commence with a dram of this mixture, proceeding gradually to the extent of half an ounce (rarely more), putting two-thirds the dose for the first half of the time (10, 15, or 20 minutes), and the other third for the remainder, always adding thirty minims of a saturated tincture of conium, with an increase if the cough be very irritable. Occasionally, I add some saturated tincture of ipecacuanha; and when the respiration is spasmodically affected, some ethereal tincture of lobelia inflata.

SCUDAMORE.

Stramonium is particularly serviceable in asthma and consumption. The root is cut, dried, and smoked from a common pipe.

Fumigation with *tar* and *turpentine* is also much in use. It

is usually applied by placing the tar on a dish over coals, and suffering it slowly to evaporate; but a far neater and more effectual mode is that practised by Sir Alexander Crichton, viz: "An ounce of potash is added to every pound of tar, in order that the latter may be deprived of its pyroligneous acid. The two ingredients being well mixed, should be first boiled for a few minutes in the open air, in order to disengage any impurities, and then be kept at a simmer in the room of the patient. This is effected by putting the mixture in an iron vessel, and placing the latter over a spirit-lamp, or some analogous contrivance. In this way, not only a chamber, but an entire house, is speedily pervaded by a most agreeable vapor, which, although it may at first excite some disposition to cough, both in healthy and sick persons, very soon, in a majority of cases, allays this symptom, and with it a great proportion of the patient's distress."—MORTON, *ut supra*.

The fumes of *rosin* were much employed by the late Dr. Parrish, of this city; he directed it to be put on a hot shovel, or coals, and allow the patient gradually to inhale it. To some persons it is very agreeable, while the lungs of others cannot tolerate it.

In some cases of croup, the inhalation of ether or Hoffman's anodyne, will often procure prompt relief, and gain time for the application of general remedies.

Chlorine gas has also been much used by inhalation, not pure, however, but mixed or diluted with atmospheric air. For this purpose a little acid of any kind may be dropped gently on a mixture of chloride of lime, by which means the chlorine is gradually disengaged, and is inhaled as it rises; or the air of a room may be pervaded with it in like manner. It has been much commended in bronchitis and the kindred affections, but is getting into disuse from its manifest inefficiency.¹

¹ *Fumigation of Chlorine.*

R. Sodii Chloridi, ℥ij.
Oxidi Manganesii, ℥v.
Aque, ℥ij.
Acidi Sulphurici, ℥j.

Mix the salt and manganese, and add at proper intervals the acid and water, mingled. When used in the wards of hospitals or other infected apartments, the patients should be previously removed. For the method of preparing the different chlorides used as disinfecting agents, see *Journal of the Philadelphia College of Pharmacy*, vol. i. No. 4.

CLASS VII.

EMMENAGOGUES.

EMMENAGOGUES are those medicines which promote the menstrual discharge, or restore it when entirely suppressed. Although by some authors the specific action of these medicines on the uterus has been questioned, if not denied, yet by others, of equal authority, it has been fully acknowledged.

It must, however, be observed, that amenorrhœa occurs under opposite states of the system; and when general *debility* appears to be the cause of the local disease, it may be advisable to accompany the emmenagogue with an active course of tonics. While, on the contrary, if the obstruction be the effect of a plethoric habit, bleeding, cathartics, &c., are essential preliminaries to the successful exhibition of a specific remedy.

POWDERS AND PILLS.

Powder of Hiera picra.

R. Pulveris Aloës et Canellæ, ʒj.

Divide in chartulas vj.—One, three times a day.

Powder of Aloes, Canella Alba, &c.

R. Pulveris Aloës, ʒjss.

—— Canellæ, gr. xvij.

—— Serpentariæ, gr. vj.—xij.

Misce, divide in pulveres vj. *Signa.*—One, every three or four hours, in sirup.

This compound, another form of the *hiera picra* of the shops, forms a very good purgative, and is usually employed as an emmenagogue.

Powder of Madder.

R. Pulveris Rubiæ, ʒss.

Signa.—A small teaspoonful to be taken morning, noon, and night, in sugar and water.

There are three or four qualities of madder, which circumstance may account for its frequent inertness.

Powder with Savin, &c.

R. Pulveris Sabinæ,

———— Zingiberis, āā ʒj.

———— Potassæ Sulphatis, ʒij.

Fiant pulveres vj. *Signa*.—One to be taken twice a day.

Powder with Savin and Cantharides.

R. Pulveris Sabinæ, ʒj.

———— Cantharidis, gr. ij.

Misce, et divide in pulveres iv. *Signa*.—One to be taken every night on going to bed, watching its effect.

Powder of Ergot.

R. Pulveris Ergotæ, ʒij.

Divide in chartulas iv.—Of which one may be taken morning and night, in sugar and water.

Drs. Chapman and Dorsey have placed this substance under the head of Emmenagogues, where it seems properly to belong.

Powder with Savin and Borax.

R. Pulveris Sabinæ,

———— Zingiberis, āā gr. vij.

Sodæ Boratis, gr. xv.

Fiat pulvis.

This powder is to be taken twice a day, in amenorrhœa with a languid pulse.

THOMSON.

Pills of Myrrh, Sulphate of Iron, &c.

R. Myrrhæ, ʒij.

Potassæ Carbonatis, ʒss.

Ferri Sulphatis, ʒij.

Sâponis, ʒss.

Fiat massa, in pilulas xl. dividenda.—Of which two may be taken morning, noon, and night. In suppression of the catamenia.

Compound Pills of Iron.

- R. Myrrhæ (in massâ), ʒij.
 Potassæ Carbonatis, ʒj.
 Tere simul in pulverem, dein adde—
 Pulveris Ferri Sulphatis, ʒj.
 Sacchari Communis, ʒj.
 Fiat massa, et divide in pilulas lxxx.

No fluid of any kind is to be used, the brown sugar giving a proper consistence.

Two or three pills are given three times a day in amenorrhœa and chlorosis.

HULSE.

Hooper's Pills.

- R. Ferri Sulphatis, ʒj.
 Pulveris Jalapæ, gr. xv.
 ——— Aloës et Canellæ, ʒj.
 Myrrhæ, gr. viij.
 Syrupi, q. s.
 Fiat massa, et divide in pilulas l.

Of these pills two or three are taken at bedtime, for several nights successively. They purge smartly, and I presume it is by virtue of this property they operate so beneficially as an emmenagogue.

CHAPMAN.

Pills with Digitalis, Myrrh, &c.

- R. Pulveris Digitalis, gr. x.
 ——— Myrrhæ, gr. xx.
 Ferri Sulphatis, gr. x.
 Syrupi, q. s.
 Fiat massa, et divide in pilulas x.
Signa.—Take one pill three times daily.

Dr. Ashwell recommends these pills in amenorrhœa combined with epilepsy.

BRIGHT.

Pills of Aloes, Sulphate of Iron, &c.

- R. Ferri Sulphatis exsiccati, ʒj.
 Pulveris Aloës, ʒij.
 ——— Caryophylli, gr. v.
 Terebinthinæ Venetæ, q. s. ut fiant pilulæ xx. One to be taken three times a day.

DECOCTIONS, INFUSIONS, &c.

Decoction of Seneka.

R. Radicis Senegæ contusæ, ʒj.

Aquæ, Oj.

Aurantii Corticis, ʒss.

Let the seneka simmer over a slow fire until the fluid be reduced one-third; add the orange-peel, cover until cool, and then strain.

Dr. Chapman observes: "My rule, in the administration of the medicine in these cases (of amenorrhœa), is to give about four ounces of the decoction, more or less, during the day, according to the circumstances of the case. But at the time when the menstrual effort is expected to be made, and until the discharge is actually induced, I push the dose as far as the stomach will allow, having given as much as two ounces every hour."—*Therapeutics*, vol. ii. p. 38.

Infusion of Ergot.

R. Ergotæ contusæ, ʒij.

Aquæ bullientis, fʒiv.

Fiat infusum.—To be given by tablespoonfuls, in protracted parturition occasioned by deficiency of contractile power in the uterus.

For the same purpose the powder may also be given in doses of ʒj., as well to assist in arresting uterine hemorrhage as to promote the expulsion of the placenta.

Decoction of Madder.

R. Pulveris Rubiæ, ʒj.

Aquæ bullientis, Oj.

Simmer for fifteen minutes, and add—

Caryophylli contusi, ʒj.

Signa.—When cool, strain and give a wineglassful every three hours, a short time before the expected return of the menstrual discharge.—See *Dewees on Females*, p. 74.

Solution of Iodide of Potassium.

R. Potassii Iodidi, ʒj.

Aquæ destillatæ, fʒj.

Fiat solutio. *Signa.*—Dose, five to twenty drops.

Injection of Aqua Ammoniac, &c.

R. Liquoris Ammoniac, gtt. xx.

Lactis, f3j.

Misce. *Signa*.—This quantity to be thrown into the vagina three or four times a day, in amenorrhœa.

Mixture of Camphor, &c.

R. Camphoræ, ʒj.

Alcoholis, q. s. ut fiat pulvis, dein adde—

Pulveris Acaciæ, ʒj.

Sacchari, ʒj.

Aquæ Cinnamomi, f3j.

Fiat mistura.

This mixture would more properly appear under the head of Narcotics, but as Dr. Dewees considers it particularly efficacious in dysmenorrhœa, we have given it a place with Emmenagogues. He directs "one-half the above draught to be given the instant pain is experienced, and if not relieved in an hour or two, the remainder to be given."—*Syst. of Midwifery*, p. 158.

Sirup of Ergot.

R. Ergotæ, gr. xx.

Extracti Opii, gr. ʒ.

Syrupi, f3viij.

Misce. Dose, two tablespoonfuls *pro re nata*, in engorgement of the uterus. LISFRANC.

Mixture with Cantharides, Hartshorn, &c.

R. Tincturæ Gentianæ Compositæ, f3jss.

———— Cantharidis, f3j.

Spiritûs Ammoniac Aromatici, f3iij.

Misce. *Signa*.—Give a teaspoonful three times daily. Useful in chlorosis, and hysterical depression. ASHWELL.

TINCTURES.

Tincture of Ergot.

R. Ergotæ, ʒijss.

Alcoholis diluti, Oj.

Macerate for fourteen days, and strain. This is the formula used in Guy's Hospital. Dose, twenty drops to a dram.

Tincture of Black Hellebore.

R. Tincturæ Hellebori, f̄ssj.

Signa.—Half a teaspoonful three times a day, gradually increasing the quantity and watching its effect.

Tincture of Black Hellebore, Myrrh, &c.

R. Tincturæ Hellebori, f̄ss.

———— Myrrhæ, f̄ssj.

———— Cantharidis, f̄ssij.

Fiat mistura.—Dose, thirty drops, three times a day, in a little sugar and water.

Dewees's Tincture of Guaiacum.

R. Pulveris Guaiaci, ʒiv.

Sodæ vel Potassæ Carbonatis, ʒiss.

Pulveris Pimentæ, ʒj.

Alcoholis diluti, Oj.

Misce.—“Digest for a few days. The volatile spirit of ammonia to be added *pro re nata*, in the proportion of a dram or two, to every four ounces of tincture.”

The above formula is different from that adopted in the Pharmacopœias, for which reason we have inserted it here. Dr. Dewees remarks that, after nearly thirty-five years' experience with this tincture, he has never known it to fail in restoring the *suppressed catamenia*, when the suppression was an *idiopathic disease*. He gives it morning, noon, and night, in doses of a teaspoonful in a little sweetened milk; and sometimes in a little Madeira, Sherry, or Teneriffe wine. It is occasionally necessary to continue the medicine for some weeks, on which occasion, the dose must be judiciously augmented.—*Syst. of Midwifery*, p. 149.

Volatile Tincture of Guaiacum and Copaiba.

R. Tincturæ Guaiaci Ammoniatae, f̄ssj.

Copaibæ, f̄ss.

Misce. *Signa.*—A teaspoonful two or three times a day.

In cases of amenorrhœa connected with general debility.

Saturated Tincture of Iodine.

R. Iodinii, ʒij.

Alcoholis, f̄ssj.

Spiritûs Lavandulæ Compositi, f̄ssij.

Fiat tinctura. *Signa.*—From five to ten drops in sweetened water twice a day, gradually increasing it.

We are informed by M. Durand, apothecary of this place, that the above is a saturated tincture. It is prescribed in amenorrhœa, gonorrhœa, leucorrhœa, &c.

From its great activity it requires to be cautiously administered.

Tincture of Cantharides.

R. Tincturæ Cantharidis, f ʒss.

Signa.—Take fifteen drops three times a day, in sugar and water.

The late Dr. Dewees was much in the habit of prescribing this article in leucorrhœa, and suppression of the menses; increasing the dose a few drops every third day, until symptoms of strangury were manifested, when he desisted altogether.

Muriated Tincture of Iron, &c.

R. Tincturæ Ferri Chloridi,

———— Aloës et Myrrhæ, āā f ʒss.

———— Castorei, f ʒij.

Misce. *Signa.*—Take ten to thirty drops three times a day in a wineglassful of cold chamomile tea.

Tincture of Hiera Picra.

R. Pulveris Aloës et Canellæ, ʒjss.

Spiritûs Vini Gallici, Oj.

Macera per dies decem et cola.

Signa.—A teaspoonful may be taken three times a day in sugar and water.

A popular and most efficient remedy in amenorrhœa.

Compound Tincture of Aloes.

R. Tincturæ Aloës et Myrrhæ, f ʒj.

Dose, a teaspoonful *pro re nata*.

Emmenagogue Suppository.

R. Olei Cacao, ʒij.

Pulveris Aloës, gr. ij.

Castorei, gr. viij.

Pulveris Assafoetidæ, gr. viij.

Fiat suppositorium.

One to be employed daily for several days at the expected period of the menses.

RICHARD.

CLASS VIII.

ANTHELMINTICS.

ANTHELMINTICS are medicines employed to destroy and bring away worms from the alimentary canal. They are also termed Vermifuges.

The only unequivocal proof of the existence of worms in the alimentary canal is the ejection of some of them *per anum*, or, as sometimes happens, by the mouth. When, in such cases, vermifuges are resorted to, they must be persevered in for several days, with the occasional intervention of an active cathartic.

POWDERS AND PILLS.

Calomel with Gamboge.

R. Hydrargyri Chloridi Mitis, gr. v.

Pulveris Gambogiæ, gr. iij. vel vj.

Misce.—For an adult. To be taken in sirup early in the morning.

Gamboge is sometimes given in doses of fifteen or twenty grains for the tape-worm, and is highly spoken of in such cases.

Calomel with Pink-root.

R. Hydrargyri Chloridi Mitis, gr. iv.

Pulveris Spigeliæ, gr. x.

Misce. *Signa*.—To be taken two mornings in succession, and on the afternoon of the second day, followed by a dose of castor-oil, or senna and manna.—For children upwards of four years old.

Pink-root with Savine, &c.

- R. Pulveris Spigeliæ,
 ——— Sennæ, āā ʒij.
 ——— Sabinæ, gr. xij.

Fiant chartulæ vj.—One every morning till three are taken, then give a dose of castor-oil; if the effect is not produced, the remainder may be exhibited after the same manner.

Common Salt with Cochineal.

- R. Sodii Chloridi, ʒijj.
 Cocci pulverizati, gr. xv.

Fiant pulveres vj. One every morning, following the last dose with some purgative medicine.

Dr. Rush speaks in high terms of this preparation.

Powder of Santonine.

- R. Santonin, gr. vj.
 Sacchari Lactis, gr. xv.

Misce, et divide in pulveres vj.

Signa.—Give one powder night and morning to a child five years old, for lumbrici.

“In many cases the worms are passed, says Dr. Wells, on the following morning, but not unfrequently it is necessary to give a second dose on the succeeding evening.”

A case of poisoning from three grains in two doses, occurred in a child four years old. (*N. Y. Journ. Pharmacy*, vol. i. p. 16.) Hence, it is necessary to exercise much caution in administering this powerful medicine.

EDITOR.

Powder of Male Fern.

- R. Pulveris Filicis Maris, ʒj.

The following is Nouffer's plan of treating cases of tape-worm:—

“Give of the powder of the Male Fern root, from one to three drams, in a large cupful of any distilled water, early in the morning. If the stomach should reject it, the powder must be repeated again, as soon as the sickness has gone off, in the same quantity. When it has been upon the stomach about two hours, a strong cathartic bolus made of calomel and gamboge (eight or ten grs. of the last, and five or six grs. of the first) must be taken, and worked off with a draught of green tea every now and then; and its operation may be still farther promoted, if necessary, by half an ounce, or ʒvj. of salts, dis-

solved in warm water. Till the worm comes away, which generally happens on the same day, the patient is to take nothing but broth. If, however, the worm should not be discharged, either wholly or in part, the powder is to be repeated with the same regimen the next day."

It has been suggested, and with some probability, that the purge directed to follow this medicine is the most efficient of the two.

To avoid the large doses necessarily given when the powder is used, an ethereal extract has been prepared, which represents all the virtues of the root in a small bulk. It is generally called the oil of fern (*Oleum Filicis Maris*), and is given in the dose of fifteen to twenty grains. It is made by exhausting 5j. of the freshly dried root by f5viij. of ether, macerating, and distilling off the ether.—DUNGLISON, *New Rem.* p. 382, 6th ed.

Oil, or Extract of Fern.

R. Olei Filicis Maris, 3ss.

Mellis Rosæ, ʒss.

Misce. *Signa*.—Take one-half at bedtime, and the remainder early in the morning, on an empty stomach.

If the worm does not come away in 6 or 8 hours after the last dose, administer some mild purgative. RADIUS.

Carbonate of Iron.

R. Ferri Subcarbonatis, 3j.

Divide in chartulas iij. One powder to be taken before breakfast.

Of this remedy, the late Professor Rush had a very favorable opinion. He gave it in doses of from 5 gr. to 3ss. to children of from one year old to ten, and sometimes in half-ounce doses, with safety and success. It is chiefly used in tape-worm.—See *Observ. and Inquiries*, p. 243.

Pills of Venice Turpentine, Calomel, &c.

R. Terebinthinae Venetae, 3j.

Extracti Hyoscyami, gr. iv.

Hydrargyri Chloridi Mitis, gr. viij.

Fiat massa, et divide in pilulas singulas, gr. ij.

Of these, four are to be taken every three hours, and during the entire period, the patient is to take only weak broth, tea, or coffee.

They are strongly recommended by some of the German practitioners in tape-worm.

INFUSIONS, DECOCTIONS, &c.

Infusion of Pink-root, &c., vulgo, Worm-tea.

R. Spigeliæ, ʒss.

Sennæ, ʒij.

Mannæ, ʒj.

Sabina, ʒij.

Fœniculi, ʒij.

Aquæ bullientis, Oj.

Fiat infusum. Dose, a tablespoonful three times daily for three days, for a child four or five years old. If it purge too actively, it should be discontinued.

Infusion of Pink-root.

R. Spigeliæ, ʒj.

Aquæ bullientis, Oj.

Fiat infusum. Of which a child may take a tablespoonful at a dose; an adult from ʒj. to ʒiv.

The spigelia sometimes produces violent effects in children, whence the necessity of caution in its use.

Decoction of the Bark of Pomegranate Root.

R. Granati Radicis Corticis, ʒjss.

Aquæ, Ojss.

To be boiled down to one-half, and given in doses of an ounce several times a day.

“The efficacy of the bark of the root of the pomegranate, as a remedy for the tape-worm, is now well established in India.”

PARIS, vol. ii. p. 173.

Infusion of Kousso.

R. Florum Brayeræ Anthelminticæ, ʒss.

Aquæ Bullientis, fʒx.

Macera per horam dimidiam. *Signa.*—The tea and flowers constitute a full dose. To be used for tape-worm.

The only preparation necessary is, that the last meal of the preceding evening should be slight. The previous evacuation of the bowels by a mild purgative or a lavement, is also desirable. The mode of administering the remedy is as follows: “A little lemon-juice is to be swallowed, and the infusion being stirred up, the whole is taken, liquid and powder, at two or three draughts, at short intervals, being washed down by cold

water and lemon-juice. To promote the operation, tea (without sugar or milk) may be taken. In three or four hours, if the remedy has not operated, a dose of castor-oil, or a saline purgative, should be administered." JON. PEREIRA.

ELECTUARIES, MIXTURES, &c.

Tin Electuary.

R. Pulveris Stanni, ʒvj. vel ʒj.

Sacchari Fæcis, vel Syrupi, fʒiv.

Misce.—Two or three tablespoonfuls may be given early in the morning; to be repeated until the whole be taken, and then followed by an active cathartic.

The efficacy of this substance is said to be increased by combination with sulphur, by which means sulphuretted hydrogen is generated, and the worm destroyed by the offensive gas. Larger doses are sometimes prescribed.

Cowhage Electuary.

R. Spicularum Mucunæ, ʒj.

Mellis liquidi, q. s. ut fiat electuarium. Dose, a teaspoonful to a child two or three years old; to be given before breakfast in the mornings of three successive days, and then followed by a brisk cathartic. In lumbrici.

Bolus with Calomel, Semen Contra, and Camphor.

R. Hydrargyri Chloridi Mitis, gr. ij.

Pulveris Artemisiæ Santonicæ, gr. viij.

Camphoræ, gr. ij. ad vj.

Syrupi, q. s. ut fiat bolus.

To be repeated *pro re nata* in children of four or six years of age.

RATIER'S *Formulary*.

Mixture with Oil of Wormseed.

R. Olei Chenopodii, fʒjss.

— Ricini, ʒiij.

— Anisi, gtt. x.

Misce, et adde—

Syrupi Rhei vel Sennæ, fʒj.

Fiat mistura. *Signa*.—Shake the bottle well before using.

Give a teaspoonful night and morning to a child two years old.

Mixture of Wormseed Oil.

R. Olei Chenopodii, fʒj.

Sacchari,

Acaciæ, āā ʒss.

Aquæ Menthæ Piperitæ, fʒijss.

Misce, secundum artem. A teaspoonful of this preparation may be given four times a day for two days in succession, and then purged off with castor-oil or salts. For children of three to eight years old.

This is a vermifuge of considerable reputation in this country. It is generally used in *lumbri*ci and *ascari*des; though we have heard of its having been successfully given for *tænia*, but in much larger doses.

Olive Oil and Ammonia.

R. Olei Olivæ, fʒviij.

Spiritûs Ammoniæ Aromatici, fʒij.

Dose, three tablespoonfuls night and morning.

This mixture is highly spoken of by some authors, and may be given in much larger quantities; if it does not purge, it should be followed by some aloetic or other cathartic preparation.

Extract of Pink-root and Senna.

R. Extracti Spigeliæ et Sennæ Fluidi, fʒij.

Signa.—Give a child two years old a teaspoonful night and morning, until purging takes place.

This preparation, originally devised by Prof. Procter, has been used successfully in Philadelphia, for the last ten years, in the treatment of *lumbri*ci.

Spirit of Turpentine

Is recommended for tape worm, in doses of ʒss. to ʒij. daily, taken in honey.

Dr. Chapman speaks highly of this remedy; but it sometimes causes strangury, and requires caution.

Spirit of Turpentine, Ether, &c.

R. Olei Terebinthinæ, fʒss.

Ætheris, fʒij.

Pulveris Acaciæ, ʒss.

Aquæ, Oj.

Misce. *Signa.*—Two tablespoonfuls to be taken morning and evening. In *tænia*.

DR. FRANK.

The effect of turpentine requires attention, for in some constitutions it causes violent strangury, in defiance of every precaution. Patients should be put on their guard against this contingency.

CLYSTERS.

Aloetic Enema.

R. Aloës, ʒj. vel ij.

Lactis, Oj.

Fiat enema.—The half of this may be given to a child eight or ten years old. In ascarides.

Enema of Camphor and Sweet Oil.

R. Camphoræ, ʒj.

Olei Olivæ, ʒij.

Misce pro enema.

This preparation is particularly adapted to *ascarides*, and should be freely thrown up the rectum for three or four successive nights. No remedy is more effectual than this in relieving the intolerable itching caused by those worms.

Enemata are the most certain of all remedies in *ascarides*; they may be made of various substances, such as solution of assafetida, of common salt, hepar sulphuris, pure olive oil, soap, aloes, &c. &c. For children, ʒss. of aloes should be dissolved in half a gill of water; to this add a tablespoonful of sweet oil, and administer the half of the mixture on going to bed. The other half the next night, if necessary.

CLASS IX.

STIMULANTS.

IN the use of stimulants, it is for the most part advisable to begin with small doses, and augment them as circumstances may require. There are cases, however, in which it is necessary to give them plentifully *ab initio*.

It will be often requisite to change not only the stimulating substance, but the part of the body to which it is applied ; thus, when the stomach fails, the rectum and skin may be acted upon with advantage.

PILLS, POWDERS, &c.

Pills of Camphor, Cantharides, &c.

R. Pulveris Cantharidis, gr. xvij.

———— Opii,

———— Camphoræ, āā gr. xxxvj.

Confectionis Rosæ, q. s.

Ut fiant pilulæ xxxvj.—Of which one may be taken every night on going to bed. In impotency from general debility.

Pills of Turpentine and Guaiacum.

R. Pulveris Guaiaci, ʒj.

Terebinthinæ Venetæ, q. s.

Ut fiant pilulæ xv. *Signa*.—One, three times a day.

These pills have a powerful effect upon the urinary organs. They are employed in gleet and leucorrhœa, but frequently produce strangury.

Pills of Cayenne Pepper.

R. Pulveris Capsici, ʒj.

Micæ Panis,

Aquæ destillatæ, āā q. s.

Ut fiant pilulæ xij. *Signa.*—One, three or four times a day. In debility of the stomach, from intemperance.

Aromatic Pills.

R. Ammonia Carbonatis, }
 Capsici, } āā ʒj.
 Caryophylli, }
 Macis, }

Olei Cari, gtt. v.

Extracti Gentianæ, gr. xij.

Syrupi, q. s.

Ut fiant pilulæ xx.—One every two hours in gout of the stomach. Dr. PARRISH.

Antiseptic Pills.

R. Camphoræ,

Potassæ Nitratis,

Pulveris Acaciæ, āā ʒj.

Fiat massa, in pilulas xv. dividenda.

Signa.—Dose, two pills every hour or two hours.

Creasote Pills.

R. Creasoti, gtt. x.

Pulveris Rhei, ʒj.

Extracti Gentianæ, gr. x.

Fiat massa in pilulas x. dividenda.

Signa.—Take one pill three times daily.

Pills of Valerianate of Quinia.

R. Quinæ Valerianatis, *Dub.* gr. xij.

Pulveris Tragacanthæ, gr. vj.

Aqua, q. s. ut fiat massa in pilulas viij. dividenda.

Signa.—Give one pill every hour or two in hemicrania.

EDITOR.

MIXTURES AND INFUSIONS.

Mixture of Carbonate of Ammonia, &c.

R. Ammoniæ Carbonatis, ʒiss.

Pulveris Sacchari,

———— Acaciæ, āā ʒiss.

Spiritûs Lavandulæ Compositi, f ʒij.

Aquæ destillatæ, vel Menthæ Viridis, f ʒiv.

Fiat mistura. *Signa*.—A tablespoonful for a dose, every two or three hours. If desirable, tinct. opii, gtt. xl. vel l. may be added to the mixture.

Draught with Valerian and Carbonate of Ammonia.

R. Olei Valerianæ, gtt. iij.

Ammoniæ Carbonatis, gr. x.

Aquæ Cinnamomi, f ʒjss.

Syrupi, f ʒss.

Fiat haustus.—To be taken every fourth hour. In nervous headache, and depression of spirits.

Mixture with Oil of Turpentine.

R. Olei Terebinthinæ, gtt. cxx.

Pulveris Acaciæ,

Sacchari, āā ʒij.

Tincturæ Opii, gtt. lx.

Spiritûs Lavandulæ Compositi, f ʒij.

Aquæ Menthæ Viridis, f ʒv.

Fiat mistura. *Signa*.—A tablespoonful for a dose every two hours.

This remedy has gained great reputation in some of the low forms of fever. The author once employed it in the dose prescribed above, in passive hemorrhage from the bowels with complete success; and in hemorrhage from the bladder, resulting from low fever, it has been given with equal advantage.

Draught with Powdered Ginger, &c.

R. Pulveris Zingiberis, gr. xv.

Ammoniæ Carbonatis, gr. viij.

Spiritûs Cinnamomi, *Lond.* f ʒij.

Aquæ, f ʒiss.

Misce.—To be taken at once. In gout or cramp of the stomach.

Camphor Mixture.

R. Camphoræ, ʒj.
 Pulveris Acaciæ,
 ——— Sacchari, āā ʒiss.
 Tincturæ Opii, gtt. xl.
 Aquæ Menthæ Viridis, f ʒiv.

Fiat mistura. *Signa*.—A tablespoonful to be given every two hours.

In reducing camphor to powder (which must be done before it can be formed into a mixture), it is necessary to add to it a few drops of alcohol; or, what is still better, to add f ʒss. of the tincture of myrrh. This renders the camphor much more miscible with water, and does not detract from the value of the preparation.

Camphor in Milk.

R. Camphoræ, ʒj.
 Lactis bullientis, f ʒiv.

Fiat solutio.—This preparation may be taken in the manner directed for the former one.

Camphor Julep.

R. Camphoræ, ʒj.
 Aquæ bullientis, f ʒviiij.

Fiat mistura.—This preparation should be set aside in a covered vessel for half an hour, and then strained. A tablespoonful for a dose *pro re nata*.

Many of the older practitioners are in the habit of using a hot camphor tea, in dysmenorrhœa, under an impression that its virtues are better developed. It certainly is stronger hot than cold, but in neither case is it equal to the officinal camphor water (aqua camphoræ) which contains three grains to the fluid-ounce when properly prepared.

EDITOR.

Camphor with Myrrh.

R. Camphoræ pulverizatæ, ʒj.
 Pulveris Myrrhæ, ʒss.
 ——— Sacchari, ʒij.
 Aquæ, f ʒvj.

Misce. *Signa*.—A tablespoonful to be given every two hours.

Mixture with Camphor and Chloroform.

- R. Camphoræ, ʒij.
 Chloroformi, ʒj.
 Vitellus Unius Ovi,
 Aquæ, f ʒiv.

Fiat mistura.

This mixture affords us the means of administering large doses of camphor, in as concentrated a form as it would be safe to give a local irritant. It can be diluted with water without precipitation. Each teaspoonful contains four grains.

SMITH.

Camphorated Enema.

- R. Enematis Communis, Oij. (See page 62.)
 Camphoræ, ʒij.
 Vitellus Unius Ovi.

Misce. One-fourth part to be thrown up the rectum in the treatment of putrid fevers, *pro re nata*. M. CHAUSSIER.

Mixture of Camphorated Opium and Guaiacum.

- R. Tincturæ Opii Camphoratæ,
 ——— Guaiaci Ammoniatæ, āā f ʒj.

Fiat mistura. *Signa*.—Two teaspoonfuls every two hours. In misplaced and retrocedent gout.

Aromatic Spirit of Hartshorn.

- R. Spiritûs Ammoniaë Aromatici, f ʒij.

Signa.—Twenty drops to be given in a wineglassful of water, *pro re nata*. In sudden exhaustion from any cause.

Yeast Mixture.

- R. Cerevisiæ Fermenti, f ʒx.
 Camphoræ, ʒss.
 Spiritûs Ætheris Nitrici, f ʒss.

Fiat mistura. *Signa*.—Take a tablespoonful every hour or two hours.

This was extensively used in the late typhus epidemic in Ireland, as a stimulant and antiseptic, and with the happiest results.

LAMPREY.

Substitute for the Fetid Spirits of Ammonia.

- R. Spiritûs Ammoniaë, f ʒj.
 Tincturæ Assafoetidæ, f ʒvij.

Misce. *Signa*.—Give a teaspoonful, in sweetened water, every four hours.

An excellent stimulus in nervous debility and hysteria.

Mixture with Dippel's Animal Oil.

R. Olei Cornu Cervi, fʒj.

Spiritûs Ætheris Compositi, fʒiij.

Misce. *Signa.*—Take twenty or thirty drops every three hours.

Used in chorea, catalepsy, &c.

RICHTER.

Emulsion of Phosphorus.

R. Phosphori puri, gr. ij.

Mucilaginis Acaciæ, q. s.

Let these articles be carefully and effectually triturated in a mortar, and add gradually as much distilled water as will make an emulsion of ʒvj.; then add—

Syrupi, fʒj.

Spiritûs Ætheris Compositi, gtt. xxx.

Fiat mistura.—Of which the usual dose is a tablespoonful every two, three, or four hours.

This preparation was originally made by the celebrated Hufeland, and is now taken from Dr. Chapman's *Therapeutics*, p. 176, vol. ii.

As the violence of the remedy, and its very great power, are generally known to physicians, we need hardly repeat the necessity there is for the greatest caution in its administration.

The subjoined is one of the many formulæ adopted in Europe.

Lobstein's Phosphorated Ether.

R. Phosphori, gr. ij.

Solve in Ætheris, fʒss.

Tincturæ Olei Menthæ Piperitæ, fʒss.

Signa.—Take from two to six drops in sugar every three hours.

The bottle should be carefully stoppered, and covered with dark paper. Used in epilepsy, paralysis, and amaurosis.

Foy.

Tincture of Arnica.

R. Arnicæ Florum, ʒjss.

Alcoholis, Oj.

Macera per dies quatuordecim et cola.

Of this tincture, Mr. Wilde gives fifteen drops in a tablespoonful of the infusion of arnica, and a little of some cordial tincture, three times a day, for the relief of tinnitus aurium. After a few days, the dose should be increased one or two drops daily, until it reaches thirty drops. It is useful as a stimulant in atonic gout.

WILDE.

Mixture with Chlorinated Lime.

R. Calcis Chlorinatæ, ʒj.

Aquæ, f ʒiij.

Solve et cola. Dein adde—

Syrupi Zingiberis, f ʒj.

Fiat mistura. *Signa*.—Take a tablespoonful every three hours. Chlorinated lime is stimulant and antiseptic, and has been used beneficially as such in typhoid, typhus, and scarlet fevers.

Linctus with Oil of Turpentine.

R. Olei Terebinthinæ, f ʒij.

Mellis optimi, f ʒj.

Fiat linctus.—Dose, a teaspoonful night and morning, with a draught of warm tea. In sciatica.

Infusion of Cayenne Pepper.

R. Pulveris Capsici, ʒiss.

Aquæ bullientis, Oss.

Fiat infusum. *Signa*.—A tablespoonful for a dose in severe colic, or gout in the stomach.

Infusion of Cloves.

R. Caryophylli contusi, ʒj.

Aquæ bullientis, Oss.

Fiat infusum. *Signa*.—To be employed in the same cases, and in similar doses as the former.

Mixture of Cubebs, &c.

R. Pulveris Cubebæ, ʒij.

Sodæ Carbonatis, ʒss.

Mucilaginis Acaciæ, f ʒvj.

Aquæ Menthæ Viridis, f ʒvj.

Fiat mistura. *Signa*.—Take a tablespoonful every hour and a half.

Mixture of Oxide of Bismuth, Cubebs, &c.

R. Pulveris Cubebæ, ʒij.

Bismuthi Subnitratis, ʒss.

Mucilaginis Acaciæ, f ʒss.

Syrupi, f ʒvj.

Aquæ, f ʒvj.

Fiat mistura. *Signa*.—Take a tablespoonful four times a day.

We have extracted the two preceding prescriptions from a

paper of Mr. John Fosbrooke, surgeon, Chitttenham. See *Medical Recorder*, for April, 1825.

They are recommended, with other similar compounds, in various affections of the mucous membranes, viz: of the bowels, bladder, trachea, &c.

Mustard Whey.

R. Lactis Vaccinæ, Oj.

Sinapis contusæ, ʒj.

Simmer until the caseous part separates, then strain, and add—
Vini Albi, f ʒvj.

Signa.—A draught of this whey may be taken every hour or two in low fever and in debilitated stomach.

Wine Whey.

R. Lactis Vaccinæ, Oss.

Vini Albi, f ʒij. vel f ʒiij.

Boil the milk, then add the wine, and strain. Grate a little nutmeg, and sweeten the clear whey to the taste.

There are few mild stimulants more employed, or more useful than wine whey. Dr. Chapman speaks of it as possessing a kindred action with carbonate of ammonia. The dose to be regulated by the circumstances of the case; from four ounces to ten or more during the day.

Mixture with Wine, &c.

R. Vitellum Unius Ovi,

Olei Cinnamomi, gtt. x.

Misce, et adde—

Vini Albi, }
Aque Cinnamomi, } āā f ʒjss.

Sacchari, ʒj.

Aque destillatæ, f ʒj.

Fiat mistura. *Signa.*—Give two or three tablespoonfuls, at short intervals, in convalescence from low fevers.

Mixture with Brandy.

R. Spiritûs Vini Gallici, f ʒij.

Lactis Vaccinæ, f ʒiv.

Sacchari, ʒss.

Myristicæ, ad gustandum.

Signa.—Take a tablespoonful every hour or two in low fevers.

Essences of Ginger.

There are several popular and useful stomachics and stimulants now sold under this title. They are easily prepared by making a saturated tincture of the root, using strong or official alcohol, and then carefully concentrating the tincture by means of a water-bath.

The Infusion of Ginger

Is also an excellent stomachic, and may be employed with great benefit in debility of the stomach and bowels.

CLASS X.

NARCOTICS.

NARCOTICS have been denominated *sedatives*, from their power of diminishing action; *anodynes*, from their capability of alleviating pain; and *hypnotics*, or *soporifics*, from their power of inducing sleep. A few words only will be necessary as general rules for their administration, and any peculiarities that individuals of the class may possess will be noticed under the appropriate prescriptions. It is well known they possess the power of fulfilling two indications; namely, of exciting and depressing the nervous energy.

To obtain the first of these results, it is necessary to begin with small doses, and repeat them frequently; while in order to arrive at the second, a full dose must be given, and not resorted to again for a considerable length of time. We should also bear in mind that the system very soon becomes habituated to their action, and the doses must be continually enlarged when it is necessary to maintain their impression. The difference between the susceptibility to their action when the body is in a morbid or healthy condition, should not be overlooked.

PILLS.

Pills of Opium.

R. Pulveris Opii, gr. xij.

Saponis, q. s.

Misce, et divide in pilulas xij.—Of which one is the ordinary dose to produce sleep.

In equal weights of powdered and soft opium, the former is much the more powerful, the latter containing a considerable proportion of water. This should be recollected in directing the medicine in the form of pills.

An opium pill that has been made long enough to become perfectly hard and dry, will be retained by an irritable stomach when a fresh one would be rejected.

Pills of Lupulin.

R. Lupulinæ, ʒss.

Contunde in mortario, et divide in pilulas x.

Signa.—Two pills may be given for a dose, and the quantity doubled if necessary.

This substance, we are told by Dr. Ives, contains all the active properties of the hop. It is slightly narcotic, and is serviceable in irritable stomach, and as a preventive of nocturnal erections in different forms of acute venereal disease.

The lupulin is sometimes administered in powder, with two parts of white sugar.

Pills of Sulphate of Morphia.

R. Morphiæ Sulphatis, gr. ij.

Confectionis Rosæ, q. s. ut fiant pilulæ xij.

Signa.—One may be given for a dose, *pro re nata*.

One grain of either of the salts of morphia is usually estimated as equal to six of opium; so that a pill of the above prescription will be equivalent to a grain of opium.

The acetate and muriate of morphia are prepared in the same way, and given in the same dose.

Pills of Lactucarium.

R. Lactucarii, gr. xij.

Confectionis Rosæ, q. s.

Divide in pilulas vj.

Signa.—One for a dose, to be repeated in two hours, if sleep be not procured.

The inspissated juice of the garden lettuce is said to act as a direct sedative, lessening the frequency of the pulse, diminishing the natural heat, and procuring sleep, without inducing the subsequent distressing effects which follow the use of opium.

The *thridace* of the French appears to be prepared by a different process from that employed in the preparation of the British lactucarium. The former is not so highly esteemed in this country as the latter.

Pills of Henbane and Ipecacuanha.

R. Extracti Hyoseyami, gr. x.

Pulveris Ipecacuanhæ, gr. v.

Misce, et divide in pilulas x.—One every hour or two until relief is procured, watching the effect.

In flatulence and irritability of the bowels.

Pills of Opium, Henbane, &c.

R. Pulveris Opii, gr. iv.
 Extracti Hyoscyami,
 ——— Conii, āā gr. xv.

Fiat massa, in pilulas x. dividenda.—One every night when an anodyne is required.

Pills of Extract of Henbane.

R. Extracti Hyoscyami, ℥j.

Divide in pilulas x.—Of which one may be given every four hours, increasing the dose if necessary.

This medicine, which is employed as a substitute for opium, does not constipate the bowels.

Pills of Extract of Stramonium.

R. Extracti Stramonii Foliorum, ℥ss.

Divide in pilulas xv. *Signa.*—One three times a day.

The powdered leaves are also prescribed in similar doses. The extract of the seeds is given in the dose of a quarter to half a grain.

Opiate Confection.

R. Confectionis Opii, ℥ss.

To be given for a dose.—One grain of opium is contained in thirty-six grains of the officinal confection.

Pills of Extract of Hemlock.

R. Extracti Conii, ℥j.

Pulveris Conii, q. s. ; ut fiat massa in pilulas xx. dividenda. One of these pills night and morning, may be taken at the commencement, gradually increasing them to two, three, four, or more, as the patient can bear them.

The efficacy of narcotic plants is influenced by soil, climate, exposure, and cultivation ; and the extracts by the season when the plants are gathered, and the mode of preparing them. Physicians would do well therefore to make themselves acquainted with the quality of the article they prescribe, inasmuch as there may be a fatal difference between two parcels of different ages, modes of preparation, &c.

Pills of Camphor.

R. Camphoræ, ℥ss.

Alcoholis,

Acaciæ, āā q. s.

Ut fiant pilule xv.—Dose, a pill every two or three hours.

Pills of Opium, Digitalis, &c.

R. Pulveris Opii,

——— Digitalis, āā gr. vj.

Confectionis Rosæ, q. s. ut fiant pilulæ xij.

Signa.—One to be taken every four hours. In asthma, &c.*Pills of Extract of Conium and Calomel.*

R. Extracti Conii, ℥ij.

Hydrargyri Chloridi Mitis, gr. xv.

Fiant pilulæ xv. *Signa.*—One three times a day. In pseudo-syphilitic affections.*Pills of Powdered Nux Vomica.*

R. Pulveris Nucis Vomicae, ℥j.

Confectionis Rosæ, q. s. ut fiant pilulæ x.

Signa.—One to be given two or three times a day, closely watching the effects. In paralysis.*Pills of the Extract of Nux Vomica.*

R. Extracti Nucis Vomicae, gr. x.

Divide in pilulas xv.—One pill twice a day, gradually augmenting the dose, and watching its effect.

Pills of Strychnia.

R. Strychniæ, gr. ij.

Confectionis Rosæ, q. s. ut fiant pilulæ xxiv.

Of which one pill may be given once or twice a day, according to the idiosyncrasy of the patient.

Strychnia forms with acids a variety of salts, which are generally very soluble, and for this reason much more active than the alkali uncombined. The commencing dose of the salt should not be greater than that of the alkaloid. It should be borne in mind that the activity of this substance is increased by all acid drinks.—See MAGENDIE'S *Formulary*, edit. 5, p. 12.

Pills of Camphor, Assafoetida, &c.

R. Camphoræ in pulverem, ℥ij.

Pulveris Assafoetidæ, ℥ij.

Extracti Belladonnæ, ℥ij.

——— Opii, gr. x.

Fiat massa, in pilulas lx. dividenda. Commence with two pills a day, and gradually increase the quantity to six in the 24 hours. Used in hysteria.

DEBREYNE.

Pills with Aconite and Dover's Powder.

R. Extracti Aconiti, gr. vj.

Pulveris Ipecacuanhæ et Opii, gr. xij.

Syrupi, q. s.

Ut fiat massa, in pilulas vj. dividenda.

Signa.—Take one pill three times a day in chronic rheumatism.

Pills of Digitalin.

R. Digitalinæ, gr. j.

Sacchari Lactis, gr. lx.

Syrupi, q. s.

Ut fiat massa, in pilulas xl. dividenda.

Signa.—Take one pill twice a day, gradually and carefully increasing the dose, until the peculiar effects of digitalis are developed.

The greatest care should be used in making these pills, to have the digitalin uniformly diffused through the sugar of milk. A safer way of administering it is in the form of solution. (See page 70.)

MIXTURES, TINCTURES, &c.

Opiate Mixture.

R. Extracti Opii, gr. v.

Sacchari, ʒj.

Aquæ Cinnamomi, fʒvj.

Fiat mistura. *Signa.*—A tablespoonful may be taken every two hours, so that the whole be consumed in twenty-four. In tetanus and colica pictonum.

Anodyne Draught.

R. Tincturæ Opii, gtt. xv. vel xxv.

Syrupi Papaveris, Lond. fʒij.

Spiritûs Cinnamomi, Lond. fʒj.

Aquæ destillatæ, fʒiiss.

Misce. *Signa.*—Anodyne draught. Laudanum is very disagreeable to some persons; combined as above, the taste is disguised.

Laudanum.

R. Tincturæ Opii, fʒss.

Signa.—Dose, twenty-five drops.

As the regulation of doses for children is, to the young practitioner, one of the most difficult parts of prescribing, and as laudanum is an article so universally employed in their diseases, we subjoin the following graduated table from Dr. Dorsey's *Syllabus*, page 36, 1817.

For a child at birth, or within a month after	$\frac{1}{2}$ to 1 drop.
Under a year old	$\frac{1}{2}$ to 2 or 3 drops.
From one to two years of age	1 to 5 "
From two to five "	2 to 8 "
From five to ten "	5 to 15 "
From ten to fifteen "	10 to 20 "
At fifteen years	15 to 20 "
For an adult	25 to 30 "

It is important in the employment of laudanum that it should be a *saturated* tincture and perfectly transparent. For these reasons it should be obtained from the apothecary in small quantities, as it is liable to deposit part of the opium; hence the liquid near the surface is weaker. Infants have been destroyed by the exhibition of *thick laudanum*. Physicians, therefore, should always examine the phial before trusting the nurse to give a dose to a child.

Black Drop.

R. Aceti Opii, f ʒss.

Signa.—Give ten drops as a dose.

This preparation is much stronger than laudanum, and has obtained celebrity from the fact that it equally promotes the ends of laudanum without affecting the head and stomach. When it cannot be had, Dr. Chapman states that, by giving the alcoholic tincture of opium in vinegar, we obtain nearly the same results.

Paregoric Elixir.

R. Tincturæ Opii Camphoratæ, f ʒj.

A teaspoonful of this in a little water is a dose for an adult. It is principally employed as an expectorant.

Sydenham's Laudanum, or Wine of Opium.

R. Vini Opii, f ʒj.

Signa.—Dose, twenty drops.

Our officinal wine of opium, now so much used as an external application in chronic inflammation of the eyes, differs only from Sydenham's laudanum, in not having an ounce of saffron added to the pint. The other ingredients are the same.

Solution of Sulphate of Morphia.

℞. Morphiae Sulphatis, gr. j.

Aquæ destillatæ, f ʒij.

Fiat solutio.—Dose, one or two teaspoonfuls.

When morphia and its salts were first introduced, the formulæ given by Magendie were generally adopted, and are still retained in New York, and some other places. These contain sixteen grains of the sulphate or acetate of morphia in a fluid-ounce of water, whereas the officinal solution has but one grain to the ounce. Practitioners settling in that vicinity should avoid confounding the two solutions. EDITOR.

The sulphate of morphia has one superiority over the acetate, namely, that the latter when dissolved in water is decomposed, two salts resulting, the one soluble, the other not.¹ From this objection the sulphate is exempt.

The muriate and nitrate of morphia possess no advantages over the sulphate and acetate.

Codéine.

The French chemists have also discovered in opium a principle allied in its effects to morphia, and called *Codéine*. In dose of a single grain it proves highly anodyne, but distresses the stomach in larger doses. Its salts are given in somewhat smaller quantities. MAGENDIE, 9^{me} ed. p. 80.

Tincture of Opium and Oil of Turpentine.

℞. Tincturæ Opii, f ʒij.

Olei Terebinthinæ, f ʒss.

Misce. *Signa*.—A teaspoonful diluted every hour or two may be given in delirium tremens, but even here its effect must be closely watched.

The turpentine may be added or omitted at the discretion of the physician, and the quantity of laudanum in some cases may be doubled. We have successfully given two teaspoonfuls every hour until ʒij. of laudanum were taken. We do not believe in the doctrine, however, that opium in this disease may be administered almost *ad libitum* without danger. On the contrary, we feel assured that death has been caused by its excessive use in delirium tremens. S. G. MORTON.

¹ This difficulty, however, is obviated by the addition of acetic acid to convert the insoluble morphia present into an acetate. EDITOR.

Tincture of Lupulin.

R. Lupulinæ puræ, ʒj.

Alcoholis, f ʒij.

Digest for six days in a close vessel. Strain, filter, and add a sufficient quantity of alcohol to make ʒiij. of the tincture. Of this, from ʒss. to ʒij. may be taken for a dose, commencing with the smaller quantity.

Tincture of Hops.

R. Tincturæ Humuli, f ʒj.

Signa.—Of this, a teaspoonful may be taken every two hours, as a narcotic in cases where opium is inadmissible.

Hops are also given in the form of infusion: for which see Tonics. Inclosed in a bag, they have acquired some celebrity as a pillow.

Camphor and Hoffman's Anodyne.

R. Aquæ Camphoræ, f ʒiv.

Spiritûs Ætheris Compositi, f ʒij.

Misce.—Dose, a dessertspoonful every hour or two hours diluted.

In nervous affections, and sleeplessness of fever.

Tincture of Digitalis.

R. Tincturæ Digitalis, f ʒij.

Of which ten drops may be given at first twice or three times a day, increasing the quantity gradually as the system becomes accustomed to it. The peculiar character of this remedy should induce caution in giving it.

Hemlock Mixture.

R. Extracti Conii, ʒss.

Syrupi Papaveris, Lond. f ʒj.

Aquæ, f ʒvij.

Misce.—A tablespoonful three or four times a day for a dose.

Tincture of Extract of Nux Vomica.

R. Extracti Nucis Vomicae, gr. iij.

Alcoholis, f ʒj.

Fiat tinctura.—This tincture to be given in doses of fifteen or twenty drops, in any kind of drink.

This preparation, and the pills made of the resin of nux vomica, have been successfully used in general and local paralysis.

Tincture of Strychnia.

R. Alcoholis, f ʒj.

Strychniæ, gr. iij.

Fiat tinctura.—This tincture to be given by drops, from six to twenty in draughts of mucilage.

Used for the same purposes as the above.

Strychnia Mixture.

R. Strychniæ, gr. j.

Acidi. Acetici, gtt. ij.

Sacchari, ʒij.

Aquæ destillatæ, f ʒij.

Fiat solutio. *Signa.*—A teaspoonful morning and evening.

Solution of Extract of Belladonna.

R. Extracti Belladonnæ, gr. iij.

Aquæ Cinnamomi, f ʒj.

Fiat solutio. *Signa.*—Two or three drops twice or thrice a day, to a child under a year old, and an additional drop for every additional year.

Belladonna has been recently used in Germany as a preventive of scarlatina. The extract, if given for several days, produces a scarlet eruption and affection of the fauces, analogous to those of scarlatina; and it is pretended that this factitious disease is sufficient to prevent the access of the real one.

Mixture with Magnesia, Assafetida, &c.

R. Magnesiæ Carbonatis, ʒss.

Tincturæ Assafoetidæ, gtt. lx.

———— Opii, gtt. xx.

Sacchari, ʒj.

Aquæ destillatæ, f ʒj.

Misce et fiat mistura.—Twenty-five drops may be given to an infant of two or four weeks old; in flatulent colic, diarrhoea, &c.

This preparation was strongly recommended and much employed by the late Dr. Dewees of this city.

If the *calcined* magnesia be used, it unites with the resin of the assafetida, forms an almost solid mass at the bottom of the vessel, and defeats the object of the prescription.

Tincture of Hemp.

R. Extracti Cannabis, gr. xxiv.

Alcoholis, f ʒj.

Fiat solutio. *Signa.*—Take ten drops every two hours.

The above is the formula of Dr. O'Shaughnessy, of Calcutta,

who has had much experience with the gunjah, or Indian hemp. In cholera, he gave ten drops every half hour to check the vomiting; in tetanus, a dram of the tincture, with the same interval, until the paroxysms ceased, or catalepsy was induced; and in acute rheumatism, twenty drops until symptoms of intoxication were manifested.

Dr. B. H. Coates has tried it in a case of tedious labor, in which it seemed to have the effect of restoring the uterine contractions by acting "as musk, castor, or assafetida frequently do in other hollow viscera, by restraining excessive and irregular contractions, and enabling the contracting organ to combine its efforts more in conformity to a healthy order, and a useful tendency and result."

The great uncertainty in the strength of the extract heretofore imported, has interfered with its introduction into practice; but there is now (1853) to be had in Philadelphia, an extract which is decidedly narcotic in the dose of a quarter to half a grain.

EDITOR.

Mixture with Oil of Bitter Almonds.

R. Olei Amygdalæ Amaræ, gtt. xx.

Alcoholis, fʒiij.

Fiat solutio. *Signa*.—Take ten drops four times daily.

Used in neuralgia, especially tic douloureux.

HORN.

CLYSTERS AND SUPPOSITORIES.

Opium Enema.

R. Pulveris Opii, gr. ij.

Mucilaginis Acaciæ, fʒss.

Lactis tepifacti, fʒij.

Misce pro enema.

Opium Enema.

R. Pulveris Opii, gr. iij. vel iv.

Adipis, ʒj.

Melt them together with a gentle heat, and when of a proper warmth, inject the mixture with a small syringe.

Laudanum Enema.

R. Tincturæ Opii, gtt. l.

Infusi Lini, fʒij. vel fʒiv.

Fiat enema.

It is generally supposed that three times the quantity of a medicine may be thrown into the rectum that may be taken into the stomach. As regards opium, however, we believe it to be too much, particularly in children. Indeed, it is an unsettled point in therapeutics, whether remedies act most powerfully upon the rectum or the stomach.

Suppository of Opium.

R. Pulveris Opii, gr. ij.

Saponis, gr. iv.

Misce bene.—Introduce into the rectum when an enema cannot be administered.

Suppository of Opium and Rhatany.

R. Olei Cacao, ℥j.

Extracti Krameriae, ℥ij.

Pulveris Opii, gr. v.

Misce secundem artem, et fiant suppositoria x. PANCOAST.

Suppository of Hemlock.

R. Ceræ Albæ, gr. xv.

Olei Cacao, ℥j.

Extracti Conii, gr. x.

Misce, s. a. The extract should be softened and added to the other ingredients as they begin to cool.

EXTERNAL USE OF NARCOTICS.

Solution of the Extract of Belladonna.

R. Extracti Belladonnæ, ℥j.

Aquæ destillatæ, f ℥iij.

Fiat solutio. *Signa*.—To be applied to the eye in cataract, for the purpose of dilating the pupil and exposing the lens.

Solution of Atropia.

R. Atropiæ, gr. ij.

Alcoholis, f ℥j.

Aquæ destillatæ, f ℥vij.

Fiat solutio. *Signa*.—Let a drop fall into the eye to dilate the pupil.

This is equally efficacious in its action, and much more elegant, than the ordinary preparations of belladonna.—DUNGLISON, *New Remedies*, 6th ed. p. 114. W. W. COOPER.

For internal use, ten drops of this solution represent one-thirtieth of a grain of the alkaloid.

Solution of the Extract of Stramonium Seeds.

This may be made of the same strength, and be used for dilating the pupil in the same way, as the solution of the extract of belladonna.

Solution of Opium in Lime-water, &c.

R. Extracti Opii, gr. j.

Liquoris Calcis,

Olei Amygdalæ, āā f ʒiij.

Fiat mistura.—Recommended by Dr. Sibergundi of Dorsten, in the treatment of sore nipples. It is to be applied to the parts on dossils of lint.

Infusion of Hemlock.

R. Conii, ʒss.

Aquæ bullientis, Oj.

Fiat infusum et cola.—To be applied as a wash to cancerous and scirrhus ulcers.

A cataplasm is also employed for the same purpose, made by mixing the powdered leaves with bread and water poultice.

Infusion of Henbane.

R. Foliorum Hyoscyami, ʒss.

Aquæ bullientis, Oj.

Fiat infusum.—Employed as an external application to cancerous, scrofulous, and other indolent ulcers.

Decoction of Stramonium in Milk.

R. Foliorum Stramonii, ʒij.

Lactis recentis, Ojss.

Boil to a pint and strain. For gouty, rheumatic, and other painful swellings.

Ointment of Belladonna.

R. Extracti Belladonnæ, ʒij.

Aquæ destillatæ, f ʒij.

Adipis, ʒij.

Fiat unguentum.—A portion to be applied to the neck of the uterus in rigidity of that part occasioned by disease, or previous to delivery. It is said to act on the same principle as it does in dilating the iris. The application to be made for some time by means of a syringe with a wide mouth.

M. CHAUSSIER.

Plaster of Opium, &c.

- R. Opii,
 Camphoræ, āā ʒss.
 Emplastri Plumbi, ʒij.
 Melt the plaster, and having previously powdered the other articles, mix them gradually. For local pains.

Plaster of Opium, &c.

- R. Pulveris Opii,
 ——— Camphoræ,
 ——— Saponis, āā ʒj.
 Tincturæ Opii, q. s. ut fiat emplastrum.

This plaster is employed in the same cases as the preceding one. Another, and the most common plan, is to spread theriaca andromachi¹ on leather of the size required, and cover it with powdered opium, sprinkling a little laudanum to promote its adhesion.

Plaster of Opium and Camphor.

- R. Pulveris Opii, ʒij.
 Camphoræ, ʒss.
 Picis Burgundicæ,
 Emplastri Plumbi, āā q. s. ut fiat emplastrum.

This plaster is highly recommended in neuralgia.

GRAVES.

Liniment of Belladonna, Cherry Laurel, &c.

- R. Extracti Belladonnæ, ʒij.
 Solve in—
 Aquæ Lauro Cerasi, ℞℥ f ʒij.
 Dein adde—
 Ætheris, f ʒj.

Misce. *Signa.*—Rub the liniment on the part, and apply flannels moistened with it.

Useful in neuralgia, in gouty and rheumatic pains, and for gathered breasts.

RANQUE.

¹ The original Theriaca Andromachi is composed of seventy-two ingredients, pulverized and rubbed up with honey into an electuary. A preparation bearing the same name, and used for the same purposes, is now made in a much less elaborate way. The word *Theriaca* is, at the present day, applied to any coarse sirup, as treacle or molasses.

Camphor and Chloroform Liniment.

R. Camphoræ, ʒjss. .
Chloroformi, fʒij.
Olei Olivæ, fʒij.
Fiat linimentum.

Useful in rheumatic and neuralgic affections.

W. B. PRICE.

CLASS XI.

ANTISPASMODICS.

ANTISPASMODICS are those medicines which allay spasm, and compose the irregular actions of the nervous system. Some of the most powerful of them are derived from the class of Narcotics.

Pills of Assafoetida and Soap.

R. Assafoetidæ, ʒj.

Saponis, gr. x.

Aquæ, q. s. ut fiant pilulæ xx.

Two or four of these pills may be taken for a dose, and repeated *pro re nata*.

Pilulæ Aloës et Assafoetidæ,

—— Galbani Compositæ,

May be given in the same doses as those made with soap only.

Opium Pills.

As under Narcotics. (Page 133.)

Pills with Castor and Salt of Amber.

R. Castorei, ʒj.

Acidi Succinici, ʒss.

Extracti Gentianæ, q. s. ut fiant pilulæ xxiv.

Dose, three pills night and morning. In hysteria.

Pills of Musk, Camphor, &c.

R. Moschi, ʒss.

Camphoræ, ʒj.

Ammoniaci, ʒij.

Opii, gr. iv.

Misce, et fiant pilulæ singulæ gr. iv.

Signa.—The dose may be four or five in the twenty-four hours.

In nervous diseases.

M. A. RICHARD.

Pills of Valerianate of Zinc.

R. Zinci Valerianatis, gr. xij.

Confectionis Rosæ, q. s.

Ut fiat massa in pilulas xij. dividenda.

Signa.—Give one pill every three hours.

Used in neuralgia, and anomalous nervous affections.

Bolus with Valerian and Rust of Iron.

R. Pulveris Valerianæ, ʒj.

Ferri Subcarbonatis, ʒss.

Mucilaginis Acaciæ, q. s. ut fiat bolus.

One, three times a day.

Bolus with Musk and Camphor.

R. Camphoræ, gr. v.

Moschi, gr. v. vel x.

Syrupi, q. s. ut fiat bolus.

Signa.—To be taken at once.

This is a full dose, which, however, can be readily modified by the practitioner.

To confine the aroma of musk, whether in the form of pills or otherwise, it should always be kept in phials.

Bolus with Musk and Carbonate of Ammonia.

R. Moschi,

Ammonia Carbonatis, āā ʒss.

Confectionis Rosæ, q. s. ut fiat bolus.

To be taken every third hour.

In mortification accompanied with spasmodic action.

Musk Mixture.

R. Moschi optimi, ʒij.

Sacchari, ʒj.

Pulveris Acaciæ, ʒj.

Aquæ destillatæ, fʒvj.

Fiat mistura. *Signa.*—A tablespoonful to be given every hour or two hours.

Mixture with Musk, &c.

R. Misturæ Moschatæ, fʒvj. (See preceding formula.)

Tincturæ Opii Camphoratæ, fʒss.

——— Valerianæ Ammoniatæ, fʒj.

Misce. *Signa.*—A teaspoonful three or four times a day. For children in pertussis.

Tincture of Assafetida, Castor, &c.

R. Tincturæ Assafœtidæ,
 ———— Castorei, āā f̄j.

Spiritûs Ammoniae Aromatici, f̄j.

Misce.—A teaspoonful may be taken in a glass of sweetened water for a dose.

Hoffman's Anodyne and Laudanum.

R. Spiritûs Ætheris Compositi, f̄jij.

Tincturæ Opii, gtt. lxxx.

Aquæ Cinnamomi, f̄vj.

Fiat mistura. *Signa*.—A tablespoonful every one or two hours.
 In hysteria, &c.

Mixture with Assafetida, &c.

R. Assafœtidæ, j.

Aquæ Menthæ Piperitæ, f̄jij.

Fiat solutio, et adde—

Tincturæ Valerianæ Ammoniata, f̄jij.

——— Castorei, f̄jij.

Ætheris, f̄j.

Signa.—A tablespoonful for a dose, largely diluted, every second hour. In hysteria.

Oil of Valerian with Spirits of Hartshorn.

R. Olei Valerianæ, gtt. viij.

Spiritûs Ammoniae Aromatici, f̄j.

Aquæ, f̄jiv.

Sacchari, j.

Misce. *Signa*.—To take a tablespoonful every two or three hours shaking the phial.

Infusion of Valerian.

R. Valerianæ, j.

Aquæ bullientis, Oj.

Put them into a covered vessel and let them stand until cold.—

Dose, a wineglassful every two or three hours.

Tincture of Valerian and Hoffman's Anodyne.

R. Spiritûs Ætheris Compositi,

Tincturæ Valerianæ, āā f̄j.

Misce. *Signa*.—A teaspoonful for a dose, with water, *pro re nata*, shaking the phial each time.

Mixture with Ether and Laudanum.

- R. Ætheris, fʒij.
 Sacchari,
 Acaciæ, āā ʒiss.
 Tincturæ Opii, gtt. lx.
 Aquæ Cinnamomi, fʒij.

Misce. *Signa*.—A teaspoonful every hour. In cramp of the stomach, &c.

Ethereal Solution of Chloride of Zinc.

- R. Zinci Chloridi, ʒj.
 Alcoholis, fʒij.
 Ætheris, fʒss.

Solve, post aliquot dies decanta.

Signa.—Take from four to eight drops, twice daily. Used in chorea, epilepsy, and anomalous nervous affections.

HUFELAND.

Embrocation of Oils of Cloves, Amber, &c.

- R. Olei Olivæ, fʒj.
 — Caryophylli,
 — Succini rectificati, āā fʒss.

Misce. Used as an embrocation in whooping-cough. ROCHE.

Assafetida Enema.

- R. Tincturæ Assafœtidæ, fʒij.
 Decocti Hordei, fʒx. vel fʒxij.

Misce pro enemate.—Tinct. opii may be added *pro re nata*. In hysteria, spasmodic colic, &c.

Musk Enema.

- R. Moschi, gr. xij.
 Sacchari, ʒij.
 Spiritûs Ammoniaë, gtt. xxx.
 Infusi Lini, fʒiv.

Fiat enema.—For children with convulsions.

CLASS XII.

SIALAGOGUES.

THOSE medicines which increase the salivary secretion are denominated Sialagogues. In forming these into a *class* in this place, we trespass on the arrangement of Dr. Chapman, but merely from a motive of convenience; for we agree with him that such a class would be better expunged from the *Materia Medica*, at the same time that it appears necessary to introduce it here, in order to place the mercurial preparations in stronger relief than could otherwise be done.

Pills of Calomel.

R. Hydrargyri Chloridi Mitis, gr. xij.

Confectionis Rosæ, q. s. ut fiant pilulæ xij.

Signa.—One, morning and evening, and, if necessary, until salivation is excited.

Pills of Corrosive Sublimate.

R. Hydrargyri Chloridi Corrosivi, gr. v.

Aquæ destillatæ, gtt. xxx. vel xl.

Confectionis Rosæ, ʒj.

Pulveris Glycyrrhizæ, q. s. ut fiant pilulæ xl.

Of which, give one pill three or four times a day. In lues venerea. As salivation rapidly follows the use of corrosive sublimate, its effect must be watched.

Dupuytren's Antisyphilitic Pills.

R. Hydrargyri Chloridi Corrosivi, gr. ss.

Extracti Cinchonæ, gr. x.

————— Opii, gr. ss.

Pulveris Cinchonæ, q. s.

Ut fiant pilulæ ij.—The two pills may be taken, one in the morning, and the other at night; but as each contains a

quarter of a grain of corrosive sublimate, their action must be very closely watched. M. Dupuytren was partial to these pills in the treatment of syphilis. He thought the opium and cinchona promoted and beneficially modified the action of the sublimate.

Pills with Corrosive Sublimate and Hemlock.

R. Hydrargyri Chloridi Corrosivi, gr. vj.
Solve in Aquæ destillatæ, q. s. et adde—
Extracti Conii, ʒj.
Pulveris Conii, q. s. ut fiat massa, in pilulas xlvij. dividenda.
Signa.—Give one pill three times a day.

Used in lues venerea.

The effect of the mercurial preparations upon children is sometimes very serious, and even fatal. So powerful upon them is the action of this medicine, that a profuse or even gentle salivation will sometimes produce mortification and destruction to the jaw, cheek, and lip. This, however, does not often happen; but to avoid it the gums should be frequently and carefully examined. It is, nevertheless, a fact, that a disease precisely similar in all its features occasionally attacks children who are badly clothed and fed, though no mercury has been given to them.

Pills of Iodide of Mercury.

R. Hydrargyri Iodidi, gr. v.
Confectionis Rosæ, q. s.
Ut fiant pilulæ xxx. Each pill will contain the sixth of a grain, which may be given three times a day. Double the quantity (two pills) is sometimes administered at first, but care should be taken not to push it too far.

Compound Pills of Iodide of Mercury.

R. Hydrargyri Iodidi, gr. ij. vel gr. iij.
Extracti Opii, gr. ij.
Lactucarii, gr. xij.
Guaiaci Resinæ, gr. xxiv.
Misce, et fiant pilulæ xxiv.—Given by M. Ricord in the syphilis of children. He directs one pill for a child six months old, and two, three, or four pills for older children.

DUNGLISON'S *New. Rem.* p. 280.

Pills of Iodide of Mercury.

- R. Hydrargyri Iodidi, gr. j.
 Extracti Juniperi vel Glycyrrhizæ, gr. xij.
 Pulveris Glycyrrhizæ, q. s.

Ut fiant pilulæ viij.—In adult patients, Magendie directs two to be given morning and evening, to be increased to three or four.
 DUNGLISON, *ut supra*.

Pills with Calomel, Camphor, &c.

- R. Hydrargyri Chloridi Mitis,
 Camphoræ, āā ʒj.
 Pulveris Opii, gr. xij.
 Syrupi, q. s. ut fiant pilulæ xx.

Signa.—One or two to be taken morning, noon, and night, when calomel by itself causes purging.

Blue Mercurial Pills.

- R. Pilulæ Hydrargyri, ʒss.
 Divide in pilulas x.

Signa.—One may be taken three or four times a day, according to circumstances.

If they should occasion purging, half a grain of opium may be added to each pill. The mass directed in the *American Pharmacopœia* contains one grain of mercury in every three grains.

Pills of the Red Oxide of Mercury.

- R. Hydrargyri Oxidi Rubri, gr. j.
 Pulveris Opii, gr. j.
 Olei Caryophylli, gtt. ij.

Fiant pilulæ iij. *Signa*.—One pill every night for a week. In venereal cases.

This preparation, in a stronger formula, was used by the celebrated John Hunter; but from its harshness it is now generally superseded by milder medicines.

Solution of Cyanuret of Mercury.

- R. Hydrargyri Cyanureti, gr. viij.
 Aquæ, ʒj.

Fiat solutio.—This is the *Liqueur anti-syphilitique* of Chaussier, who prefers it to corrosive sublimate on account of its greater solubility and more speedy action. By giving a teaspoonful of the solution, we give one-sixteenth of a grain of the cyanuret, which may be repeated morning, noon, and night, watching its effects.

Powder of Mercury and Chalk with Ipecacuanha.

R. Hydrargyri cum Cretâ, ʒj.

Pulveris Ipecacuanhæ, ʒss.

Fiant chartulæ vj. *Signa.*—One night and morning in sirup.

Commended by Dr. Paris, in dyspepsia, in which the biliary secretion is impaired. He prefers it under these circumstances to every other form of mercury.

Masticatory with Pellitory, &c.

R. Pulveris Pyrethri,

Mastiches, āā ʒj.

Let these be dissolved together by a moderate application of heat; then divide the mass into two equal parts. One of these lumps may be chewed at pleasure. In toothache, and paralytic affections of the tongue.

Mixture of Pellitory, Opium, &c.

R. Radicis Pyrethri contusæ, ʒss.

Aceti destillati, f ʒvj.

Opii, gr. iij.

Digest in a sand-bath for an hour. *Signa.*—A spoonful to be held in the mouth and frequently renewed. Used in the same cases as the former.

There are several other substances, such as polygala senega, sal ammoniac, nitric acid, &c., which occasionally excite salivation; but they are seldom employed for that purpose.

CLASS XIII.

TONICS.

TONICS are those medicines which impart strength to the system, without subjecting it to any preternatural excitement during their operation. They stimulate, however, in a secondary manner, by gradually increasing the force of the circulation, and by restoring the digestive and secretory functions to a more healthy state. Considerable perseverance is often necessary in giving any one of this class a fair trial; and when one fails, we may often resort with success to another. Much also may sometimes be gained by alternating them.

[It is a well-known fact that some medicines possessed of tonic properties, in small or ordinary doses, have the singular power, when administered more freely, of producing such changes in the economy as to render them applicable to the cure of intermittent disease. The numerous remedies of this class, and their varied modes of exhibition, render it advisable, in a work of this kind, to have some mode of distinguishing them from ordinary tonics. For convenience of reference, therefore, two sub-classes have been made, viz: 1. Of Anti-Intermittents, and 2. Of Ordinary Tonics. EDITOR.]

SUB-CLASS I.—ANTI-INTERMITTENTS.

At the head of this class, stand confessedly Peruvian bark and its numerous preparations.

POWDERS AND PILLS.

Powder of Peruvian Bark.

R. Pulveris Cinchonæ, ʒj.

A dram (about a teaspoonful) to be taken every hour in port-wine, water, or milk, during the intermission of fever.

Some practitioners suspend the bark till within a few hours of the expected paroxysm ; and a few others administer it during the hot stage of the disease. It is better to commence immediately after the paroxysm has ended, and give a dose every hour, until the expected period of its return ; and to continue it in smaller quantities for several days.

Powder of Peruvian Bark and Cloves.

R. Pulveris Cinchonæ,
Potassæ Bitartratis, āā ʒj.
Pulveris Caryophylli, ʒj.

Misce.—Dose, a dram and a half every second hour.

This compound will sometimes succeed when the bark alone fails.

Powder of Peruvian Bark and Sulphate of Magnesia.

R. Pulveris Cinchonæ,
Magnesiæ Sulphatis, āā ʒvj.

Fiat pulvis, et divide in partes æquales iv.—One to be taken every two hours, when purging is required.

Powder of Peruvian Bark, Snake-root, &c.

R. Pulveris Cinchonæ, ʒss.
——— Serpentariæ, ʒj.
Sodæ Bicarbonatis, ʒij.

Divide in chartulas iv. *Signa.*—One to be given every two hours. In obstinate intermittents.

If the bark should purge when given alone, a little opium may be added ; when it oppresses the stomach, cloves, as in a former prescription ; if costiveness should follow, rhubarb or sulphate of magnesia. The alkali in the above preparation will correct acidity.

Powder of Prussiate of Iron and Guaiacum.

R. Ferri Ferrocyanureti,
Pulveris Guaiaci, āā ʒj.

Misce, et divide in chartulas xij. *Signa.*—One, three times a day. In obstinate intermittents.

Pills of Sulphate of Quinia.

R. Pilulæ Quiniæ Sulphatis, No. x.

Signa.—Take two pills every hour, in the apyrexia of intermittent fever, until eight or ten are taken.

The officinal pill contains one grain. Sometimes it is desirable to administer a larger quantity. In such cases, Mr. E. Parrish, of this city, recommends the following form, viz:—

R. Quiniæ Sulphatis, gr. xx.

Acidi Sulphurici Aromatici, gtt. xv.

“Drop the acid into the sulphate of quinia on a tile or slab, and triturate it with a spatula until it assumes a pilular consistence; then divide into the required number of pills.” Made in this way a three-grain pill is not inconveniently large.

In prescribing quinia in the apyrexia of intermittents, and especially in that of neuralgia, it is often necessary to increase the dose beyond the officinal formula. For example, three, four, or five grains may be requisite in those cases in which the neuralgic paroxysms are violent and the intermission short; and this quantity must be repeated every two or three hours, or even every hour. The principal inconveniences arising from the free exhibition of quinia are a painful sense of fulness, noise, and dizziness of the head, constriction of the chest, and purging. The latter may be checked by accompanying each pill with a few drops of laudanum, or by a tablespoonful of the simple aqueous infusion of camphor. If the affection of the head or chest becomes severe, the medicine must be suspended or diminished in quantity.

Pills of Sulphate of Quinia and Carbonate of Ammonia.

R. Quiniæ Sulphatis, gr. xij.

Ammoniæ Carbonatis, gr. xxiv.

Fiant pilulæ xij. *Signa.*—One every hour, beginning six hours before the expected paroxysm.

These pills are large, but they will act when quinia alone proves unavailing. Quinia may also be combined advantageously with the oil of black pepper, or other essential oils, in case of irritable stomach.

Pills of Sulphate of Cinchonia.

R. Cinchonæ Sulphatis, ʒss.

Confectionis Rosæ, q. s.

Ut fiat massa, in pilulas xx. dividenda.

Signa.—Take two pills every hour, in the apyrexia of intermittent fever, until eight are taken.

Experiments made in the Pennsylvania Hospital, prove the sulphate of cinchonia to be an efficient substitute for quinia, in a somewhat larger dose.

EDITOR.

Pills of Salicine.

R. Salicin, gr. xxiv.

Mucilaginis Acaciæ, q. s.

Ut fiant pilulæ viij.—A pill to be taken three or four times a day.

Salicine possesses decided anti-intermittent powers. It has to be used, however, in doses double or treble those of the sulphate of quinia.

Pills of Piperine.

R. Piperin, gr. xij.

Extracti Gentianæ, q. s. ut fiant pilulæ xij.

One every hour during the apyrexia of fever.

This is one of the preparations lately introduced into practice. It is a white crystalline substance procured from the black pepper, and was thought to possess considerable febrifuge powers. It is stated by Pelletier, however, that its supposed virtues depend only on the acrid concrete oil with which it is associated, and that when perfectly pure it is quite inert.

Pills of Arsenic, &c.

R. Acidi Arseniosi, gr. j.

Pulveris Opii, gr. iij.

Saponis, gr. viij.

Pulveris Glycyrrhizæ, q. s. ut fiat massa in pilulas xx. dividenda.

Signa.—Give one pill two, three, or four times a day, and increase gradually, according to circumstances, watching the effect. In intermittent fever, and local diseases of an intermittent character.

Pills of Muriate of Ammonia, Arsenic, &c.

R. Ammoniæ Muriatis, ℥ss.

Pulveris Opii, gr. viij.

Acidi Arseniosi, gr. iv.

Syrupi, q. s. ut fiant pilulæ xxxij.

Of which one may be given three times a day. In intermittent fever.

Each of these pills contains one-eighth of a grain of arsenious acid. The utmost care is requisite in compounding this and all other formulæ for solid arsenic, at the same time that its specific action must be closely watched.

Pills with the Sulphate of Copper.

- R. Cupri Sulphatis, gr. iv.
 Extracti Cinchonæ Rubræ, gr. xxxij.
 Syrupi, q. s. ut fiat massa, in pilulas xvj. dividenda.
 One to be taken three times a day. In obstinate intermittents.

Pills of Extract of Bark, &c.

- R. Extracti Cinchonæ Rubræ,
 ——— Gentianæ, āā ʒj.
 Ferri Sulphatis, ʒss.
 Pulveris Myrrhæ, ʒj.
 Olei Cari, gtt. x.
 Syrupi Zingiberis, q. s. ut fiant pilulæ lx.
Signa.—Three to be taken three times a day.

Pills of Sulphate of Copper and Opium.

- R. Cupri Sulphatis, gr. iij.
 Pulveris Opii, gr. iv.
 ——— Acaciæ, gr. x.
 Syrupi, q. s.
 Ut fiat massa in pilulas xij. dividenda.
Signa.—Give one pill every three hours in obstinate intermittents. This combination is sometimes serviceable, when quinia fails.

CHAPMAN.

Pills of Sulphate of Bebeerine.

- R. Bebeerinæ Sulphatis, ʒij.
 Confectionis Rosæ, q. s.
 Fiat massa, et divide in pilulas xl.
Signa.—Give two pills every two hours in intermittent fevers.
 This article has lately attracted attention in Edinburgh and Paris, and the experiments thus far reported warrant a more extended trial of its powers.

EDITOR.

DECOCTIONS AND INFUSIONS.

Decoction of Cinchona.

- R. Cinchonæ Rubræ contusæ, ʒj.
 Aquæ, Oj.
 Boil for ten minutes—and while hot add—
 Serpentariæ Radicis, ʒss.
 Aurantii Corticis, ʒij.
 And suffer them to infuse for half an hour near the fire in a covered vessel.—Dose, a wineglassful every hour.

If the bark is boiled longer than ten minutes, it deposits extractive and resinous matter, becomes paler, and is less efficient. The last ingredients are directed to be infused because their aromatic properties would be dissipated by decoction.

Decoction of Bark, &c.

- R. Decocti Cinchonæ Rubræ, f ʒvj.
 Tincturæ Cinchonæ Compositæ, f ʒj.
 Acidi Sulphurici Aromatici, gtt. x.
 Misce.—A tablespoonful may be taken every hour or two hours
pro re nata.

By this preparation we obtain both the aqueous and alcoholic extracts of the bark.

Infusion of Cinchona with Lime-water.

- R. Pulveris Cinchonæ Rubræ, ʒij.
 Liquoris Calcis, Oij.
 Misce, et fiat infusum.

This infusion was formerly much employed. A wineglassful may be taken three or four times per diem, or once every hour, in intermittent and other forms of fever.

Infusion of Bark.

- R. Pulveris Cinchonæ Rubræ, ʒj.
 Acidi Sulphurici Aromatici, f ʒj.
 Aquæ, Oj.
 Macera per horas decem et cola.
Signa.—Take a wineglassful every two hours.

This infusion represents the virtues of the bark more fully than either of the foregoing.

Infusion of Boneset.

- R. Eupatorii, ʒj.
 Aquæ bullientis, Oj.
 Infuse for two hours, and strain.—Of this a wineglassful may be taken every one or two hours, cold. In intermittent fever.

SOLUTIONS, MIXTURES, &c.

Solution of the Sulphate of Quinia.

- R. Quiniæ Sulphatis, gr. viij.
 Acidi Sulphurici Aromatici, gtt. x.
 Aquæ, f ʒj.

Fiat solutio.

A teaspoonful contains one grain of the salt.

Mixture of Sulphate of Quinia.

- R. Quiniæ Sulphatis, gr. xx.
 Acidi Sulphurici, gtt. j.
 Sacchari, ʒj.
 Aquæ Cinnamomi, f ʒijss.

Misce. *Signa*.—A teaspoonful for a dose, every hour during the apyrexia of intermittents.

By the addition of sulphuric acid, a super-sulphate of quinia is formed, which is soluble in the aqueous menstruum.

Sulphate of Quinia in Sirup.

- R. Quiniæ Sulphatis, gr. xvj.
 Syrupi Zingiberis vel Limonis, f ʒij.

Misce. *Signa*.—A teaspoonful to be given every hour during the apyrexia.

The French physicians employ a wine and tincture of quinia, which, however, are not superior to the compound used in this country. Both in Europe and America, and especially in Italy, the dose is carried to a much greater extent than we have directed. Eight, ten, fifteen, twenty, and even thirty grains have been given by the Italians, without manifest injury and with successful results. Such doses, however, are seldom admissible in this country; and a cure can generally be accomplished by giving sixteen or twenty grains in twenty-four hours, repeated in half doses for three or four successive days.

Mixture with Ferrocyanate of Quinia.

- R. Quiniæ Ferrocyanatis, gr. iv.
 Alcoholis, f ʒj.

Solve et adde—

Aquæ Camphoræ, f ʒvij.

Misce.—This preparation is said to be more active than the sulphate, whence it should be given in less doses; a teaspoonful

of the solution (which will contain half a grain of the salt) may be taken every hour or two hours, according to circumstances. Pills of the ferrocyanate of quinia are prepared with simple mucilage, and administered like the sulphate.

The acetate, citrate, muriate, nitrate, and other salts of quinia, have also been employed in Europe, and occasionally in this country. They possess no advantages over the ordinary preparations, and are given in much the same quantity.

Mixture with Sulphate of Quinia and Tartaric Acid.

R. Quiniæ Sulphatis, gr. vj.

Acidi Tartarici, gr. iij.

Syrupi, f ʒj.

Misce. Dose, a teaspoonful.

The advantages of this preparation, are increased solubility and diminished bitterness of the quinine salt. CASORATI.

Mixture of Sulphate of Quinia and Tannic Acid.

R. Quiniæ Sulphatis, gr. x.

Acidi Tannici, gr. ij.

Aquæ, f ʒvj.

Syrupi Aurantii Corticis, f ʒij.

Misce. *Signa.*—A teaspoonful every two hours.

The object proposed in this formula by Dr. Thomas of Baltimore is to diminish the intense bitterness of the sulphate. This is accomplished by the formation (in part at least) of an insoluble tannate. Whether it is as efficient as a more soluble salt, remains to be proved.

R. H. THOMAS.

Mixture with Tincture of Bark and Citrate of Potash.

R. Succı Limonis, f ʒjss.

Potassæ Carbonatis, ʒj.

Tincturæ Cinchonæ, f ʒj.

Aquæ Cinnamomi, f ʒiij.

Misce. *Signa.*—A tablespoonful every two hours. To promote insensible perspiration while taking the bark.

Sirup of Cinchonia.

R. Cinchoniæ Sulphatis, gr. xxiv.

Syrupi, Oss.

Fiat mistura.—A tablespoonful will contain a grain and a half of the salt, which is a full dose; to be repeated *pro re nata*.

Mixture of Red Bark in Port Wine, &c.

- R. Pulveris Cinchonæ Rubræ, ʒss.
 Theriacæ Andromachi, ʒj.
 Succī Limonis, f ʒij.
 Vini Rubri, f ʒiv.

Misce. *Signa*.—A wineglassful every second hour during the apyrexia of intermittent fever.

We find this prescription strongly recommended in the *N. A. Medical and Surgical Journal*, by Dr. Meigs.

Mixture with Decoction and Tincture of Bark, &c.

- R. Decocti Cinchonæ Rubræ, f ʒiij.
 Tincturæ Cinchonæ Compositæ, f ʒj.
 Pulveris Cinchonæ, ʒij.
 Syrupi, f ʒss.

Fiat mistura.—A tablespoonful to be taken every one or two hours during the absence of fever.

Mixture with Extract of Bark, &c.

- R. Extracti Cinchonæ Rubræ, ʒj.
 Decocti Cinchonæ Rubræ, f ʒvj.
 Tincturæ Cardamomi, f ʒiv.

Misce.—A tablespoonful for a dose, as above directed.

Mixture of Fowler's Solution, &c.

- R. Liquoris Potassæ Arsenitis, gtt. lx.
 Tincturæ Opii, gtt. xxx.
 Spiritūs Lavandulæ Compositi, f ʒj.
 Aquæ Cinnamomi, f ʒiv.

Fiat mistura. *Signa*.—A tablespoonful for an adult, a teaspoonful for a child; to be administered every two or three hours, during the apyrexia of intermittents, watching the effect.

Arsenical Mixture.

- R. Liquoris Potassæ Arsenitis, gtt. x.
 Aquæ destillatæ, f ʒj.
 Tincturæ Opii, gtt. x.
 Spiritūs Lavandulæ Compositi, f ʒss.
- To be taken at a dose.

This mixture is recommended by Dr. Graves, when the simple arsenical solution irritates the stomach.

Enema of Extract of Bark, &c.

R. Extracti Cinchonæ Rubræ, ʒss.

Aquæ tepidæ, f ʒiv.

Solve; dein adde—

Olei Olivæ, f ʒss.

Tincturæ Opii, gtt. x.

Fiat enema.—To be administered every fourth hour. In intermittents.

When preferred, the powdered bark may be used, say ʒij. mixed with some of the decoction instead of common water. Bark is also employed externally in the form of a waistcoat, worn next the skin. A poultice made by mixing the powder and decoction into a paste, may be applied to the region of the stomach; and lastly, the decoction is used as a semicupium.

Enema of Quinia.

Quinia is employed in enemata, and in this form has an immediate and powerful effect on intermittent fevers. From twelve to twenty grains of the sulphate of quinia, dissolved in a tea-cupful of flaxseed or other mucilage, may be administered at a time, and repeated in four or six hours.

SUB-CLASS II.—ORDINARY TONICS.

POWDERS AND PILLS.

Powder of Carbonate of Iron.

R. Ferri Subcarbonatis, ʒj.

Divide in chartulas vj. *Signa*.—One every six hours in molasses. Chorea, neuralgia, &c.

The dose may be carried as far as ʒss. every four or six hours, but must sometimes be preceded by antiphlogistic treatment.

Powder of Colomba, Ginger, &c.

R. Pulveris Colombæ,

———— Ferri Subcarbonatis,

———— Rhei,

———— Zingiberis, āā ʒj.

Misce, et fiant pulveres xij.—One to be taken three times a day, in molasses.

Iron by Hydrogen.

R. Ferri Pulveris, ℥j.

Divide in pulveres xij.

Signa.—Take one powder three times a day in sirup or molasses.

This has been much used in anæmia and chlorosis. It is one of the best preparations of iron.

Powder of Belladonna, Quinia, and Rhubarb.

R. Pulveris Belladonnæ Radicis, gr. jss.

Quiniæ Muriatis, gr. iv.

Pulveris Rhei, gr. xv.

Misce, et divide in pulveres x.

Signa.—Take one powder morning, noon, and night.

This has been employed successfully in engorgements of the liver and spleen, following intermittent fever. SCHWABE.

Powders with Ipecacuanha, Iron, &c.

R. Ferri Subcarbonatis, gr. xlvij.

Pulveris Ipecacuanhæ, gr. vj.

Hydrargyri cum Cretâ, gr. xij.

Misce, et divide in pulveres vj.

Signa.—Take one powder twice a day.

Used with decidedly good effect in chlorosis. ASHWELL.

Powder of Angustura Bark.

R. Pulveris Angusturæ, ℥ss.

Divide in chartulas vj.—Of which one may be taken three or four times a day, in water, or any other vehicle.

This bark has been successfully administered in uterine hemorrhage.

Powder of Colomba and Tartrate of Iron.

R. Ferri et Potassæ Tartratis, ℥ij.

Pulveris Colombæ, ℥ss.

Fiant pulveres iv. *Signa.*—One every three or four hours, in sirup.

Powder with Rust of Iron, &c.

R. Ferri Sulphatis, ℥ij.

— Subcarbonatis, ℥iss.

Divide in pulveres xij. *Signa.*—One to be taken morning, noon, and night, in sirup or molasses.

Pills of Protocarbonate of Iron.

R. Pilulæ Ferri Carbonatis, ʒj.

Ferri Subcarbonatis, q. s.

Ut fiant pilulæ xx. *Signa.*—A pill to be taken morning, noon, and night. This is one of the latest and best formulæ for the chalybeate preparations.

Lactate of Iron.

This salt of iron has lately been introduced into notice, but it has no obvious advantage over the ordinary preparations.—The dose is from gr. v. to ʒj.

Pills of Cinchona and Camphor.

R. Extracti Cinchonæ Rubræ, ʒj.

——— Opii, gr. j.

Camphoræ, gr. xij.

Pulveris Cinchonæ, q. s.

Ut fiant pilulæ xij.—One to be taken three or four times a day.

RATIER.

Pills of Ammoniuuret of Copper.

R. Cupri Ammoniati, ʒj.

Confectionis Rosæ, q. s. ut fiant pilulæ xl.

Signa.—One or two to be taken at bedtime. In epilepsy.

Febrifuge Bolus.

R. Pulveris Cinchonæ Rubræ, ʒj.

Ammoniæ Muriatis,

——— Carbonatis, āā gr. xij.

Antimonii et Potassæ Tartratis, gr. xvij.

Syrupi, q. s. ut fiat massa et divide in partes xlvij.

Signa.—One to be taken four times a day, during the apyrexia of fever.

HÔTEL DIEU.

Pills of Ammoniated Iron, &c.

R. Ferri Ammoniati, ʒj.

Pulveris Rhei, gr. viij.

Confectionis Rosæ, q. s. ut fiat massa in pilulas viij. dividenda.

This formula is recommended in rachitis, and is to be continued with one pill a day for a fortnight, and then intermitted for the same length of time, and again resumed. The object of this direction is to prevent the system from becoming habituated to the medicine.

Pills of Sulphate of Iron and Quinia.

R. Quiniæ Sulphatis, gr. xij.

Ferri Sulphatis, gr. xxiv.

Pulveris Opii, gr. iij.

Syrupi, q. s.

Misce, et fiant pilulæ xij. *Signa.*—Take two pills three times daily. Used in chlorosis and anæmia. KIRKBY.

Pills of Iron and Quinia.

R. Pilulæ Ferri Carbonatis, ℥ss.

Quiniæ Sulphatis, gr. x.

Fiat massa, et divide in pilulas xv. *Signa.*—One pill every four hours. THOMAS.

Pills of Iron, Red Pepper, &c.

R. Pulveris Aloës,

Ferri Sulphatis, āā ʒj.

Mastiches, gr. x.

Pulveris Capsici, ʒj.

Syrupi, q. s.

Fiat massa in pilulas xx. dividenda. *Signa.*—Take one pill every four hours. In chlorosis. COPLAND.

Mitchell's Tonic Pills.

R. Extracti Quassiae, ʒij.

———— Conii, gr. x.

Ferri Subcarbonatis, gr. x.

Liquoris Potassæ Arsenitis, gtt. x.

Fiat massa in pilulas xl. dividenda. *Signa.*—Take one pill three times daily. J. K. MITCHELL.

Pills of Nitrate of Silver.

R. Argenti Nitratis, gr. iv.

Pulveris Glycyrrhizæ, gr. xij.

Syrupi, q. s.

Fiat massa, et divide in pilulas xij.

Signa.—Take one pill every three hours.

Pills of Nitrate of Silver, &c.

R. Argenti Nitratis, gr. v. vel x.

Pulveris Opii, ʒss.

———— Camphoræ,

———— Myristicæ, āā ʒj.

Mucilaginis Acaciæ, q. s. ut fiat massa, et divide in pilulas xx. Dose, a pill morning and evening. In chorea and epilepsy.

Pills of Iodide of Manganese.

R. Potassii Iodidi,
Manganesiæ Sulphatis, exsiccatae, āā ʒj.
Mellis, q. s.

Fiat massa in pilulas xxx. dividenda.

Signa.—To be kept in a well-stopped bottle. Take one pill morning and night.

The salts of manganese have recently been introduced as substitutes for the corresponding preparations of iron.

HANNON.

Pills of Iron and Gentian.

R. Ferri Pulveris, ʒj.
Extracti Gentianæ, ʒij.

Fiat massa, et divide in pilulas xx.

Signa.—Take one pill three times a day.

Pills of the Sulphate of Iron.

R. Ferri Sulphatis, ʒj.

Extracti Gentianæ, q. s. ut fiat massa, et divide in pilulas
xxx.

One to be taken morning, noon, and night. In dyspepsia, &c.

Pills of the Extract of Quassia.

R. Extracti Quassiae, ʒj.

Fiant pilulæ xx.

Of which one may be taken three or four times a day.

This is one of the purest bitters, and when a simple tonic is required, a more appropriate preparation cannot be substituted.

Pills of Oxide of Zinc.

R. Zinci Oxidi, ʒij.

Confectionis Rosæ, q. s. ut fiant pilulæ x.

Signa.—One three or four times a day. In epilepsy, chorea, &c.

Dr. Chapman speaks favorably of this medicine, and recommends to begin with the above dose, and gradually to increase it.

Pills of the Nitrate of Bismuth.

R. Bismuthi Subnitratiss, ʒj.

Mucilaginis Acaciæ, q. s. ut fiant pilulæ xxx.

One to be taken every two hours. In dyspepsia.

Pills of Nitromuriate of Gold.

R. Auri Nitromuriatis, gr. v.

Pulveris Glycyrrhizæ, ʒjss.

Mucilaginis Acaciæ, q. s. ut fiat massa, in pilulas lxxv. dividenda.

Signa.—One or two thrice a day. In syphilis, &c.

This metal is placed under the head of tonics by Dr. Chapman and we have followed his example. Although once much extolled, it is now seldom used.

[By the term nitro-muriate, the author means what is now known as the chloride of gold. According to the *Codex*, it is prepared by heating in a sand-bath, "one part of fine leaf gold, with three parts of nitro-muriatic acid, and evaporating the solution carefully until the odor of chlorine is perceptible. It is then set aside to cool and crystallize." (DUNGLISON.) The preparations of gold are again coming into use. EDITOR.]

Mustard Seed.

R. Seminum Sinapis Albi, ʒj.

Signa.—A teaspoonful of the unbruised seeds two or three times a day. They are beneficial in costive habits, when the constipation depends upon a torpid state of the muscular coat of the intestines.—PARIS *on Diet*, p. 187.

Electuary of Cinchona, Rust of Iron, &c.

R. Pulveris Cinchonæ,

———— Ferri Subcarbonatis, āā ʒj.

Copaibæ, q. s. ut fiat electuarium.

To be made (by the patient) into pills of the ordinary size, of which four may be taken three times a day.

Pills of the Oxide of Silver.

R. Argenti Oxidi, gr. vj.

Pulveris Glycyrrhizæ, gr. xij.

Syrupi, q. s.

Ut fiat massa in pilulas xij. dividenda.

Signa.—Take one pill three times daily.

Used with much success in pyrosis. Also highly recommended in menorrhagia.

INFUSIONS AND DECOCTIONS.

Infusion of Sage, Boneset, &c.

- R. *Salviæ*,
Eupatorii, āā ʒss.
Cascarillæ, ʒj.

Infuse in three half-pints of boiling water until cold. Dose, a wineglassful every three or four hours. In hectic fever.

Infusion of Colomba and Ginger.

- R. *Colombæ contusæ*, ʒj.
Zingiberis, ʒij.
Aquæ bullientis, Oj.

Fiat infusum.—Give of the strained liquor (cold) a wineglassful every two hours.

This infusion, freely used, is adapted to chronic diarrhœa.

Infusion of Colomba, Rhubarb, &c.

- R. *Cari contusi*,
Radicis Colombæ contusæ,
 ——— *Rhei contusæ*, āā ʒj.
Aquæ ferventis, fʒvijj.

Digest for two hours, and strain.

- R. *Liquoris colati*, fʒijjss.
Tincturæ Rhei, fʒj.
Syrupi Zingiberis, fʒij.

Misc.—Dose, a teaspoonful to a tablespoonful for children, according to their age. In diarrhœa.

Infusion of Wild-cherry Bark.

- R. *Pruni Virginiani*, ʒj.
Aurantii Corticis, ʒij.
Aquæ, Oj.

Infuse the bark alone for six hours, and then add the orange-peel. A wineglassful may be taken every hour or two in consumptive cases, asthma, &c.

Infusion of Virginia Snake-root.

- R. *Serpentariæ*, ʒj.
Aquæ bullientis, Oj.

Infuse in a covered vessel for one hour. A wineglassful every hour or two hours, *pro re nata*.

Infusion of Quassia, &c.

R. Quassiae,
Serpentariæ,
Aurantii Corticis, āā ʒss.
Aquæ bullientis, Oij.

Fiat infusum et cola.—A teacupful to be taken cold, three times a day.

Cold Infusion of Chamomile and Orange Peel.

R. Anthemidis, ʒj.
Aurantii Corticis, ʒss.
Aquæ, Oij.

Macerate for twenty-four hours.—Dose, a teacupful three or four times a day.

The infusion of chamomile made with cold, is in general more grateful to the patient than when made with boiling water. The same remark applies to infusions generally, and they are also less liable to ferment.

Infusion of Hops.

R. Humuli, ʒj.
Aquæ ferventis, Oj.

Infuse for two hours, and strain by expression. Of this a wineglassful may be taken three or four times a day. In dyspepsia.

Compound Infusion of Gentian.

R. Gentianæ concisæ, ʒss.
Aurantii Corticis, ʒij.
Cardamomi, ʒss.
Aquæ ferventis, Oj.

Fiat infusum.

Gentian is an excellent bitter. The above preparation is given in wineglassful doses, in debility of the digestive organs, &c.

Decoction of Angustura Bark.

R. Angusturæ contusæ, ʒj.
Aquæ, Ojss.

Boil for fifteen or twenty minutes and strain. Of this decoction a wineglassful may be taken every two hours.

Decoction of Dog-wood Bark.

R. Cornûs Floridæ contusi, ʒj.

Aquæ, Oj.

Boil for twenty or thirty minutes, and strain.—A wineglassful may be given every hour, as a substitute for Peruvian bark in intermittents, or as an ordinary tonic, every two hours.

Decoction of Bark and Valerian.

R. Cinchonæ Rubræ contusæ, ʒj.

Boil in a pint of water for ten minutes, and strain.

R. Valerianæ contusæ, ʒj.

Aquæ bullientis, Oj.

Infuse for one hour, and strain.—Add the decoction of bark to this infusion, and give a teacupful cold, three or four times a day.

This was a favorite prescription with the late Dr. Parrish, in rheumatic and nervous headache and hemicrania.

Acorn Coffee.

R. Pulveris Glandis Quercûs torrefactæ, ʒj.

Aquæ bullientis, Oj.

Fiat infusum.—Three or four teacupfuls may be taken during the day, and augmented according to circumstances.

The above is a favorite remedy with the Germans. "It is one of the most successful in mesenteric atrophy, commencing rachitis, glandular swellings, asthma, and cough. Continued for a long time, it is one of the most powerful means we possess for destroying the scrofulous disposition."—HUFELAND, *Treatise on Scrofula*, p. 215.

MIXTURES, TINCTURES, &c.

Griffith's Myrrh Mixture.

R. Myrrhæ,

Sacchari, āā ʒj.

Potassæ Carbonatis, gr. xxv.

Tere simul, et adde gradatim—

Aquæ Rosæ, f ʒvijs.

Spiritûs Lavandulæ, f ʒss.

Dein adde—

Ferri Sulphatis crystallizati, in pulverem contriti, ʒj.

Fiat mistura.—Dose, a tablespoonful according to circumstances.

Given as a tonic in phthisis.

Mixture of Sulphuric Acid, &c.

R. Ferri Sulphatis, ʒj.

Acidi Sulphurici, gtt. iv. vel vj.

Sacchari, ʒj.

Aquæ, f ʒiv.

Misce. *Signa*.—A teaspoonful for a dose, to be repeated every two, three, or four hours, much diluted.

Vinous Tincture of Gentian, &c.

R. Pulveris Gentianæ,

Corticis Aurantii, āā ʒss.

Vini Rubri, Oj.

Macerate for three days, and give a wineglassful two or three times a day. In dyspepsia.

Mixture of Sulphate of Iron and Elixir of Vitriol.

R. Ferri Sulphatis, gr. iv.

Acidi Sulphurici Aromatici, gtt. xx.

Aquæ destillatæ, f ʒj.

Fiat mistura. *Signa*.—A teaspoonful to be taken three times a day in a wineglass of water.

Vinegar Draught.

R. Acidi Acetici, f ʒj.

Tincturæ Cardamomi Compositæ, f ʒss.

Syrupi, f ʒss.

Aquæ, f ʒx.

Misce.—To be taken in such portions as the stomach can bear. In sick headache.

Tincture of Cinchona, Valerian, &c.

R. Tincturæ Cinchonæ,

———— Valerianæ, āā f ʒj.

———— Cardamomi, f ʒij.

Aquæ Ment hæ Piperitæ, f ʒiv.

Fiat mistura.—Of this mixture a tablespoonful may be given every third hour. It is especially designed as a tonic in nervous temperaments.

Compound Tincture of Aloes, Gentian, &c.

- R. Aloës, ʒj.
 Pulveris Zedoariæ,
 ——— Gentianæ,
 ——— Croci,
 ——— Rhei,
 ——— Agarici, āā ʒj.
 Syrupi, f ʒij.

To be mixed into a quart of brandy. This is the celebrated Baume de vie, or Elixir of Life. It is a powerful tonic, and one of the most effectual febrifuge medicines. In intermittents, especially, it is justly celebrated. Dose, a tablespoonful three times a day, in water.

Acidulated Tincture of Gentian.

- R. Tincturæ Gentianæ Compositæ, f ʒiv.
 Acidi Sulphurici Aromatici, f ʒss.

Misce.—A small teaspoonful to be taken in sugar and water, three or four times a day.

Huxham's Tincture of Bark.

- R. Tincturæ Cinchonæ Compositæ, f ʒj.

Signa.—A teaspoonful for a dose, every two hours.

In the same doses as the above may be taken,

Tinctura Colombæ—et
 Tinctura Gentianæ Compositæ.

Muriated Tincture of Iron.

- R. Tincturæ Ferri Chloridi, f ʒj.

From five to twenty drops to be taken in a glass of cold chamomile tea, twice or thrice a day. In passive hemorrhage.

Nitrate of Silver Mixture.

- R. Argenti Nitratis, gr. $\frac{1}{4}$ —j.
 Aquæ destillatæ, f ʒij.
 Pulveris Acaciæ, ʒij.
 Sacchari, ʒij.

Fiat mistura. *Signa.*—A teaspoonful every two hours.

Used in the diarrhœa of newly-weaned infants. This is accompanied by the occasional injection of gr. $\frac{1}{4}$ of the salt in mucilage.

HIRSCH.

Solution of Acetate of Strychnia.

R. Strychniæ Acetatis, gr. j.

Acidi Acetici, ℥xx.

Alcoholis, f ʒij.

Aquæ, f ʒvj.

Fiat mistura. *Signa*.—Take ten drops three times a day.

Useful as a tonic, in all cases of nervous exhaustion.

MARSHALL HALL.

Solution of Persesquintrate of Iron.

R. Liquoris Ferri Nitratis, f ʒj.

Syrupi, f ʒj.

Aquæ, f ʒvj.

Fiat mistura. *Signa*.—Give a teaspoonful every three hours.

This preparation is very serviceable in all forms of diarrhœa, but more especially in chronic mucous diarrhœa, hemorrhage from the bowels, uterine hemorrhage, &c.

Sirup of Proto-Nitrate of Iron.

R. Sulphate of Iron, ʒviiij.

Carbonate of Soda, ʒx.

White Sugar, ʒxx.

Nitric Acid (*sp. gr.* 1.42) f ʒv. and f ʒv.

Boiling Water,

Simple Sirup, of each, sufficient.

“Dissolve the sulphate of iron, and carbonate of soda, each in two pints of the water, filter, and add to each solution two ounces of simple sirup. Mix the solutions, and allow the precipitate to subside. Pour off the supernatant liquid, and wash the precipitated carbonate carefully with sweetened water, until the washings have no longer a saline taste. Collect the precipitate upon a fine muslin strainer, and with gentle pressure express as much of the water as possible. Transfer to a porcelain capsule, and add gradually the nitric acid, previously diluted with an equal measure of water. Mix the sugar with the solution, and dissolve over a water-bath, stirring from time to time with a glass rod. When done, the sirup should be made to measure thirty fluidounces, by the addition of a sufficient quantity of water. This sirup contains ten grains of dry nitrate of iron, to each fluidram, and the dose varies from twenty to forty drops.”

LIVERMORE.

Prof. Procter has published another formula, *Journ. Pharmacy*, vol. 23, p. 315.

This preparation is much used in Philadelphia as a substitute for the persesquinitrate of iron, and as the mode of making it is not generally known, the whole process is given.

EDITOR.

Sirup of Iodide of Iron.

R. Liquoris Ferri Iodidi, f ʒss.

Signa.—Take twenty drops, every three hours, in sweetened water.

This valuable preparation is tonic, alterative, and emmenagogue.

Sirup of Iodide of Iron and Manganese.

R. Liquoris Ferri et Mangesiæ Iodidi, f ʒss.

Signa.—Take twenty drops every three hours, in sweetened water.

This is used for the same purposes as the preceding. The salts of manganese are beginning to attract attention. The formula for its preparation, by Prof. Procter, will be found in the *American Journal of Pharmacy*, vol. 25, p. 198.

CLASS XIV.

ALTERATIVES.

ALTERATIVES have been defined—medicines that re-establish the healthy functions of the animal economy, without producing any active evacuation. This class of remedies, therefore, embraces numerous preparations which produce a slow but decided effect on the various secreting organs, sometimes without any sensible increase of the secretions themselves, but in other instances obviously augmenting or magnifying them. Such is the action of minute doses of mercury, iodine, and other substances upon the glandular apparatus. These medicines are generally classed with stimulants and tonics, some of them with narcotics, and others have been placed with those preparations to which they seem most allied when given in an overdose—sialagogues, for example. Without attempting to investigate this question farther than regards practical convenience, we proceed in this place to indicate the more active and important Alteratives; merely premising the familiar truth, that medicines of almost every class of the *materia medica* become alteratives by being administered in very small doses at intervals of a few hours.

PILLS AND POWDERS.

Mercury.

The mercurial preparations constitute the most decided and efficient medicines of this class. In addition to several formulæ inserted under the head of Sialagogues, the following may be appropriately noticed in this place.

Calomel Pills.

R. Hydrargyri Chloridi Mitis, gr. ij.

Confectionis Rosæ, q. s. ut fiant pilulæ xij.

Of which one may be taken every hour or two hours. In *marasmus*, &c. &c.

The use of calomel in minute doses has become very general, and deservedly so. In the hepatic and intestinal complaints of children, it is productive of the happiest results when carefully managed.

In infants, for example, the sixteenth or eighth of a grain, repeated every hour or two hours, according to age and circumstance, will often check diarrhœa, cholera infantum, and dysentery, which have resisted every other treatment. If the discharges are attended by pain and tenesmus, small doses of powdered opium or laudanum may be added; but, as a general rule, the anodyne had better be given by injection.

The preparation called blue-pill (*pilula hydrargyri*) is indicated in similar conditions, and is especially beneficial whenever an alterative is called for. A grain given at bedtime, or even a grain in divided portions through the day, will often produce the desired effect. Hence its extensive use in chronic and obscure derangements of the digestive organs, especially of the stomach and liver; in the whole class of syphilitic affections; glandular enlargements, &c.

In like manner, all the mercurial preparations mentioned under the head of *SIALAGOGUES*, become alteratives by cautious administration in doses more or less minute, and especially when they are given in conjunction with the preparations of *sarsaparilla*.

Pills of Calomel, Quinia, Opium, &c.

R. Hydrargyri Chloridi Mitis, gr. vj.

Pulveris Opii, gr. iij.

Quiniæ Sulphatis, gr. xij.

Syrupi, q. s. ut fiat massa in pilulæ xij. dividenda.

Signa.—One night and morning.

Pills of Blue Mass, Quinia, &c.

R. Pilulæ Hydrargyri,

Quiniæ Sulphatis,

Pulveris Aloës, āā gr. xij.

Syrupi Rhei Aromatici, q. s. ut fiant pilulæ xij.

Signa.—One, three or four times a day.

Either of the above prescriptions is adapted to the condition that follows bilious intermittent or remittent fever. They may be denominated tonic alteratives.

Pills of Red Iodide of Mercury.

R. Hydrargyri Iodidi Rubri, gr. ss.

Extracti Juniperi vel Glycyrrhizæ, gr. viij.

Divide in pilulas viij. *Signa.*—Two in the morning and two at night; dose to be gradually and cautiously augmented.

Corrosive Sublimate Pills.

R. Hydrargyri Chloridi Corrosivi, gr. j.

Solve in aquam destillatam, dein adde—

Micæ Panis,

Sacchari, āā gr. x. vel q. s.

Ut fiant pilulæ xx.

Signa.—Take one pill every four hours as an alterative.

DZONDI.

Arsenic.

Arsenic is a powerful alterative, and one that should be administered with the utmost circumspection. In addition to the formulæ already given, the following may be found useful.

Pills of the Arseniate of Iron.

R. Ferri Arseniatis, gr. iij.

Extracti Lupuli, *Ed.* 3j.

Pulveris Altheæ, 3ss.

Syrupi, q. s. ut fiat massa.

To be divided into forty-eight pills, of which one is given daily.

Prescribed by Dr. Biett in scrofulous, cancerous, and herpetic diseases.

Asiatic Pills.

R. Acidi Arseniosi, gr. j.

Pulveris Piperis Nigri, gr. xij.

——— Acaciæ, gr. ij.

Aquæ, q. s.

Misce, et divide in pilulas xij.

Signa.—Take one pill morning and night. Much used for tuberculous lepra. BIETT.

Iodide of Arsenic Pills.

R. Arsenici Iodidi, gr. j.—ij.

Extracti Conii, ʒij.

Fiat massa, et divide in pilulas xvj.

Signa.—Take one pill morning and night. Given in lepra, impetigo, and cancerous disease.

Pills of Iodide of Silver.

R. Argenti Iodidi,
 Potassæ Nitratis, āā gr. x.
 Tere simul ut fiat pulvis subtilis, dein adde—
 Pulveris Glycyrrhizæ, ʒss.
 Sacchari, ʒj.
 Mucilaginis Acaciæ, q. s.
 Ut fiant pilulæ xl.
Signa.—Take one pill three times daily.

Used successfully in the gastric affections of the Irish peasantry.
 C. PATTERSON.

Pills of Bromide of Iron.

R. Ferri Bromidi pulverizati, gr. xij.
 Confectionis Rosæ, gr. xvij.
 Acaciæ, gr. xij.
 Misce, et fiant pilulæ xx. Two pills to be taken in the morning, and two in the evening, or, one three times a day. In hypertrophy of the heart, and scrofulous affections.

Dr. Robert Dick strongly recommends the use of the bromide of iron in dyspeptics with strumous habits.

MAGENDIE.

Pills of Calomel and Ox Gall.

R. Hydrargyri Chloridi Mitis, ʒj.
 Fellis Bovini inspissati, gr. xv.
 Saponis, gr. x.
 Extracti Taraxaci, ʒss.
 Fiat massa in pilulas xx. dividenda.
Signa.—Take two pills at bedtime.

These pills are alterative and deobstruent.

Brucia.

The alkaline principle called Brucia, or Brucine, is contained in the false *Angustura* bark, *nux vomica*, and *St. Ignatius's* bean. It possesses the properties of *strychnia*, but in less degree. It may be administered in doses of half a grain to a grain; but the minimum dose should be first given, and the quantity gradually augmented.

Pills of Brucia.

R. Bruciæ, gr. xij.

Confectionis Rosæ, q. s. ut fiat massa.

Divide in pilulas xxiv.

Signa.—One pill to be taken four times a day, and gradually increased.

Burnt Sponge.

R. Spongiæ Ustæ, ʒj.

Sacchari ʒj.

Misce, et divide in chartulas vj.—One three times a day, gradually augmenting the dose. In bronchocele.

It is well known that iodine was first discovered in sea-weed and sponges; and the latter having long had the reputation of being a cure for goitre, it was found that iodine was the active remedial agent. We give above a prescription for the use of Sponge as formerly in vogue, but now nearly superseded by the preparations of pure iodine.

TINCTURES, MIXTURES, &c.

Preparations of Iodine.

That iodine is one of the most powerful alteratives at present known, there can be no question. There are few diseases dependent upon or connected with morbid secretion of the glandular structures, but what are more or less modified by its use, while in others it acts as a direct curative agent. It is chiefly, however, in scrofulous, glandular, and cutaneous affections that it acts most beneficially, dispersing indurations of the glands, and restoring the secretions of the cutaneous and mucous apparatus; whence its advantages in eruptive diseases, bronchitis, leucorrhœa, amenorrhœa, &c. &c. It has been found to restore suppressed hemorrhoidal and catamenial discharges, and when pushed too far, has even produced alarming hemorrhage of the lungs, nose, and uterus. These facts should make the practitioner extremely cautious in its administration, beginning with small doses, and watching their effects. Should it produce any of the inconveniences already mentioned, or dizziness, nausea, purging, burning of the skin or swelling of the gums, it should be at once suspended, or given less frequently and in smaller doses.

Ethereal Tincture of Iodine.

R. Iodinii, gr. vj.
 Ætheris, fʒij.

Misce.—Thirty drops contain one grain of iodine.—Dose, ten drops two or three times a day, largely diluted. In cutaneous and glandular diseases.

Compound Tincture of Iodine.

R. Iodinii, ʒss.
 Potassii Iodidi, ʒj.
 Alcoholis, Oj.

Fiat tinctura.—Dose, five to fifteen drops, *pro re nata*.

Mixture of Iodide of Potassium, &c.

R. Potassii Iodidi, gr. x. vel ʒj.
 Magnesiae Sulphatis, ʒss.
 Potassae et Antimonii Tartratis, gr. ss.
 Aquae destillatae, fʒvj.

Misce. *Signa*.—A teaspoonful three or four times a day. In scrofula.

Saturated Tincture of Iodine.

R. Iodinii, ʒij.
 Alcoholis, fʒj.
 Spiritûs Lavandulae Compositi, fʒij.

Fiat solutio.—This is Mr. Durand's saturated solution. The dose is ten or twenty drops, which may be gradually increased, watching its effects.

Iodine with Iodide of Potassium.

R. Iodinii, gr. iij.
 Potassii Iodidi, gr. vj.
 Aquae destillatae, fʒj.

Solve. *Signa*.—Dose, six to ten or twenty drops thrice a day in a draught of cold water.

If it produce dizziness, pain in the bowels, or other unpleasant symptoms, diminish the quantity.

The author of *Illustrations of Pulmonary Consumption*, strongly recommends this formula in catarrh, phthisis, &c. It has a powerful effect in glandular disease, and all affections of the mucous membranes.

S. G. MORTON.

Mixture of Tincture of Iodine, &c.

- R. Tincturæ Iodinii, f ʒj.
 Mucilaginis Acaciæ, f ʒij.
 Aquæ destillatæ, f ʒvj.

Fiat mistura. *Signa*.—A tablespoonful every two hours. Employed in cases of ulceration accompanied by purulent discharge from the meatus auditorius, and in the scrofulous diathesis.

Solution of Iodide of Iron.

- R. Ferri Iodidi, ʒj.
 Aquæ destillatæ, f ʒj.

Fiat solutio.—Dose, six to ten drops, morning, noon, and night, in a wineglass of cold water.

Solution of Iodide of Potassium.

- R. Potassii Iodidi, gr. xxxvj.
 Aquæ destillatæ, f ʒj.

Fiat solutio.—Dose, ten to twenty drops, morning, noon, and night, in a wineglass of sweetened water. Each dose will contain between one and two grains of the iodide.

MAGENDIE.

Lugol's Solution.

- R. Liquoris Iodinii Compositi, f ʒss.

Signa.—Give ten drops in sweetened water three times a day.

Magendie's Anti-epileptic Iodine Solution.

- R. Potassii Iodidi, ʒiv.
 Iodinii, gr. ij.
 Aquæ Menthæ Piperitæ, f ʒvj.

Fiat solutio.—Dose, a teaspoonful thrice per diem.

Mixture with Iodide of Potassium.

- R. Potassii Iodidi, ʒss.
 Syrupi Zingiberis, f ʒj.
 Aquæ, f ʒv.

Fiat mistura. *Signa*.—Take a tablespoonful three times a day.

Solution of the Ioduretted Hydriodate of Potash.

- R. Iodinii, ʒj.
 Potassii Iodidi, ʒij.
 Aquæ destillatæ, f ʒvij.

Fiat solutio.—Dose, six drops, morning and evening, in a wineglass of water. In scrofulous and scirrhus diseases.

DUNGLISON'S *New Remedies*, 6th ed. p. 598.

Mixture of Iodo-hydrargyrate of Potash.

- R. Potassii Iodidi, gr. iijss.
 Hydrargyri Iodidi Rubri, gr. ivss.
 Aquæ destillatæ, f 3j.

Dissolve first the iodide of potassium, and then the biniodide of mercury, in the water. The compound salt in this solution may be assumed at eight grains, although there is a small excess of the iodide of potassium. The dose is from two to five drops, containing from the thirtieth to the twelfth part of a grain, which may be repeated three times a day, much diluted.

DR. CHANNING.

Ioduretted Sulphuric Ether.

- R. Ætheris, f 5j.
 Iodinii, gr. vj.

Solve. *Signa.*—Six drops constitute a dose. To be largely diluted when taken.

Thirty drops of this solution contain a grain of iodine.

Iodine Waters.

The physicians of Europe, and especially of Paris, have made extensive and salutary use of iodine waters, both internally and externally. In these, the iodine is very much diluted, and generally with the addition of more or less common salt. These waters are prepared on a large scale and with great precision by the pharmaceutists of Paris; but they have been but partially introduced into this country. M. Magendie, however, gives formulæ for some extemporaneous iodine waters, which may answer all the purposes of the more elaborate preparations. The following is an example:—

- R. Potassii Iodidi, gr. vj.
 Iodinii, gr. j.
 Aquæ, Oij.

Fiat solutio.—This solution is directed to be drank at meals, in place of common water.

Donovan's Solution.

- R. Liquoris Arsenici et Hydrargyri Iodidi, f 5ss.

Signa.—Give ten drops three times a day in sweetened water.

Used in various skin diseases, and for venereal eruptions.

Bromine.

This substance, which is allied to chlorine and iodine, in its habitudes, is obtained from salt springs and other mineral waters. In medicine, it is chiefly used in combination with potassa.

Solution of Bromine.

R. Brominii, f ʒss.

Aquæ, f ʒijss.

Misce. *Signa.*—Dose, five drops three times daily.

POURCHE.

Solution of Bromide of Potassium.

R. Potassii Bromidi, ʒj.—ij.

Syrupi Aurantii Corticis, f ʒj.

Aquæ, f ʒiiij.

Fiat mistura. *Signa.*—Give a tablespoonful three times a day.

Mixture of Iodine and Arsenic.

R. Liquoris Iodinii Compositi, f ʒij.

———— Potassæ Arsenitis, f ʒj.

Misce. *Signa.*—Take five drops three times daily.

Pearson's Solution.

R. Sodæ Arseniatis, gr. j.

Aquæ, f ʒj.

Fiat solutio. *Signa.*—Dose, twelve to twenty drops, three times a day.

Used in the scaly and other skin diseases.

An arseniate of ammonia was introduced by Bielt, and employed by him in the same strength as the foregoing solution, *i. e.* one grain to the fluidounce, and given in the dose of twenty drops. Its virtues are similar to those of Fowler's and Pearson's solutions.

Phosphate of Ammonia.

R. Ammoniæ Phosphatis, ʒss.

Aquæ destillatæ, f ʒvj.

Fiat solutio.—A tablespoonful to be taken every four hours.

The preceding formula is that of Dr. Buckler, who recommends it in those cases of gout and rheumatism in which little acid is present in the urine; as well as in all the chronic modifications of those diseases, especially when they are associated with chronic thickening of the white tissues.

Strychnia.

We have (under the head of Narcotics) noticed several prescriptions for using this extremely poisonous substance; but as

it is chiefly given as an alterative, we now add some additional modes of administration. Strychnia is much in vogue in the whole class of nervous diseases, paralysis, whether general or local, neuralgia, chorea, hysteria, &c. It is also employed in syphilis, amenorrhœa, dyspepsia, bronchitis, and, in fact, in all cases where an alterative course is indicated.

Solution of Acetate of Strychnia.

R. Strychniæ Acetatis, gr. iij.

Alcoholis, f 3j.

Aquæ Cinnamomi, f 3vij.

Misce.—Dose, five drops twice a day, cautiously increasing it.

DUNGLISON, *New Remedies*, 6th ed. p. 664.

Tincture of Acetate of Strychnia.

R. Strychniæ Acetatis, gr. jss.

Alcoholis, f 3ss.

Fiat tinctura.—Dose, five to ten drops thrice per diem. In syphilitic pains of the bones, &c. DUNGLISON, *ut supra*.

A sulphate of strychnia is also in use. It is prepared like the acetate, of course substituting sulphuric acid. It has no advantages over the preparations already given, and is little used.

See MAGENDIE, *Formulaire*, 9ème éd. p. 30.

Mixture of Brucia.

R. Bruciæ, gr. vj.

Aquæ destillatæ, f 3iv.

Sacchari, 3ij.

Fiat mistura. Dose half a tablespoonful night and morning. This and the following preparation are used in paralysis and the entire series of spasmodic affections.

Tincture of Brucia.

R. Bruciæ, gr. xvij.

Alcoholis, f 3j.

Fiat tinctura.—From six to ten drops may be taken in a draught of water.

Cod-Liver Oil (Gadus Morrhua).

This preparation, which is now divested in a great measure of its nauseous smell and taste, is given in doses of half an ounce, three or four times a day. To children, a teaspoonful. It is of much service in all strumous affections, and in the

chronic forms of gout and rheumatism. It may be taken alone, or in milk; most persons prefer taking it on the froth of porter. Others swallow it in coffee, in lemon-juice, or with some of the aromatic waters.

Dr. Bradshaw recommends a thin infusion of flaxseed, flavored with lemon-peel, and sweetened to please the palate. He says: "I pour a small quantity of infusion into a glass, on which is added the oil, and again another small quantity of infusion upon it. In this manner, the dose may be swallowed, and will slip down without offending the palate, and generally is well retained even by irritable stomachs." The same author says: "Another mode has been successful when many have failed, and will be found equally valuable: It is the yolk of an egg beat up with boiling water and sugar, with which may be mixed a dessertspoonful of brandy; and on this the dose may be given without producing that nausea so frequently experienced when conveyed by other vehicles."

In the Editor's experience, the form of emulsion has proved to be one of the least satisfactory modes of exhibiting this article. He is in the habit of administering it in the pure state, directing the patient to chew a piece of orange-peel before and after each draught; or, what is equally effectual, to rinse the mouth, and gargle the throat with vinegar, before and after the dose.

Mixture with Cod-Liver Oil.

R. Olei Morrhuæ, f3ss.
Liquoris Potassæ, gtt. xl.
Aquæ Ment hæ Piperitæ, f3ss.
Misce, et fiat haustus.

The draught to be washed down with a teaspoonful of lemon-juice to liberate the oil on the stomach. DUNGLISON, *New Rem.* 6th ed. p. 557.

PERCIVAL.

Mixture of Iodine and Cod-Liver Oil.

R. Iodinii, gr. jss.
Olei Morrhuæ, f3v.
Rub the iodine with a dram of the oil, and then carefully add the remainder.
Signa.—Dose, a tablespoonful three times a day.

The quantity of iodine can be increased without materially changing the appearance of the solution. FLEISCHMANN.

Mixture with Corrosive Sublimate.

R. Hydrargyri Chloridi Corrosivi, gr. ij.

Aquæ destillatæ, fʒvj.

Spiritûs Cinnamomi,

Syrupi, āā fʒj.

Misce.—Dose, one or two large spoonfuls, twice or thrice a day.
In venereal cases.

Nitro-muriatic Acid.

R. Acidi Nitrici, fʒss.

— Muriatici, fʒj.

Aquæ, fʒviij.

Misce, et adde—

Spiritûs Ætheris Nitrici, fʒj. vel. fʒij.

Fiat mistura. *Signa.*—From twenty drops to half a tea-spoonful to be given every three hours, in a wineglass of sweetened water.

These acids, in the same proportions, are also employed as a bath in hepatic derangements. To accomplish this end, put a teaspoonful of the compound acid into a quart of warm water, and apply it to the region of the liver night and morning, by means of a soft sponge.

Diluted Nitric Acid.

R. Acidi Nitrici, fʒss. vel fʒj.

Aquæ, Oij.

Sacchari, ʒj.

Fiat mistura.—One-eighth of this quantity may be taken daily, in divided doses. In lues venerea.

FERRIAR, *Med. Hist.*

Mixture of Nitric Acid.

R. Acidi Nitrici, fʒss.

Pulveris Acaciæ,

Sacchari, āā ʒij.

Aquæ, fʒviij.

Fiat mistura. *Signa.*—A small tablespoonful largely diluted, and sweetened, *pro re nata*.

Nitric acid may be prescribed as a tonic in the above form, or simply diluted with water. It is best taken through a quill, as the teeth suffer when it comes in contact with them. It sometimes arrests the colliquative sweats of hectic fever when other remedies fail.

Artificial Harrowgate Water.

R. Potassæ Sulphatis cum Sulphure, *Ed.* 3j.

—— Bitartratis, 3ss.

Magnesiæ Sulphatis, 3vj.

Aquæ destillatæ, Oij.

Solve. *Signa.*—One-half, taken in draughts, constitutes a dose.

The above ought to be taken in the morning before breakfast, and be followed by a walk, to produce the desired effect.

Antacid Tincture.

R. Pulveris Guaiaci, 3j.

Terebinthinæ Canadensis, 3j.

Hydrargyri Chloridi Corrosivi, ʒj.

Olei Sassafras, f3ij.

Alcoholis, f3viij.

Dissolve the corrosive sublimate in one-half the spirit, and then add to it the other ingredients.—Of this from ten to twenty drops may be taken morning and evening, in wine or water.

This form of exhibiting corrosive sublimate is employed by Dr. Emerson and other practitioners of this city, in lues venerea, &c.

CLASS XV.

ASTRINGENTS.

DOCTOR CULLEN describes astringents to be "such substances as, applied to the human body, produce contraction and condensation in the soft solids, and thereby increase their density and force of cohesion." "And by the operation of this corrugating power," adds Dr. Murray, "either directly exerted on a part, or extended by sympathetic action, the morbid affections arising from a state of relaxation are supposed to be removed."

In the use of this class of medicines a caution suggests itself; namely, that they should not be given to arrest evacuations that are critical, and designed by nature to relieve a plethoric state of the system, unless such an evacuation exists to an alarming extent.

POWDERS, PILLS, &c.

Compound Powder of Alum.

R. Aluminis, ℥j.

Pulveris Kino, gr. v.

Fiant pulveres v. *Signa.*—One every two hours. In obstinate diarrhoea.

This compound, though much employed by some practitioners, contains substances which would be incompatible with each other if given in solution. In menorrhagia, &c.

Powder of Sugar of Lead and Calomel.

R. Hydrargyri Chloridi Mitis, gr. j. vel ij.

Plumbi Acetatis, gr. ss. ad gr. j.

Fiant pulveres iv.—One every three hours. In cholera infantum.

The above dose is adapted to any period between ten and twenty months of age.

DR. T. D. MITCHELL.

Powder of Sugar of Lead, Calomel, and Ipecacuanha.

- R. Hydrargyri Chloridi Mitis,
 Pulveris Ipecacuanhæ, āā gr. ij.
 Plumbi Acetatis, gr. viij.
 Misce, et divide in pulveres viij.
Signa.—Give one powder every three hours.

Used in cholera infantum. The Editor, from his connection with a large charitable institution, has had repeated opportunities of testing the various astringent preparations employed in this complaint, and he has found none to answer better than this formula, either as it stands, or with the addition of twenty-four grains of prepared chalk. The quantity of sugar of lead appears large, but experience proves that it may be given, both with safety and benefit, to a child from one to three years old.

D. F. CONDIE.

Powder with Alum and Opium.

- R. Aluminis, ʒss.
 Pulveris Opīi, gr. iij.
 Fiant pulveres vj. *Signa.*—One to be taken every three or four hours. In diarrhœa.

Powders of Nitrate of Bismuth, &c.

- R. Bismuthi Subnitratis, ʒj.
 Pulveris Acaciæ, ʒss.
 Magnesiæ, ʒj.
 Misce, et divide in pulveres xij.
Signa.—Give one powder every four or six hours.

Used in the diarrhœa of phthisis.

T. THOMSON.

Powders of Nitrate of Bismuth.

- R. Bismuthi Subnitratis, gr. iij.—vj.
 Sacchari, gr. x.
 Fiant pulveres vj.
Signa.—Place a powder on the tongue of a child (one year old), three or four times daily.

Used in the diarrhœa of children.

RECAMIER, TROUSSEAU.

Pills of Acetate of Lead and Calomel.

R. Plumbi Acetatis, ʒss.

Hydrargyri Chloridi Mitis, gr. v.

Confectionis Rosæ, q. s.

Ut fiant pilulæ x.—One to be given every two, three, or four hours, according to circumstances. In hemorrhage, and especially in hematemesis.

Pills of Sugar of Lead and Opium.

R. Plumbi Acetatis, ʒj.

Opii, gr. j.

Misce, et divide in pilulas xij.—A pill to be taken every hour at first, and then every two hours. This is Dr. Graves's prescription for the treatment of cholera.

Pills of Sugar of Lead and Opium.

R. Plumbi Acetatis, gr. xij.

Pulveris Opii, gr. vj.

Confectionis Rosæ, q. s. ut fiat massa in pilulas vj. dividenda.

One to be taken for a dose, according to circumstances. In hæmoptysis and other hemorrhages.

Pills of Alum, Catechu, &c.

R. Aluminis, gr. vj.

Extracti Opii, gr. j.

Catechu, gr. vj.

To be divided into six pills, one of which is to be given every two, three, or four hours. In passive hemorrhage, and in atonic mucous discharges.

Pills of Rhatany.

R. Extracti Krameriæ, ʒj.

Pulveris Kino, q. s.

Ut fiant pilulæ viij. *Signa.*—A pill to be given every hour or two hours. In the profluvia.

Pills of Creasote.

R. Creasoti, gtt. x.

Pulveris Glycyrrhizæ,

Mucilaginis Acaciæ, āā q. s.

Ut fiant pilulæ xx.—Dose, a pill thrice in the day. In bronchitis, neuralgia, &c.

Pills of Kino, Opium, &c.

R. Pulveris Kino, gr. xx.

————— Opii, gr. ij.

Mucilaginis Acaciæ, q. s. ut fiant pilulæ iv.

Of which take one every four hours. In diarrhœa, pyrosis, &c.

Monesia.

This substance is the product of a tree that grows in Chili, but its botanical affinities are unknown. It is given in the form of extract, twelve or twenty grains daily, divided into three-grain pills. There is also a tincture of monesia, of which the dose is fʒj. twice *per diem*. In leucorrhœa, diarrhœa, menorrhagia, &c. &c.

Tannin Pills.

R. Acidi Tannici, gr. viij. vel xij.

Mucilaginis Acaciæ, q. s.

Ut fiant pilulæ viij. One may be given every two or three hours. In gonorrhœa, &c.

Pills of Tannin and Morphia.

R. Acidi Tannici, ʒj.

Morphiæ Sulphatis, gr. j.

Mucilaginis Acaciæ, q. s.

Ut fiant pilulæ x.

One to be given every two or three hours. Blue mass may be added according to circumstances. In diarrhœa and dysentery.

Pills of Gallic Acid.

R. Acidi Gallici, ʒj.

Extracti Gentianæ, gr. x.

Syrupi, q. s.

Fiat massa in pilulas x. dividenda.

Signa.—Give one pill every three hours. Used in hæmaturia, and other hemorrhages.

Bolus with Alum and Extract of Bark.

R. Aluminis,

Extracti Cinchonæ Rubræ,

Myristicæ, āā ʒss.

Syrupi, q. s. ut fiat bolus.

To be given in uterine and other hemorrhages, and repeated *pro re nata*.

MIXTURES, INFUSIONS, &c.

Infusion of Matico.

R. Matico concisi, ʒj.

Aquæ bullientis, Oj.

Macera per horas duas et cola.

Signa.—Take a wineglassful every two or three hours.

This has been used to check the discharge in hæmoptysis, hæmatemesis, and dysentery.

Parrish's Camphor Mixture.

R. Aquæ Camphoræ, fʒiij.

Spiritûs Lavandulæ Compositi, fʒj.

Sacchari, ʒj.

Fiat mistura. *Signa.*—Give a tablespoonful every two hours in diarrhœa and cholera morbus, adding ten drops of laudanum when there is much pain.

JOS. PARRISH.

Camphor, Laudanum, and Nitric Acid, or Hope's Mixture.

R. Aquæ Camphoræ, fʒiv.

Acidi Nitrici, gtt. iv.

Tincturæ Opii, gtt. xl. vel lx.

Fiat mistura.—Dose, a tablespoonful every two hours. In diarrhœa and dysentery.

This is a well known and very efficacious preparation analogous to *Hope's mixture*. If the nitric acid is added in larger proportion, it is liable to defeat the object of the prescription. The original formula contained a fluidram of nitrous acid, forty drops of laudanum, and eight fluidounces of camphor water, and was given in doses of two fluidounces every three hours.

HOPE.

Chalk Mixture.

R. Cretæ Præparatæ, ʒiss.

Sacchari,

Acaciæ, āā ʒj.

Aquæ destillatæ, fʒiv.

Olei Cinnamomi, gtt. ij.

Tincturæ Opii, gtt. xl. vel lx.

Misce.—Dose, a tablespoonful every two hours. In diarrhœa and dysentery.

When a strongly astringent chalk mixture is desirable, we add to the above formula Tinct. Kino, fʒij. or the same quantity of tincture of krameria, catechu, or nutgalls.

Jackson's Mixture.

R. Spiritûs Lavandulæ Compositi,
Tincturæ Camphoræ, āā f̄ss.

————— Opii,

Spiritûs Ætheris Compositi, āā f̄ij.

Misce. *Signa*.—Give twenty drops at short intervals, in cholera morbus, and the early stages of cholera, to allay nausea and relieve pain. S. JACKSON.

Although this preparation is not strictly astringent, yet it may be classed with great propriety with Hooper's and Parrish's mixtures.

Infusion of Galls.

R. Gallæ contusæ, ʒij.

Aquæ bullientis, f̄xx.

Infuse for four hours, and strain.—To be taken in doses of a tablespoonful every hour or two hours. In diarrhoea. It may be rendered more agreeable by the addition of some aromatic.

Infusion of Galls and Chalk.

R. Acaciæ, ʒj.

Cretæ Præparatæ, ʒij. vel ʒss.

Infusi Gallæ, f̄iv.

Tincturæ Opii, f̄ss.

Fiat mistura.—Dose, a tablespoonful every two hours. In diarrhoea.

Infusion of Logwood.

R. Hæmatoxyli concisi, ʒss.

Aquæ bullientis, Oj.

Infuse for two or three hours, and strain. Of this a tablespoonful is given to a child every two or three hours. In cholera and diarrhoea.

Mixture of Extract of Logwood, &c.

R. Extracti Hæmatoxyli, ʒiij.

Tincturæ Catechu, f̄ij.

Aquæ, f̄vij.

Misce. *Signa*.—Two tablespoonfuls to be given once in three or four hours. In the advanced stages of diarrhoea and dysentery.

Mixture with Extract of Bark and Alum.

- R. Extracti Cinchonæ Rubræ, ʒss. vel ʒj.
 Aluminis, ʒij.
 Aquæ Cinnamomi, fʒiij.
 Syrupi, fʒj.

Misce. *Signa*.—Two teaspoonfuls every two hours. In low diseases accompanied with hemorrhage.

Mixture with Tincture of Catechu, &c.

- R. Tincturæ Catechu, fʒj.
 ——— Opii, gtt. lx.
 Acaciæ, ʒij.
 Aquæ Cinnamomi, fʒvj.

Misce. *Signa*.—A tablespoonful every two hours. In diarrhoea.

Creasote Mixture.

- R. Creasoti, gtt. vj.
 Mucilaginis Acaciæ, fʒiv.
 Sacchari, ʒj.

Fiat mistura.—Dose, a tablespoonful diluted, every two or three hours. In hemorrhages.

Alcoholic Solution of Creasote.

- R. Creasoti, gtt. j.
 Alcoholis, gtt. xvj.

Misce.—A drop or two of this solution dropped into the cavity of an aching tooth, is said to afford immediate relief.

MARINUS.

Electuary of Catechu and Opium.

- R. Electuarii Catechu, *Ed.* ʒj.

Suited to cases of diarrhoea, in teaspoonful doses. Ten scruples contain one grain of opium.

Alum Whey.

- R. Lactis Vaccinæ bullientis, Oj.
 Pulveris Aluminis, ʒij.

Boil them until the coagulum separates, and strain. A cupful to be taken occasionally. In diabetes, &c.

Peruvian Bark with Lime-water.

- R. Pulveris Cinchonæ Rubræ, ʒss.
 Liquoris Calcis, fʒviiij.

Misce. *Signa*.—A tablespoonful every two hours, or oftener if necessary. In diarrhoea and menorrhagia.

Sirup of Galls, Brandy, &c.

R. Gallæ contusæ, ʒij.

Spiritus Vini Gallici, fʒviij.

Mix.—Burn out the spirit; then add—

Corticis Cinnamomi,

Macis, āā ʒij.

While hot, add sugar by burning it over the inflamed brandy until a sirup is formed. Decant the pure sirup. Dose, a tablespoonful every two hours. In chronic and obstinate diarrhœa.

JOS. PARRISH.

Solution of Sulphate of Copper, &c.

R. Cupri Sulphatis, gr. iiij.

Acidi Sulphurici, gtt. x.

Aquæ destillatæ, ʒj. vel fʒij.

Fiat solutio. *Signa.*—From twenty to forty drops to be given every one or two hours in sirup, *pro re nata*. In menorrhagia.

Mixture of Extract of Rhatany, &c.

R. Extracti Krameriæ, ʒj.

Syrupi Papaveris, *Lond.*

Aquæ Rosæ, āā fʒij.

Fiat solutio. *Signa.*—A teaspoonful for a dose, *pro re nata*. In passive hemorrhage and chronic dysentery.

M. FOUQUIER.

Solution of Acetate of Lead.

R. Plumbi Acetatis, gr. v.

Aceti, gtt. v.

Sacchari, ʒj.

Aquæ, fʒj.

Fiat mistura. *Signa.*—Give a teaspoonful to a child every hour or two, to allay the vomiting in cholera infantum.

D. F. CONDIE.

Enema of Acetate of Lead, &c.

R. Plumbi Acetatis, ʒj.

Tincturæ Opii, gtt. lx.

Aquæ tepidæ, fʒij.

Fiat solutio.—To be thrown up the rectum in uterine hemorrhage.

DEWEES'S *Syst. of Mid.* p. 420.

CLASS XVI.

ABSORBENTS, OR ANTACIDS.

ABSORBENTS or Antacids form a class of remedies for correcting acidity in the stomach and bowels. As a general rule they should be preceded by an emetic or cathartic, or both. When a permanent effect is desired, they are usually combined with tonics, aromatics, and sometimes with narcotics. When given alone, they too frequently afford but transient relief.

POWDERS AND PILLS.

Compound Cretaceous Powder with Opium.

- R. Cretæ Præparatæ, ℥iss.
Pulveris Cinnamomi, ℥j.
——— Piperis Longi, gr. viij.
——— Opii, gr. vj.

Divide in chartulas xij.—One three or four times a day. In diarrhœa.

The above formula, with a slight alteration, is from the *London Pharmacopœia*.

Powder with Crab's Claws and Rhubarb.

- R. Chelæ Cancrorum, ℥j.
Pulveris Rhei, ℥ij.

Misce, et divide in pulveres iv.—Of which one may be taken once or twice a day, according to circumstances. In acidity of the primæ viæ.

This preparation is called by the several names of crab's claws, eyes, and stones. It is a carbonate of lime united with some animal gelatine, which, like prepared oyster-shells, has little or no superiority over the prepared chalk.

Powder with Magnesia, Fennel-seed, &c.

R. Magnesiae, ʒss.

Pulveris Fœniculi,

———— Cinnamomi, āā ʒj.

Fiat pulvis.—Dose, a small teaspoonful three or four times a day.

Powder with Soda, Magnesia, &c.

R. Magnesiae, ʒss.

Sodæ Bicarbonatis,

Pulveris Zingiberis, āā ʒj.

Misce. *Signa*.—A small teaspoonful occasionally. In cardialgia.

Powder with Magnesia and Colombo.

R. Magnesiae, ʒjss.

Pulveris Colombæ, ʒj.

Misce.—Divide in chartulas vj.—Of which one may be taken three times a day, in sugar and water. In cardialgia.

Pills of Soda, Rhubarb, &c.

R. Pulveris Rhei,

Sodæ Carbonatis exsiccatae,

Extracti Gentianæ, āā ʒj.

Hydrargyri Chloridi Mitis, gr. iij.

Fiant pilulæ xx. *Signa*.—Two occasionally. In dyspepsia.

MIXTURES, SOLUTIONS, &c.

Chalk Mixture.

R. Cretæ Præparatae, ʒiss. vel ʒij.

Sacchari,

Pulveris Acaciæ, āā ʒj.

Aquæ Menthæ Piperitæ, f ʒvj.

Fiat mistura.—A tablespoonful for a dose every two hours. When administered in diarrhoea, astringents and laudanum are added.

Solution of Salt of Tartar.

R. Potassæ Carbonatis, ʒj.

Sacchari, ʒj.

Aquæ Menthæ Piperitæ, f ʒiv.

Spiritûs Lavandulæ Compositi, f ʒij.

Tincturæ Opii, gtt. xl.

Misce. *Signa*.—A tablespoonful every hour or two hours. In sickness of the stomach occasioned by acidity.

The opium in the above prescription may be omitted at the discretion of the practitioner.

Mixture of Salt of Tartar, Ammonia, &c.

R. Spiritûs Ammoniaë Aromatici, f ʒj.

Potassæ Carbonatis, ʒij.

Aquæ Cinnamomi, f ʒiv.

Misce. *Signa.*—A large teaspoonful for a dose *pro re nata*. In pyrosis.

Lime-water and Milk.

R. Liqûoris Calcis,

Lactis recentis, āā f ʒiij.

Misce. *Signa.*—One or two tablespoonfuls for a dose. In sickness of the stomach few remedies are more prompt in their effect. An antacid remedy admirably adapted to children, is prepared by pouring four ounces of boiling-water on a heaped teaspoonful of calcined magnesia, and giving a dessertspoonful of the clear solution every hour or two. It is highly aperient.

Mixture with Ammonia, Magnesia, &c.

R. Magnesiae, ʒj.

Spiritûs Ammoniaë Aromatici, f ʒj.

—— Cinnamomi, *Lond.* f ʒiij.

Aquæ, f ʒvj.

Misce. *Signa.*—A tablespoonful every two hours. In the sickness accompanying pregnancy.

Mixture of Magnesia, Camphor, &c.

R. Magnesiae, ʒj.

Camphoræ, ʒss.

Sacchari,

Acaciaë, āā ʒij.

Ætheris, f ʒss.

Aquæ destillatæ, f ʒiv.

Misce. *Signa.*—A tablespoonful for a dose four or five times a day. In flatulency and irritable stomach.

Mixture of Colombo, &c.

R. Colombæ, ʒss.

Coque in Aquæ, f ʒviiij. ad f ʒv.

Cola et adde—

Potassæ Carbonatis, ʒss.

Succi Limonis, f ʒiij.

Tincturæ Opii, gtt. xij.

Misce.—From a tea to a tablespoonful may be given every hour. Anti-emetic.

A hot infusion of bruised colombo, is preferable to the above decoction, inasmuch as it contains all the virtues of the root, without the starch.

EDITOR.

Anti-emetic Mixture.

R. Tincturæ Aurantii, *Loul.* f 3ss.

———— Aloës,

———— Castorei, āā f 3j.

Misce. *Signa.*—Dose, from forty to fifty drops three or four times a day.

In continued or chronic vomiting, occasioned by atony or spasm of the stomach, the above is strongly recommended by Hufeland in his Treatise on Scrofula.

Mixture with Carbonate of Soda, &c.

R. Sodæ Carbonatis, ʒiv.

Infusi Gentianæ Compositæ,

Aquæ Cinnamomi, āā f 3iij.

Tincturæ Cardamomi, f 3ss.

Fiat mistura. *Signa.*—A tablespoonful every two or three hours. In acidity, flatulence of the stomach, &c.

Dyspeptic Lye.

Take of

Hickory ashes,	1 quart,
Soot,	6 ounces.
Boiling water	1 gallon.

Mix, and let them stand for twenty-four hours, frequently stirring the ingredients. A teacupful may be given of this liquor (decanted), three times a day. It should be decanted always at the end of twenty-four hours, for, by standing on the materials a long time, the water acquires *caustic properties*, and may do serious injury.

This domestic remedy we have thought best to give, as we received it, in the English language. It deserves an insertion, from the circumstance that it was employed by the late Dr. Physick in his own case, and was of decided advantage when the usual alkaline remedies had failed.

BOOK II.

OF

EXTERNAL APPLICATIONS.

CLASS XVII.

CAUSTICS, OR ESCHAROTICS.

CAUSTICS or cauterics are divided into the actual and potential. The first consists of fire itself; and the second, of those substances which destroy the living solids, either by excessive stimulation, or, as some suppose, by producing a chemical decomposition.

Firing.

A modified form of the actual cautery has attracted some attention in Europe, from its great efficacy in the relief of neuralgic and other pains, paralysis, sciatica, and chronic rheumatism.

Dr. Corrigan uses a simple instrument, consisting of a thick iron wire shank, two inches long, inserted in a small wooden handle, and having on its extremity, which is slightly curved, a button of iron a quarter of an inch thick, and half an inch in diameter, the whole instrument being only six inches long. The face of the button for application is *quite flat*. If the instrument be larger than this, it will require too much time to heat it; if smaller, it will not retain the heat a sufficient length of time. In using the instrument, it is necessary to hold the button over the flame of a spirit-lamp, keeping the forefinger, which holds the wire, at the distance of half an inch from the button. As soon as the finger feels uncomfortably hot, the instrument is ready for use. It is applied as quickly as possible, the skin being tipped successively, at intervals of half an inch, over the whole affected part, as lightly and rapidly as possible, always taking care to bring the flat surface of the button fairly in contact with the skin. In this way, the process of firing a whole limb, or the loins, making about one hundred applications, does not occupy a minute. You can ascertain at once whether the heat be sufficient. If you look sideways at the spots as you touch them, you will observe that each spot the

iron has touched immediately becomes of a glistening white. The iron is never redhot, and does not make an eschar.—BRAITHWAITE'S *Retrospect*, No. 13, p. 55.

The use of the actual cautery as a caustic, extends back to the most ancient times, but the particular modes of its application belong more properly to the professed treatises on surgery than to a work of this kind.

Caustic Potash.

The *causticum commune acerrimum* is the pure potash; when mixed with lime, its activity is lessened, and it is then called *Causticum commune mitius*. For the best mode of applying this substance, see the remarks on Issues. As this caustic spreads rapidly and eats deeply, care should be taken not to leave it on too long, when in the neighborhood of large blood-vessels. When, however, we wish to arrest its effects, they may be neutralized at once by vinegar, or vinegar and water.

Common Caustic with Opium.

R. Potassæ cum Calce, ʒij.

Pulveris Opii, ʒss.

Saponis Mollis, q. s.

Misce.—This preparation is employed for the same purpose as the common caustic.

Vienna Paste.

R. Potassæ, ʒv.

Calcis, ʒvj.

Alcoholis, q. s.

Ut fiat magma.

The paste is spread on the part to be cauterized, and is allowed to remain for ten or fifteen minutes, while the surrounding skin is protected by adhesive plaster.

Lunar Caustic.

This caustic is the *Argenti Nitras* deprived of its water of crystallization, fused and cast into moulds. Its mode of application is universally known. It is free from the inconvenience of the former.

Burnt Alum.

The *Alumen Ustum* is made by depriving this salt of its water of crystallization by heat. It is applied in powder to the fungous granulation of ulcers; and, mixed with an equal part of powdered sugar, is used to remove specks or nebulae of the cornea.

Chloride of Zinc.

This is a powerful escharotic, and is safe and certain in cases that require the destruction of a considerable thickness of the living tissues; "never causing violent pain in the affected parts, or any kind of irritative fever, or disorder of the digestive organs."

"The chloride of zinc may be applied as a caustic by means of a moistened hair pencil, either alone or mixed with an equal portion of oxide of zinc, or sulphate of lime, or according to the following form:—

Canquoin's Caustic Paste.

R. Zinci Chloridi, partem, j.

Farinæ, partem, jss.

Antimonii Terchloridi, partem, ss.

Aquæ, q. s.

Ut fiat pasta." DUNGLISON'S *New Remedies*, 6th ed. p. 694.

Caustic Paste of Chloride of Zinc.

R. Zinci Chloridi,

Calceis Sulphatis, āā ʒss.

Misce.—This preparation is very liable to deliquescence, to prevent which a little flour may be added at the time of applying it.

Mineral Acids.

The mineral acids are all escharotic, though seldom employed, on account of the difficulty of managing fluids when used in this way. *Nitric acid*, much diluted, is sometimes applied to foul ulcers in warm weather, to destroy the animalculæ which are apt to be generated in such sores.

Acid Nitrate of Mercury.

R. Hydrargyri Pernitratis Liquoris, *Dub.* f ʒss.

This is a powerful caustic, and is much employed in England and on the Continent to destroy malignant ulcerations, particularly those of a cancerous character. It is applied by means of a camel-hair pencil, and the parts are then covered with lint.

White Oxide of Arsenic, or Arsenious Acid.

R. Acidi Arseniosi, ʒj.

Aquæ, f ʒij.

Fiat solutio.

This solution is occasionally applied to cancerous sores, and was formerly used in a powder or paste as an empirical remedy. But it is now seldom employed, from the circumstance of its producing constitutional irritation. Sir Astley Cooper mentions cases in which, from its absorption when applied to open sores, partial if not complete paralysis of the limbs was the consequence.

Ointment of Arsenic.

R. Acidi Arseniosi, ʒj.

Adipis,

Cerati Cetacei, āā ʒvj.

Melt the cerate and lard over a slow fire, and then pour it on the white oxide of arsenic, and triturate it in a glass mortar until it is *intimately* united. *Pharm. Chirurg.*—For cancers, &c.

In hot weather, arsenic may be advantageously rubbed up with basilicon, to prevent its spreading on the sound parts. A scruple of the one to half an ounce of the other, makes a strong ointment.

The same care is required in its use as in using the solution.

Dubois's Arsenical Powder.

R. Acidi Arseniosi, ʒss.

Hydrargyri Sulphureti Rubri, ʒj.

Sanguinis Draconis, ʒss.

Fiat pulvis.—Chiefly used in the French hospitals as a caustic to cancerous affections. RATIER.

Arsenic is sometimes applied in the form of a paste made with water, which is carefully laid over the part to be removed, and renewed according to circumstances. When the slough forms, it should be dressed with a poultice, or with cerate spread on lint. It acts rapidly and requires watching, a remark that applies equally to all these pastes.

Solution of Muriate of Ammonia, &c.

R. Ammoniae Muriatis, ʒj.

Aceti destillati, f ʒij.

Aquæ, f ʒiv.

Fiat solutio.—To be applied to venereal warts.

BELL on Venereal.

Chromic Acid.

This acid has lately been employed in Germany as an escharotic. It is said to be efficient, manageable, and less painful

than the ordinary applications. A concentrated solution is applied by means of a glass rod. The solid chromic acid has to be used with care, on account of its penetrating action.

Moxa.

This is another method of using the actual cautery; and, as it is engaging the attention of physicians, we insert the usual mode of applying it. The Chinese employ this remedy by means of the mugwort. They separate this soft, lanuginous substance by beating, and form it into a cone; the base of this is attached, by being wetted, to the skin, and the point set on fire.

Dr. Boyle recommends dipping cotton in a strong solution of nitrate of potash, and inclosing pieces of it thoroughly dried in paper cylinders, about one inch in diameter, and another inch in height. Dr. Burne employs wood in a state of dry rot, touchwood. This only requires to be shaped properly, and burns, he says, with certainty and steadiness, and is preferable to the other modes suggested; some of which, as common cotton or lint, requiring the use of the bellows, and, when they have been dipped in nitre, demanding care to prevent the scintillations from injuring the surrounding parts.

The cylinders or cones of touchwood may be confined to the skin by a little adhesive plaster fixed to the edges.

It has been proposed to place pieces of cold iron in the neighborhood of the part while the moxa is burning, in order to render the sensations more vague, and therefore less painful.

Moxa is used in diseases of the spine, paralysis, &c. &c.

SUB-CLASS.—ISSUES AND SETONS.

These remedies are employed as substitutes for blisters, especially when a permanent discharge is desired.

There are three kinds of issues in common use, by either of which we may procure a discharge of purulent matter for any length of time that may be requisite. They are applied in different situations, according to circumstances, and some one of the following parts is generally chosen: the nape of the neck; the hollow of the deltoid muscle, and between the shoulders or over the ribs in front. In the inferior extremities they are most conveniently formed in the inside of the leg, either above or below the knee.

1. *Blister Issue.*

To form this issue, it is merely necessary to apply a blister to any part in the usual manner; the skin being thus removed, the discharge must be promoted by means of stimulating ointments, as unguentum cantharidis, ceratum resinæ, ung. sabinæ, etc. These may be alternated with mild liniments, in order to prevent the ulceration and consequent discharge from becoming excessive.

2. *The Pea Issue.*

There are two modes of forming this issue; the first is made by means of a lancet; the part is merely taken in the thumb and finger, and an incision made by passing the lancet through the skin. The wound should be kept open by means of orange peas, the common pea, a bean, or a bit of orris root.

The second is formed with caustic potash.

The following extracts from the *Surgeon's Vade Mecum* will exemplify the manner of forming the issue with caustic:—

“The part is first to be covered with a piece of adhesive plaster, in the centre of which a circular hole has been cut of the size of the intended issue; in this hole the caustic *kali purum*, made into a paste with soap, is to be placed. The whole is then to be covered with another plaster, secured by a bandage, and suffered to remain for *ten* or *twelve* hours. Generally, in two or three days an eschar begins to separate, when the opening should be filled with the substance made choice of.” These substances are enumerated above.

3. *The Seton.*

To form this artificial discharge, take the seton needle, to which is attached a skein of silk, the latter being partially anointed with some oleaginous substance. The needle is to be passed completely through the part chosen for the operation, after which it is removed, and the ends of the silk left hanging from the wound. It should be dressed every day, or twice a day, with some mild ointment; and if this be not sufficient to keep up the discharge, the more stimulating unguents may be used, as mentioned when speaking of the blister issue.

CLASS XVIII.

EPISPASTICS.

“By the ancients, all the external applications which redden or inflame the skin were called *Epispastics*, and these were designated according to the several degrees of effect: the slightest, entitled *phœnigmoi*; the next, *sinapismi*; the more active, *vesicatorii*; and the strongest, *caustici*.”

At the present day, the term is restricted to those articles which produce a serous discharge beneath the cuticle. The blister of common language.

As a general rule, blisters should remain on the patient six or eight hours, in order to insure their full effect. There are individuals, however, whose skins are so delicate, that a shorter application will answer every purpose; in children, it is seldom necessary to leave them on longer than three or four hours. When they are applied to the scalp, twelve hours are almost universally required; and Dr. Chapman directs the hair to be shaved, if possible, some hours before the application is made.

The usual dressing, when a blister is taken off, is basilicon ointment, or simple cerate, according to the urgency of the case, or irritability of the system. In order to keep it open, the weak epispastic, or savin ointment, will be found sufficient.

In acute disease, these remedies ought never to precede those means which have a tendency to reduce inflammatory action. It is generally admitted that, otherwise employed, they do harm: although there are individuals of acknowledged experience who say “they can bring the patient to the blistering point at pleasure,” and in pleurisy, and some other acute diseases, apply them soon after a free bleeding. Blisters should be placed as near the affected part as possible. In fevers, they are usually put on the inside of the forearms or legs, back of the neck, or between the shoulders.

When the system is very prostrate, they should be applied

to the inside of the thighs, or the last two places mentioned; for, when the circulation is languid in the extremities, they seldom draw well, and are even liable to produce gangrene by exhausting the little vitality that remains.

When, from peculiarity of constitution, *strangury* is produced, we must take them off after three or four hours, and bathe the part with sweet oil, or apply a poultice, giving at the same time diluent drinks, such as flaxseed tea or barley water, with the sweet spirits of nitre; camphor or parsley tea; bladders also, half filled with warm water, should be applied to the pubes. An opium suppository or injection will be found very advantageous. A poultice of ground flaxseed or starch is often serviceable, and a warm bath affords a prompt relief. Prevention, however, is better than cure; and in irritable constitutions, or in persons disposed to strangury, the blistering ointment, before being spread, should be rubbed up with three or four grains of opium, and three or four times as much camphor, before it is spread. This preventive seldom fails. Blisters are rendered much less irritating by covering them with a fine gauze, which, if it be pressed in close contact with the plaster, will not prevent or materially retard its operation. It is seldom necessary to keep a blister on longer than to *redde[n] the skin*, when a simple poultice will complete the vesication. In children, this precaution is imperative.

Blistering Cerate.

R. Cerati Cantharidis, q. s.

Ut fiat emplastrum epispasticum, 6 × 6.

Epispastic Ointment.

R. Ceræ flavæ,

Resinæ,

Adipis, āā ʒvj.

Melt over a slow fire, and add slowly—

Pulveris Cantharidis ʒvijss.

Misce, et fiat unguentum.

In employing this ointment, or the preceding cerate, the margin of the blister should be covered with adhesive plaster, in order to confine it to its place, and prevent its creating unnecessary irritation. Another mode is to draw straps across it diagonally, which plan we generally prefer; it prevents the blister from rising in the middle, and thereby partially defeating its object.

Instantaneous Vesication.

The French practitioners have a process for raising an instant blister. A piece of linen or paper, being cut to the requisite size, is immersed in spirits of wine or brandy, or even Cologne water, oil of turpentine, or spirits of camphor, whichever is most readily obtained. It is then laid on the part to be blistered, taking care that the moisture does not wet the surrounding surface. The flame of a lighted taper is then applied quickly over the surface of the paper so as to cause ignition, which is almost instantaneous. At the conclusion of this operation, the cuticle is found detached and raised from the true skin beneath.

Cantharidin.

This substance, more or less impure, has been used during the last few years, as a substitute for the common blistering cerate. The pure cantharidin is too expensive for general use, but an impure article, sufficiently concentrated for ordinary purposes, can be readily obtained. The cantharidin blistering tissues, and vesicating taffetas, now offered for sale, are of this character:—

Ether Cantharidalis, or Green Oil of Flies.

R. Pulveris Cantharidis, partem unam.

Ætheris, partes duas.

Digere per dies tres et exprime.

If this extract be applied twice without any admixture, by means of a hair pencil, it produces, in children, free vesication in an hour or two; and if applied three times in an adult, the same effect takes place in three or four hours. The ether evaporates, and the remaining green fixed oil operates like the cantharides in substance.

OETTINGER.

Collodium Vesicans.

R. Ætheris Cantharidalis (prepared as above),

Collodii, partes æquales.

Misce.

It is computed that two scruples of this collodium are equal to four drams of common blistering plaster. It is applied by a brush two or three times over the affected part.

OETTINGER.

Extract of Cantharides.

During the past winter (1853), a simple aqueous extract of Spanish Flies, prepared by Professor Procter, was used extensively in the Pennsylvania Hospital, and found to be a most efficient substitute for the blistering cerate. The application of it for six hours raised as free a blister as the cerate did in eight.

Since the discovery, by the same gentleman, that cantharidin is soluble in oil, oleaginous extracts have been devised, thus proving the general correctness of Galen's plan of preparing an ointment by macerating the entire insect in melted lard for twenty-four hours, and then straining by expression. An oleaginous extract, prepared by digesting the powdered flies in melted lard, is a most efficient substitute for the officinal cerate.

CLASS XIX.

RUBEFACIENTS.

THE substances called Rubefacients not unfrequently vesicate on very delicate skins. The term, therefore, is relative. They are employed for the same purpose as blisters, and are often good substitutes for them.

Mustard Plaster.

R. Pulveris Sinapis, ℥ij.

Aquæ tepidæ, q. s.

Ut fiat cataplasma.

It is a very general practice to use vinegar in making this plaster, but if the mustard be pure, the chemical reactions, on which the irritant properties chiefly depend, are better developed with water than vinegar. In fact, a high heat, or the use of a strong acid solution materially checks the process.

The uses of the mustard poultice are too well known to require description. It is frequently made by adding an equal quantity of wheat or rye flour; this should always be done for children, or other persons of delicate skins. It cannot generally be borne longer than a half hour or hour.

Garlic Cataplasma.

The garlic cataplasma is employed with the same intention as the mustard, though much less active. It may be made by bruising the cloves of garlic, and mixing them in a common poultice of bread and milk. Onions may be substituted for garlic.

Cataplasma of Mustard, Pepper, &c.

R. Seminum Sinapis contus. ℔ss.

Pulveris Piperis Nigri,

———— Zingiberis, āā ℥j.

Aquæ bullientis, q. s. ut fiat cataplasma.

To be applied to the region of the stomach in severe pain or sickness; or to the soles of the feet when revulsion is required.

Spice Plaster.

R. Pulveris Caryophylli,
 ——— Cinnamomi,
 ——— Capsici, āā ʒss.

Misce, et adde—

Farinæ,

Spiritus Vini Gallici, āā q. s. ut fiat cataplasma.

Let the plaster, thus prepared, be applied hot to the epigastric region.

Clove Plaster.

Quilt some powdered cloves in flannel, and wring them out in hot brandy. Thus prepared, they form an excellent external application in colic, and in irritability of the stomach, &c.

Warm Strengthening Plaster.

R. Picis Burgundicæ, ʒv.
 Unguenti Cantharidis, ʒj.

Fiat emplastrum secundum artem.—This plaster is very irritating, and sometimes draws a blister.

Mercurial Ointment, Camphor, &c.

R. Unguenti Hydrargyri, ʒj.
 Olei Terebinthinæ,
 Camphoræ, āā ʒij.
 Cerati Simplicis, ʒj.

Fiat unguentum secundum artem. Used as a rubefacient over the hypochondriac region, when blisters are precluded.

Tartar-emetie Plaster.

This may be made by sprinkling the powdered Tartrate of Antimony and Potash on the surface of a Burgundy pitch or adhesive plaster, or on one made by spreading basilicon, or simple ointment, on leather or linen. In either way the margin should be left clean.

Tartar emetic requires a longer time to produce its effect than the ordinary blister; the irritation is of a peculiar and very painful kind, and its effects must be watched with extreme care.

Tartar-emetie Ointment.

R. Antimonii et Potassæ Tartratis, ʒj.
 Adipis, ʒj.

Misce secundum artem.

This ointment, when applied to the skin, is a slow but certain means of exciting an eruption. We have given the formula of Dr. Hannay, of Glasgow, in Scotland. The officinal formula is two drams to the ounce, a better strength for ordinary purposes.

Tartar-emetic and Corrosive Sublimate.

R. Antimonii et Potassæ Tartratis, ʒj.

Hydrargyri Chloridi Corrosivi, gr. v.

Aquæ, f ʒj.

Spiritûs Lavandulæ Compositi, f ʒj.

Solve sales in aqua, dein adde spiritum.

This solution is applied by wetting the fingers with it, and rubbing it on the part. The rubbing should be continued for five or ten minutes, and after some hours a pustular eruption will appear.—Dr. Hannay's essay, *De Tartratis Antimonii usu externo*, p. 26.

It has been used in pertussis, dyspnœa, catarrh, &c., and in the removal of incipient tumors.

It has been ascertained by the illustrious Jenner and others, that tartar emetic, employed for an undue length of time externally, will produce violent effects on the system.

We find in a note in Beck's *Medical Jurisprudence*, vol. ii. p. 283, that Dr. Francis mentions two cases wherein tartar-emetic ointment, excessively used, produced effects similar to those at times arising from the outward application of arsenic. Magendie and Brodie ascertained by experiments upon animals, that inserted into wounds its effects resemble those produced by internal exhibition.

Granville's Antidynous Lotions.

These preparations act with great celerity, and are now in common use. They are formed of *three ingredients*, mixed in certain proportions, according to the urgency of the case.

Viz. 1. Of a saturated solution of ammonia; 2. Of the spirits of rosemary; and 3. Of spirits of camphor.

1. *The milder Ammoniated Lotion.*

R. Liquoris Ammoniæ fortioris, f ʒj.

Spiritûs Rosmarini, f ʒvj.

Tincturæ Camphoræ, f ʒij.

Misce.

The stronger Ammoniated Lotion.

R. Liquoris Ammoniae fortioris, f 3x.

Spiritus Rosmarini, f 3iv.

Tincturae Camphorae, f 3ij.

Misce.

The only point in which these formulæ differ from the original ones by Dr. Granville is in the preparation of the spirit of rosemary. Our pharmacopœia directs the oil of rosemary to be dissolved in alcohol. "Dr. Granville had it prepared by taking two lbs. of the tips or small leaves of fresh rosemary, and eight pints of alcohol, which are infused together for twenty-four hours in a covered vessel, and after adding as much water as will first prevent the empyreumatic smell, distil over *seven* pints."

As to the method of applying these counter-irritants, we shall use the words of their inventor, Dr. Granville. "First, impregnate with the colorless and transparent liquid, either a piece of linen folded six or seven times to the size of the part intended to be covered, or a piece of thick and coarse flannel; and, secondly, lay either of them on the spot, pressing with the hand at the same time, very steadily and firmly, the said linen or flannel, over which there should be placed a thick towel, doubled several times, so that not only the evaporation of the lotion may be impeded, but the hand employed in pressing the application to the part may not suffer damage from any evaporation, or from contact with the liquid. In some parts of the body more convenient than the others, the readiest and most effectual mode of pressing down the application is, by tying over it a towel or thick bandage; but to this mode there is the objection that we cannot, under certain circumstances, infect the part as quickly and as often as required, so as to judge, from the effect of the application, when to stop, or how long to persevere in using it."²

When applied to the temples, forehead, jaws, or chin, great care must be taken to prevent the effluvia from affecting the eyes and nostrils. The skin should be examined in a minute or two after the application, in order to see the effect produced. When the pain is severe after the lotion is removed, simple diluent washes, and warm water, which is perhaps the best, will speedily allay it. Vesication will generally take place in from two to five minutes, and even in a minute in some persons; but where cauterization is requisite, ten or twelve minutes may be necessary. The irritated or denuded surface may be dressed

² Granville. Counter-Irritation, its Principles and Practice, p. 39.

according to existing indications, with a simple poultice, tepid fomentations, cerate, or irritating ointments.

Gondret's Pommade Ammoniacale.

R. Adipis, partes xxxij.

Olei Amygdalæ, partes ij.

Liquoris Ammoniaë (*sp. gr.* 0.905), partes xvij.

Melt the lard, mix the oil with it, and pour them into a wide-mouthed phial with a glass stopper; then add the ammonia, close the bottle, mix the contents together by shaking, and keep the mixture in a cool place. If the skin is to be excited, perspiration to be re-established, or some subcutaneous engorgement to be dissipated, light and hasty practices accomplish these objects. If a rubefacient effect is desirable, spread the ointment thickly on linen, and apply it for one or two minutes. For vesication, continue the application for five or six minutes; and, by a farther continuance, absolute cauterization is speedily produced.

These powerful preparations are used in a great variety of diseases, and in all cases when speedy and powerful counter-irritation is demanded; as in neuralgia and rheumatic affections generally, both acute and chronic; spasms, convulsions, gout, paralysis, inflammatory diseases of internal organs, diseases of the heart, sprains, &c. &c.

Solution of Caustic Potash.

R. Potassæ, ʒiss.

Aquæ, fʒij.

Fiat solutio.—Applied to the spine in tetanus.

Dr. Joseph Hartshorne speaks in high terms of the beneficial effect of this solution in the above disease. The best mode of applying it, so as merely to produce a rubefacient action, is to tie a piece of sponge to a stick or common table fork, and apply it from the nape of the neck to the sacrum. When the solution is strong, the effect is powerful.

Tincture of Cayenne Pepper.

R. Pulveris Capsici, ʒij.

Alcoholis Diluti, fʒviij.

Misce.—This preparation forms an excellent rubefacient in low fever. Dr. Chapman recommends it strongly in sciatica and other rheumatic affections.

Turpentine Liniment.

R. Olei Terebinthinæ, f 3ij.
 — Olivæ, f 3ij.
 Tincturæ Camphoræ, f 3j.
 Liquoris Ammoniaë, f 3j.
 Fiat linimentum.

Decoction of Turpentine and Cantharides.

R. Pulveris Cantharidis, 3j.
 Olei Terebinthinæ, f 3viij.
 Digest together for three hours, then filter.

This is the process usually followed by our apothecaries; but we conceive thirty minutes' boiling would be quite sufficient, and the cantharides be less likely to be injured by the heat. It was first made by Dr. Hartshorne of this city, who has used it with great success in the typhoid state of fever. It is merely rubbed over the surface of the body. S. G. MORTON.

Liniment of Cantharides, &c.

R. Camphoræ, 3iij.
 Solve in Decocti Cantharidis cum Terebinthinâ f 3ss.—et adde—
 Unguenti Hydrargyri,
 ——— Simplicis, āā 3j.
 Fiat linimentum.—This preparation is employed in low fever, when it is desirable to rouse the system, and at the same time affect it with mercury; also in tumors.

Liniment of Croton Oil.

R. Olei Tiglii,
 — Olivæ, āā f 3ss.
 Fiat linimentum.—Croton oil is one of our safest and most efficient rubefacients. Rubbed for a few minutes, or sometimes even for a few seconds, on the skin, it produces irritation and redness, followed in some hours by a pustular eruption. If the effect is too severe at first, it is readily counteracted by a poultice, preceded by washing with mild soap and water to remove the oil. In some persons the oil may be applied pure.

Volatile Liniment.

R. Olei Olivæ,
 Liquoris Ammoniaë, āā f 3j.
 Misc.—Applied to the neck in inflammation of the throat.

Soap Liniment, &c.

R. Linimenti Saponis Camphorati, f ʒij.
 Liquoris Ammoniae, f ʒj.
 Tincturæ Opii, f ʒss.
 Fiat linimentum.—To be used as above.

Iodine Paint.

R. Iodinii, gr. lxiv.
 Potassii Iodidi, gr. xxx.
 Alcoholis, f ʒj.

Fiat solutio.

KING'S COLLEGE HOSPITAL.

Iodine is one of our best rubefacients and counter-irritants. This and the following formula furnish solutions that prove satisfactory in practice.

Ethereal Solution of Iodine.

R. Ætheris, f ʒj.
 Potassii Iodidi, gr. xv.
 Iodinii, q. s. ad saturandum.

Misce.

"It is applied in the same way as the tincture, by means of a camel-hair pencil rubbed over the part, until it begins to produce a burning sensation in the part; then cover it with a pledget of wadding so as to prevent evaporation."

R. THOMSON.

Liniment with Croton Oil and Potash.

R. Olei Tiglii,
 Liquoris Potassæ, āā ℥xv.

Misce, et adde—

Aquæ Rosæ, f ʒj.

Fiat linimentum.

This may be rubbed on the affected part twice or three times a day, until minute pustules appear.

JAMES ALLEN.

Cowhage Ointment.

R. Mucunæ, gr. viij.
 Adipis, ʒj.

Fiat unguentum.

The ointment must be rubbed in from ten to twenty minutes. The immediate effect is a sensation resembling stinging with nettles, but this passes off in less than half an hour. The skin generally becomes covered with white flat papulæ which soon disappear.

It is used in the same cases as tartar emetic or the Croton oil liniment.

BLATIN.

CLASS XX.

ANOMALOUS EXTERNAL APPLICATIONS.

It was the original intention of the Author of this work to arrange all the prescriptions, whether for internal or external use, under their appropriate therapeutic heads, and a few of the earlier editions conformed, more or less, to that view. But the introduction of a great number of new remedies, the true value of which can only be determined by careful trials, has caused much difficulty in arranging them properly in that respect. This remark is particularly applicable to the host of articles used, under various forms, in the skin diseases. To remedy this, and to facilitate reference as much as possible, the Editor of the tenth edition has added three new classes, of which the present constitutes one. Class XX. is arranged on a pharmaceutical, rather than a therapeutical basis, and its subdivisions are named according to the form of preparation employed. Viz. Collyria, Injections, Gargles, Liniments, Ointments, Cataplasms, Baths, Fomentations, and Fumigations.

A.—COLLYRIA.

These are preparations applied to the eyes. They are sometimes dry, but generally liquid—consisting of infusions, decoctions, or distilled waters, with the addition of various substances.

POWDERS.

Dry Collyrium of Oxide of Zinc and Sugar.

R. Sacchari,
Zinci Oxidi, āā partes æquales.
Tere in pulverem.

M. RECAMIER.

Dry Collyrium of Red Precipitate, &c.

R. Sacchari, ʒij.
 Hydrargyri Oxidi Rubri, gr. x.
 Zinci Oxidi, ʒj.
 Fiat pulvis subtilissimus.

DUPUYTREN.

This and the preceding powder are used in opacities of the cornea, by blowing them into the eye through a quill.

Dry Collyrium of Opium, &c.

R. Pulveris Opii, gr. iv.
 Hydrargyri Chloridi Mitis,
 Sacchari, āā ʒj.
 Tere bene.

These compounds are blown into the eyes for the removal of specks on the cornea resulting from inflammation.

Powder of Blue Vitriol, &c.

R. Cupri Sulphatis,
 Boli Armeniæ, āā ʒj.
 Camphoræ, ʒij.

Fiat pulvis.—Put ʒj. of this powder into a pint of boiling water; filter the solution, and mix ʒj. of it with ʒj. of clear water, and apply it to the purulent eyes of infants by means of a syringe,
pro re nata.

COOPER'S Surgery.

Lapis Divinus.

R. Aluminis,
 Potassæ Nitratis, } āā ʒj.
 Cupri Sulphatis, }
 Camphoræ, ʒss.

The alum, nitre, and sulphate of copper are to be fused together in a glazed earthen crucible, to drive off the water of crystallization, and the camphor is to be added near the end of the process. The cold mass is to be broken in pieces, and preserved in a closely stoppered bottle. This article is much employed on the Continent as an eyewash, in the strength of two grains and upwards to a fluidounce of distilled water.

SOLUTIONS AND MIXTURES.

Alum Collyrium.

R. Pulveris Aluminis, gr. xij. vel xx.
 Aquæ Rosæ, fʒiv.

Fiat solutio.—To be applied to the eye in chronic inflammation.

Collyrium of Iodide of Potassium.

- R. Aquæ Rosæ, f 3vj.
 Potassii Iodidi, gr. xxiv.
 Iodinii, gr. j. vel ij.

Fiat collyrium. To be applied to the eyes three or four times in the day. In scrofulous ophthalmia, and ulcerations of the cornea and conjunctiva. RYAN.

Collyrium of Green Vitriol.

- R. Ferri Sulphatis, gr. ij.
 Aquæ, f 5j.

Fiat solutio.—Used in chronic ophthalmia.

Collyrium of Subacetate of Lead, &c.

- R. Liquoris Plumbi Subacetatis, gtt. xij.
 Vini Opii, gtt. xl.
 Aquæ Rosæ, f 3iv.

Fiat collyrium.—This should not be used when there are ulcers on the cornea.

Collyrium of Vinegar, &c.

- R. Aceti destillati, f 3j.
 Alcoholis, f 3ss.
 Aquæ Rosæ, f 3viij.

Fiat mistura. *Signa*.—To be applied to weak eyes, and to ophthalmia after depletion.

Collyrium of Acetate of Zinc.

- R. Zinci Acetatis, gr. iv.
 Aquæ Rosæ, f 3j.

Fiat solutio.—Used like the preceding.

Opiate Collyrium.

- R. Pulveris Opii, gr. j.
 Camphoræ, gr. v.
 Mucilaginis Acaciæ, f 3j.
 Misce, et fiat collyrium.

Alum Curd.

- R. Pulveris Aluminis, 3ss.
 Albumen unius Ovi.

Agitate them well until a coagulum be formed.—To be applied on a linen rag to inflamed eyes. It should be removed in three or four hours.

Collyrium of Nitrate of Silver.

R. Argenti Nitratis, gr. j. vel ij. ad gr. xx.

Aquæ destillatæ, fʒij.

Fiat solutio.—In acute ophthalmia, and to promote the absorption of nebulae of the cornea.

Collyrium of Sulphate of Cadmium, &c.

R. Cadmii Sulphatis, gr. ij.

Aquæ Rosæ, fʒij.

Vini Opii, fʒj.

Fiat mistura.—Used for opacities of the cornea, and in chronic ophthalmia.

The quantity of sulphate of cadmium is gradually increased to four grains in an ounce.

FRONMÜLLER.

Collyrium of Sulphate of Copper.

R. Cupri Sulphatis, gr. vj.

Camphoræ, ʒj.

Aquæ ferventis, fʒviiij.

Rub the camphor with the water, then strain, and add the sulphate of copper.

This is nearly the same as the formula recommended by Mr. Ware in the treatment of purulent ophthalmia.

Strychnia Collyrium.

R. Strychniæ, gr. ij. vel iv.

Acidi Acetici diluti,

Aquæ destillatæ, āā fʒj.

Fiat solutio.—A drop or two, diluted or not according to circumstances, to be applied to the eye several times a day.

DR. RYAN on Amaurosis.

Collyrium of Corrosive Sublimate.

R. Hydrargyri Chloridi Corrosivi, gr. ij.

Aquæ destillatæ, fʒviij.

Fiat solutio.

The above may be advantageously used in gonorrhœal ophthalmia, as also in that of a scrofulous character.

Collyrium of Opium and Camphor.

R. Extracti Opii, gr. x.

Camphoræ, gr. vj.

Aquæ ferventis, f̄3xij.

Rub the camphor and opium well together in a mortar, and add the water. Then strain before using.

Conradi's Collyrium.

R. Hydrargyri Chloridi Corrosivi, gr. j.

Decocti Cydonii, Lond. f̄3j.

Aquæ Rosæ, f̄3vj.

Vini Opii, f̄3j.

Fiat mistura.

This collyrium is praised by most of the German oculists in cases of chronic ophthalmia.

Bates's Camphorated Water.

R. Cupri Sulphatis,

Boli Armeniæ, āā gr. viij.

Camphoræ, gr. ij.

Aquæ bullientis, f̄3viij.

Fiat infusum et cola.

This is used as a lotion in the purulent ophthalmia of infants.

Collyrium of Acetate of Ammonia and Camphor.

R. Liquoris Ammonia Acetatis, f̄3ij.

Aquæ Camphoræ, f̄3vj.

Misce. *Signa.*—For eyes requiring a mild astringent and stimulant application.

Collyrium of Acetate of Ammonia with Opium.

R. Liquoris Ammonia Acetatis, f̄3ij.

Aquæ ferventis, f̄3vj.

Extracti Opii, gr. x.

Dissolve the extract of opium in the hot water, strain, and add the solution of acetate of ammonia. In acute ophthalmia, when considerable pain remains after depletion.

Emollient Collyrium.

R. Radicis Altheæ, 3ij.

Aquæ destillatæ, Oj.

Infuse for three hours near the fire, and strain.—In inflammation of the conjunctiva accompanied with violent irritation.

Collyrium of Poppies, &c.

R. Decocti Papaveris, *Lond.* f̄iv.

Aquæ Rosæ,

——— Camphoræ, āā f̄ij.

Misce.—Either of the last two collyria is adapted to the early stages of ophthalmia.

These collyria can be most successfully applied to the eyes by introducing the point of a small syringe into the outer angle, and gradually injecting them.

Anodyne Collyrium.

R. Colchici Radicis contusi, ʒj.

Infusi Lini bullientis, f̄iv.

Tincturæ Opii, f̄ij.

Fiat mistura.—For ophthalmia accompanied with great sensibility.

M. JADELLOT.

Collyrium with Conia.

R. Coniæ, partes ij.

Alcoholis, partes xij.

Aquæ destillatæ, partes cc.

Fiat solutio.—Employed in scrofulous ophthalmia, with intolerance of light, by rubbing it around the orbits, or dropping it in the eye several times a day.

Conia, the active principle of conium maculatum, is a most virulent poison, and therefore care is requisite, even in its external use.

FRONMÜLLER.

Guthrie's Ointment.

R. Argenti Nitratis, gr. ij. ad x.

Liquoris Plumbi Subacetatis, gtt. xv.

Unguenti Cetacei, *Lond.* ʒj.

Fiat unguentum.

“The manner of using the ointment, is by introducing between the lids a portion, larger or smaller, as the case may seem to require it, from the size of a large pin's head to that of a garden pea.”

GUTHRIE.

Ointment of Red Precipitate, &c.

R. Hydrargyri Oxidi Rubri, gr. x.

Zinci Sulphatis, ʒj.

Adipis, ʒij.

Fiat unguentum.

Recommended by Professor Dupuytren, in obstinate chronic ophthalmia depending on a scrofulous diathesis.

B.—INJECTIONS.

Injections are liquid compositions of emollient, astringent, or stimulating substances. The parts into which they are thrown are the external auditory passages, the canal of the urethra, the bladder, and also fistulous cavities, the tunica vaginalis in the case of hydrocele, and the vagina.

A.—FOR THE EAR.

Injection with Camphor, &c.

R. Camphoræ, ʒiv.

Olei Cajuputi, fʒvij.

— Amygdalæ, fʒxv.

Tere simul.—A few drops to be dropped in the ear, in cases of dulness of hearing, from hardened wax, or dryness of the membrane.

RADIUS.

Goulard's Extract, and Rose Water.

R. Liquoris Plumbi Subacetatis, fʒj.

Aquæ Rosæ, fʒj.

Misce.—This is to be dropped into the ear until the meatus is full, and be allowed to remain there for a few minutes.

WILDE.

Mixture with Bates's Alum Water.

R. Liquoris Aluminis Compositi, Lond. fʒj.

Aquæ Rosæ, fʒj.

Misce.

Used in otorrhœa, in the same way as the preceding.

WILDE.

Mixture with Aqua Sappharina.

R. Liquoris Cupri Ammonio-Sulphatis, Lond. fʒj.

Aquæ destillatæ, fʒj.

Miscé.

Like the preceding, this is used as an astringent in otorrhœa.

WILDE.

Solution of Nitrate of Silver.

R. Argenti Nitratis, gr. x.

Aquæ destillatæ, f 3j.

Fiat solutio.

Used in simple otorrhœa by painting the meatus and membrana tympani with a fine camel-hair pencil, dipped in the solution, every second or third day; and one of the three foregoing solutions is used in the interim, night and morning.

WILDE.

Injection of Pyroligneous Acid.

R. Acidi Pyrolignei, *Dub.* f 3ij.

Aquæ destillatæ f 3vj.

Fiat injectio.—To be injected twice a day with a silver syringe, in purulent discharge from the meatus auditorius.

BUCHANAN *on the Ear.*

Injection of Catechu.

R. Catechu, gr. xij.

Aquæ bullientis, f 3vj.

Fiat injectio.—To be used four or five times a day. *Idem.*

Pyroligneous Acid, Ether, &c.

R. Acidi Pyrolignei, *Dub.*

Spiritus Ætheris Sulphurici, *Ed.*

Olei Terebinthinæ, aa partes equales.

Misce. *Signa.*—Introduce two drops into the meatus auditorius every night, when the cerumen is deficient in quantity.—*Idem.*

Oxgall and Peruvian Balsam, for the Ear.

R. Fellis Bovini, f 3iij.

Balsami Peruviani, f 3j.

Misce.—To be dropped occasionally into the ear to correct a fetid discharge, syringing the ear every day with a weak solution of soap and water.

DR. HUGH SMITH.

Injection of Creasote.

R. Creasoti, gtt. ij. vel iv.

Syrupi, f 3j.

Aquæ, f 3vij.

Misce.

To be used like the foregoing.

EDITOR.

Solution of Nitrate of Silver.

R. Argenti Nitratis, ʒss.—j.

Aquæ destillatæ, f ʒj.

Misce.

We have given above a weaker formula for the application of lunar caustic to the membrani tympani. The present solution is intended to be applied every third day to the *outer half* of the meatus, and adjacent portion of the concha, in chronic inflammation or thickening of the membrane lining the tympanic cavity.

J. TOYNBEE.

Solution of Sulphate of Cadmium.

R. Cadmii Sulphatis, gr. viij.

Aquæ, f ʒj.

Fiat solutio.

Used in otorrhœa.

LINCKE.

B.—FOR THE URETHRA.

When injections are employed in the cure of gonorrhœa, two circumstances merit especial attention, viz., the manner of injecting them and the frequency of using them. When the common syringe is used, the pipe should be perfectly smooth, short and conical, so that it cannot pass more than a quarter of an inch into the urethra. The patient should oil the pipe before using it, seat himself on the edge of a chair so as to avoid pressing the perineum, draw the penis forward on the syringe with the left hand, and throw up the injection to the seat of disease (previously ascertained) with the other. As respects the frequency of injecting, it may be observed, that five or six times a day are not too often, and that the cure will be retarded by using these remedies at longer intervals. We cannot forbear remarking that gonorrhœal inflammation, situated near the extremity of the penis, can be often cured by a *routine plan*; but when seated higher up the canal, and affecting Cowper's gland or the prostate, it becomes a serious disease, the cure of which ought not to be undertaken by persons ignorant of the structure of the parts and the nature of morbid action.

Nitrate of Silver.

This article is extensively used, both in gonorrhœa and gleet. Some practitioners employ a weak solution, of two or three grains, to the ounce of rose water, twice a day. Others prefer

ten grains to the ounce and employ the injection every second or third day. Another class prefer an ointment of ten or twenty grains to the ounce, and smear this over a bougie, which is then passed into the urethra. In acute gonorrhœa, either of the latter plans will frequently arrest the disease in a few days. In gleet, perhaps, the first plan is preferable.

Injection of Corrosive Sublimate, &c.

R. Zinci Sulphatis, gr. x.
Hydrargyri Chloridi Corrosivi, gr. ij.
Aquæ Rosæ, f ʒviij.
Fiat injectio.—In gleet.

This powerful remedy should be employed with great caution.

Injection of Sulphate of Zinc, &c.

R. Zinci Sulphatis, gr. x.
Pulveris Acaciæ, ʒij.
Tincturæ Opii, f ʒj.
Aquæ destillatæ, f ʒviij.
Fiat injectio.—In gonorrhœa virulenta.

Injection of Muriate of Ammonia, &c.

R. Ammoniæ Muriatis, gr. j. vel ij.
Hydrargyri Chloridi Corrosivi, gr. ij.
Aquæ destillatæ, f ʒiv.
Fiat injectio.—In chronic gonorrhœa.

Injection of Acetate of Zinc.

R. Zinci Acetatis, gr. viij.
Aquæ Rosæ, f ʒiv.
Fiat injectio.—In gonorrhœa virulenta.

For all these preparations, rose water is an admirable menstruum; it possesses the advantages of being slightly astringent and rendered pure by distillation, and is therefore particularly applicable to the formation of injections and collyria.

Injection of Chloride of Zinc.

R. Zinci Chloridi, gr. j. vel ij.
Aquæ, f ʒj.
Fiat solutio.
Used in gonorrhœa.

GODDARD.

Injection of Sulphate of Iron.

R. Ferri Sulphatis, gr. viij.

Aquæ, f 3j.

Fiat solutio.

Used in gleet.

Injection of Iodide of Iron.

R. Ferri Iodidi, ʒss. vel j.

Aquæ, f 3viij.

Fiat solutio.

Used in blennorrhœa.

RICORD.

Injection of Sulphate of Quinia.

R. Quiniæ Sulphatis, gr. ij. ad viij.

Aquæ, f 3j.

Acidi Sulphurici diluti, q. s.

Ut fiat solutio.

Used in gonorrhœa.

GODDARD.

Injection of Armenian Bole.

R. Zinci Sulphatis, ʒj.

Boli Armenia, ʒij.

Pulveris Acaciæ, ʒij.

Aquæ Rosæ, f 3viij.

Fiat injectio.—In gonorrhœa.

Injection of Calomel, &c.

R. Hydrargyri Chloridi Mitis, ʒj.

Acaciæ, ʒij.

Tincturæ Opii, f 3ss.

Aquæ, f 3viij.

Fiat injectio.—For gonorrhœa or gleet.

Injection of Sulphate of Copper.

R. Cupri Sulphatis, gr. vj.

Aquæ destillatæ, f 3vj.

Tincturæ Opii, f 3j.

Fiat solutio.—To be thrown up the urethra in chronic gonorrhœa.

Injection of Tea.

R. Foliorum Theæ Viridis, ʒj.

Aquæ ferventis, f 3vj.

Fiat infusum et cola.—For an injection in gonorrhœa virulenta.

Injection of Ammoniated Copper.

R. Cupri Ammoniatī, gr. v.

Aquæ Rosæ, f̄viiij.

Fiat solutio.—In chronic gonorrhœa.

Solution of Tannin.

R. Acidi Tannici, gr. xxxij.

Aquæ destillatæ, f̄viiij.

Solve.—Employed as a wash to chronic ulcers, and as an injection in obstinate blennorrhœa.—BERAL, in DUNGLISON'S *New Remedies*, 6th ed. p. 45.

Injection of Strychnia.

R. Strychniæ, gr. j.

Acidi Nitrici, gtt. ij.

Aquæ, f̄j.

Fiat solutio. *Signa*.—Inject a dram three times a day after urination. Used in gleet. JOHNSON.

C.—GARGLES AND MOUTH WASHES.

Gargles, like injections, are astringent and stimulating; and to be of any service must be frequently applied, and persevered in for some time.

Gargle of Brandy.

R. Spiritûs Vini Gallici,

Aquæ, aâ partes æquales.

Brandy and water form an excellent gargle for some cases of inflamed fauces.

Gargle of Galls and Alcohol.

R. Infusi Gallæ (ȝij. ad Oss.) f̄viiij.

Alcoholis, f̄j.

Fiat gargarysma.—In relaxation of the uvula and soft palate. Should be applied with a camel-hair brush.

Gargle of Infusion of Oak Bark.

R. Corticis Quercûs Albæ, ȝj.

Aquæ, Ojss.

Boil to a pint, and strain; then add—

Aluminis, ȝss.

Spiritûs Vini Gallici, f̄ij.

Misce.

Gargle of Sulphate of Quinia, &c.

R. Quiniæ Sulphatis, gr. xij.

Cupri Sulphatis, gr. xvj.

Acidi Sulphurici Aromatici, f3j.

Aquæ, f3viiij.

Fiat mistura. *Signa.*—To be used three or four times a day, in obstinate sore throat. JOS. HARTSHORNE.

Gargle of Borax, &c.

R. Sodæ Boratis, 3j.

Tincturæ Myrrhæ, f3ss.

Mellis despumati, f3j.

Aquæ Rosæ, f3iv.

Fiat gargarysma.—To be used as a gargle or mouth wash, in ptyalism or scorbutic affections of the gums.

Gargle of Sulphate of Zinc, &c.

R. Zinci Sulphatis, 3j.

Mellis, f3ss.

Tincturæ Myrrhæ, f3j.

Spiritûs Vini Gallici, f3j.

Aquæ Rosæ, f3vj.

Fiat gargarysma.—To be applied to ulcerated gums, occasioned by excessive salivation. Water can be added if it be too strong.

In mercurial sore mouth, common lead water forms a most soothing and useful application. It makes the teeth black—an effect, however, which is only temporary.

Dr. Blake's Toothache Solution.

R. Aluminis (in pulverem subtilissimum trituranum), 3ij.

Spiritûs Ætheris Nitrici, f3vij.

Fiat mistura.—Said by Dr. Blake, of London, to be an infallible cure for *toothache*, when not connected with rheumatism.

Gargle of Alum.

R. Aluminis, 3ij.

Aquæ, f3iv.

Solve.—This is Dr. Cavarra's gargle for *fætor oris*, or offensive breath.

Gargle of Pomegranate.

R. Granati Fructûs Corticis, ʒss.

Rosæ Gallicæ, ʒij.

Aquæ bullientis, f ʒvj.

Fiat infusum. Cola et adde—

Mellis despumati, f ʒj.

Aluminis, ʒij.

Fiat gargarysma.—For inflammation of the throat and fauces.

Muriatic Acid Gargle.

R. Infusi Cinchonæ Rubræ, f ʒiv.

Mellis despumati, f ʒj.

Acidi Muriatici, gtt. x. vel xvij.

Fiat gargarysma.—In chronic inflammation, when the pain and swelling have been reduced by depletion.

RATIER'S *Formulary*.

Detergent Gargle of Sulphuric Acid, &c.

R. Decocti Hordei, f ʒiv.

Mellis despumati, f ʒss.

Acidi Sulphurici, gtt. x. vel xx.

Fiat gargarysma.

RATIER.

Mercurial Gargle.

R. Hydrargyri Chloridi Corrosivi, gr. j.

Mellis despumati, f ʒss.

Aquæ destillatæ, f ʒiv.

Fiat solutio.—For venereal sore throat.

RATIER.

Acetous Gargle.

R. Acidi Acetici, f ʒij.

Ammoniæ Muriatis, ʒj.

Mellis, f ʒiss.

Aquæ, f ʒxij.

Fiat gargarysma.—For inflamed fauces.

RATIER.

Astringent Gargle of Acetate of Lead, &c.

R. Plumbi Acetatis, ʒss.

Syrupi, f ʒj.

Decocti Hordei, Oj.

Fiat solutio.—In mercurial sore mouth, &c.

RATIER.

Chloride of Lime Gargle.

R. Calcis Chlorinatæ, ℥ij.

Aquæ, Oj.

Solve et cola. Dein adde—

Mellis, f ℥j.

Misce.—Used in ulceration of the mouth and throat.

Disinfecting Mouth Wash.

R. Calcis Chlorinatæ, ℥iij.

Aquæ destillatæ, f ℥ij.

Tere simul et filtra. Dein adde—

Olei Rosæ, ℥v.

Alcoholis, f ℥ij.

Mellis Rosæ, f ℥ss.

Misce.—Used in mercurial sore mouth.

Cayenne Pepper Gargle.

R. Tincturæ Capsici, f ℥ss.

Aquæ Rosæ, f ℥viiij.

Misce.—A stimulating gargle used in scarlet fever.

GRIFFITHS.

Mel Ægyptiacum.

R. Linimenti Æruginis, Lond. f ℥j.

This is applied with a brush to venereal ulcers of the throat. Diluted with water (one dram to the ounce), it forms a detergent gargle.

Gargle with Iodine and Opium.

R. Tincturæ Iodini, f ℥j.—ij.

———— Opii, f ℥j.

Aquæ, f ℥vj.

Fiat gargarysma.

Signa.—To be well shaken when used.

This is employed three or four times daily in ulcerated sore throat.

Solution of Iodide of Zinc.

R. Zinci Iodidi, gr. x.—xxx.

Aquæ destillatæ, f ℥j.

Fiat solutio.

Applied daily to enlarged tonsils, by a piece of sponge tied to a quill.

J. J. ROSS.

Mixture of Sulphuric Acid and Honey.

R. Acidi Sulphurici, f ʒss.
 Mellis, f ʒj.
 Fiat linimentum.

Used for aphthæ, by touching the ulcerated surfaces frequently with a soft pencil.

Wash of Sulphate of Copper, &c.

R. Cupri Sulphatis, ʒij.
 Pulveris Cinchonæ, ʒss.
 Aquæ destillatæ, f ʒiv.

Misce. *Signa*.—In the gangrenous sore mouth of children, applied to the full extent of the ulceration and excoriation.

DR. B. H. COATES.

Sulphate of zinc is often employed in like doses, in this affection.

Gargle of Alum, Sage Tea, and Honey.

R. Foliorum Salviæ, ʒss.
 Aquæ bullientis, Oss.
 Fiat infusum, et cola, dein adde—
 Aluminis, ʒss.
 Mellis, f ʒj.
 Fiat mistura.

This domestic remedy is a very effectual one in cases of sore throat.

Linctus with Muriatic Acid, &c.

R. Acidi Muriatici, f ʒss.
 Mellis,
 Aquæ Rosæ, āā f ʒj.

Misce. *Signa*.—To be applied to scorbutic gums three or four times a day.

D.—LINIMENTS, LOTIONS, &c.

Lotion of Lime-water, Zinc, &c.

R. Zinci Oxidi, gr. xij.
 Cupri Sulphatis, gr. iij. vel iv.
 Mellis, f ʒj.
 Liquoris Calcis, f ʒij.

Fiat lotio.—To be applied to phagedenic ulcers of the genitals.

Vide SWEDIAUR on Syphilis.

Liniment of Lime-water and Linseed Oil (Carron Oil).

R. Olei Lini, f ʒiij.

Liquoris Calcis, f ʒvj.

Fiat linimentum.—To be applied to burns and scalds by means of a feather or soft rag.

Lotion of Lime-water and Alcohol.

R. Alcoholis, f ʒij.

Liquoris Calcis, Oss.

Fiat mistura.—In the same cases as the above.

Lotion of Goulard's Extract of Lead, &c.

R. Liquoris Plumbi Subacetatis, f ʒj.

Tincturæ Camphoræ, f ʒiij.

Aquæ, Oj.

Fiat lotio.—To be used as the former.

This compound is much recommended by some of the French surgeons. Applied by means of lint to old ulcers.

Lotion of Chloride of Zinc.

R. Zinci Chloridi, gr. ij.

Aquæ destillatæ, f ʒj.

Misce.—Applied on lint to chronic and indolent ulcers.

HANCKE.

Lotion of Sulphuret of Potassium, &c.

1. R. Potassii Sulphureti, ʒj. vel ʒij.

Aquæ, Oj.

Misce.

2. R. Acidi Muriatici, f ʒj. vel f ʒij.

Aquæ destillatæ, Oij.

Mix one ounce of each lotion with four ounces of warm water, and apply by means of a sponge. In psora.

M. ALIBERT.

Lotion of Subacetate of Lead.

R. Confectionis Rosæ, ʒj.

Mellis,

Liquoris Plumbi Subacetatis, āā f ʒss.

Tincturæ Opii, gtt. lx.

Misce. *Signa*.—Applied to painful venereal ulcers, and to chancres after the use of lunar caustic.

Compound Lotion of Sulphuret of Potassium.

- R. Potassii Sulphureti, ʒss.
 Saponis Veneti, ʒj.
 Alcoholis, f ʒiv.
 Tincturæ Myrrhæ, f ʒss.
 Liquoris Calcis, Oj.

Fiat mistura secundum artem.—To be applied to the head night and morning, in tinea capitis. The hair must be shaved previous to its employment.

Solution of Phosphoric Acid.

- R. Acidi Phosphorici, ʒj.
 Aquæ destillatæ, f ʒviij.
 Fiat solutio.

Dr. Lentin, a German physician, has employed this solution on compresses applied to ulcers situated over carious bones.—*SPILLAN'S Supplement to the British Pharmacopæias.*

Astringent Fomentation.

- R. Radicis Bistortæ,
 Granati Fructûs Corticis, āā ʒij.
 Ammoniæ Muriatis, ʒij.
 Vini Rubri, Oj.

Misce. *Signa*.—For indolent inflammatory swellings.

RATIER'S Formulary.

Lotions of Prussic Acid.

Prussic acid (Acidum hydrocyanicum dilutum) is used externally, in the proportion of ʒij. to Oj. of water, or other fluids. In cancerous ulcers; dartre; and injected into the vagina in cancer of the uterus.—See *MAGENDIE'S Formulary*, edit. 5th, p. 66.

Lotion of Corrosive Sublimate.

- R. Hydrargyri Chloridi Corrosivi, ʒj.
 Aquæ, f ʒvj.
 Solve et adde—
 Ammoniæ Muriatis, ʒij.
 Potassæ Nitratis, ʒss.

Fiat lotio.—Principally used as a wash in itch.

GOOD'S Study of Med. vol. iv.

Yellow Wash.

- R. Hydrargyri Chloridi Corrosivi, gr. j. vel. iij.
 Liquoris Calcis, f ʒj.
 Fiat solutio.

For ordinary purposes one grain to the ounce will be sufficient. In phagedenic and venereal ulcers.

Black Wash.

R. Hydrargyri Chloridi Mitis, ʒj.

Liquoris Calcis, fʒiv.

Misce. *Signa*.—The bottle to be well shaken before using.

This prescription is used in the same cases as the other. The red oxide in that, and the black oxide of mercury in this, are precipitated by the lime-water. They are sometimes administered in the form of injections in gleet and gonorrhœa.

Lotion of Cyanuret of Potassium.

R. Potassii Cyanureti, gr. xij.

Misturæ Amygdalæ, fʒvj.

Fiat lotio. In lichen, and other chronic eruptions.

Liniment of Chlorine.

R. Chlorinii Liquoris, *Dub.* fʒj.

Olei Olivæ, fʒj.

Fiat linimentum.—Applied externally to itch, herpes, and tinea capitis. DUNGLISON'S *New Remedies*, 6th ed. p. 195.

Iodine Liniment.

R. Linimenti Saponis Camphorati, fʒj.

Tincturæ Iodinii, fʒj.

Misce.

Creasote Liniment.

R. Creasoti, gtt. x.

Olei Olivæ, fʒj.

Fiat linimentum.—In herpes, &c.

Creasote Lotion.

R. Creasoti, gtt. x.

Aceti, fʒij.

Aquæ destillatæ, fʒij.

Fiat lotio.—Applied in phagedenic ulceration.

An admirable wash for herpetic eruptions may be prepared by mixing half an ounce of Wetherill's distilled creasote water with four times the quantity of pure water. This may be applied with safety to the skin of infants; but, if too active, it should be still farther diluted.

Camphorated Oil.

R. Camphoræ, ʒj.

Olei Olivæ, f ʒiv.

Tere simul, et fiat solutio; tunc adde—

Olei Terebinthinæ, f ʒjss.

To be applied by rubbing with a piece of flannel. In deep-seated rheumatic pains.

Epithem with Tincture of Cantharides.

R. Pulveris Cantharidis, ʒj.

Alcoholis, f ʒiv.

Digere cum leni calore per dies duas, cola et adde—

Camphoræ, ʒss.

Signa.—To be rubbed on the parts affected. In paralysis.

Epithem with Camphor and Carbonate of Ammonia.

R. Tincturæ Camphoræ, f ʒiv.

Ammonia Carbonatis, ʒss.

Olei Juniperi,

— Succini, āā f ʒij.

Misce.—To be employed in the same cases as the former.

Liniment with Camphor, &c.

R. Tincturæ Camphoræ, f ʒiij.

Acidi Acetici, f ʒj.

Fiat embrocatio.—To be applied to bruises, rheumatic parts, &c.

Liniment with Ammonia and Olive Oil; vulgo, Volatile Liniment.

R. Olei Olivæ,

Liquoris Ammonia, āā f ʒj.

Misce.—Applied to the neck in inflammation of the throat.

Liniment of Ammonia and Tartarized Antimony.

R. Linimenti Ammonia, f ʒj.

Antimonii et Potassæ Tartratis, ʒj.

Misce.—Applied to indolent tumors, and other swellings.

Liniment of Oil of Amber, &c.

R. Olei Succini, f ʒij.

— Olivæ, f ʒss.

Tincturæ Opii, f ʒij.

Spiritus Vini Gallici, f ʒiij.

Fiat linimentum. Signa.—To be rubbed between the shoulders.

In pertussis.

Liniment with Nux Vomica.

R. Tincturæ Nucis Vomicae, f 3j.

Liquoris Ammoniae, f 3ij.

Misce.

This is applied by friction in paralytic and nervous diseases.

MAGENDIE.

Anti-psoric Lotion of Dupuytren.

R. Potassii Sulphureti, 3iv.

Aquæ, Oj.

Acidi Sulphurici, f 3iv.

Misce.—To be applied two or three times a day on the parts covered with pustules, and with this treatment the simple bath is conjoined.

Lotion of Myrrh.

R. Tincturæ Myrrhæ,

Liquoris Calcis, aa f 3j.

Misce.

This lotion is recommended by Dr. Kirkland for removing fungous flesh.

Compound Opiate Liniment.

R. Tincturæ Opii,

Spiritus Ætheris Sulphurici, Ed.

Tincturæ Camphoræ, aa f 3ij.

Fiat embrocatio.—To be applied to rheumatic parts.

Also, in odontalgia, pain in the region of the ear, &c.

Opiate Liniment.

R. Olei Olivæ, f 3ij.

Tincturæ Opii, f 3ij.

Liquoris Plumbi Subacetatis, f 3ss.

Misce.

Embrocation of Acetate of Ammonia.

R. Ammoniae Carbonatis, 3ij.

Acidi Acetici, q. s. ad saturandum.

Alcoholis diluti, Ojss.

Mix the ammonia with the acid, and after effervescence ceases, add the spirit.

In rheumatic affections; or mixed with linseed meal, it may be used as a poultice to chronic inflammations of the joints, &c.

Compound Liniment of Petroleum.

R. Petrolei Barbadosensis, f 3j.

Camphoræ, 3ss.

Alcoholis, gtt. xl.

Fiat linimentum.—Powder the camphor by adding the spirit, and then mix the whole in a mortar.

Glycerin.

This principle, which is demulcent and antiseptic, forms a very useful addition to lotions and gargles. It has been used extensively by Mr. Startin, surgeon to the London Cutaneous Institution, and he has published several formulæ for its employment. Two of them are subjoined.

Lotion with Borax and Glycerin.

R. Sodæ Boratis, 3ss. ad 3j.

Aquæ Rosæ, f 3vijss.

Glycerinæ, f 3ss.

Fiat mistura.

Used for chapped hands, fissures of the lips, and irritation of the skin of any kind.

Lotion of Ammonia, Glycerin, &c.

R. Spiritûs Ammoniaë Aromatici, f 3j.

Glycerinæ, f 3ss.

Tincturæ Cantharidis, f 3j. ad f 3ij.

Aquæ Rosmarini, f 3vij.

Misce. *Signa*.—To be used with a wet hair-brush once or twice a day.

Employed for alopecia following fevers, or for falling off of the hair, dryness or want of action of the scalp, thinness of the hair, &c.

STARTIN.

Solution of Tartar Emetic.

R. Antimonii et Potassæ Tartratis, ʒj.

Aquæ destillatæ, f 3j.

Fiat solutio.

Used to cleanse foul ulcers, and to repress fungous growths, and venereal warts.

SIR W. BLIZARD.

¹ When aromatic waters are prescribed, which are not officinal, the compounder can readily prepare them by rubbing two drops of the essential oil, with a few grains of carbonate of magnesia, and a fluidounce of water, and then filtering the solution through bibulous paper.

Solution of Tannin.

R. Acidi Tannici, gr. v.
Aquæ destillatæ, f 3j.

Fiat solutio.

Used for sore nipples, by applying it to the part on lint covered with oiled silk. DRUITT.

Lotion of Green Vitriol.

R. Ferri Sulphatis, 3j.
Aquæ, Oj.

Fiat lotio.

Used in erysipelas, by applying it to the part every two or three hours. VELPEAU.

Lotion of Borax and Morphia.

R. Sodæ Boratis, 3ss.
Morphiæ Sulphatis, gr. vj.
Aquæ Rosæ, f 3viij.

Fiat lotio.

Used in pruritus vulvæ. Wash the surfaces with soap and tepid water, and dry them before applying the lotion, which is to be done three times daily. C. D. MEIGS.

Lotion of Hyposulphite of Soda, &c.

R. Sodæ Hyposulphitis, 3j. ad ij.
Aluminæ Sulphatis, 3j. ad ij.
Aquæ Rosæ, f 3viijss.

Fiat lotio.

To the above a tablespoonful of some flavoring ingredient, as the Eau de Cologne, may be added. Used in acne by washing the part with a linen rag dipped in the solution twice or thrice a day. STARTIN.

Labarraque's Disinfecting Liquid.

R. Liquoris Sodæ Chlorinatæ, f 3ij.

This stimulant and antiseptic solution has been extensively employed in low forms of disease, as typhus, scarlatina maligna, &c. It is also used as a gargle, as an injection in coryza and ozæna, and as a wash for various foul ulcers. As a gargle or injection half a fluidram to an ounce of water, and as a wash one or two fluidrams to the ounce, are about the proper proportions.

E.—OINTMENTS.

Ointment of Red Precipitate.

R. Hydrargyri Oxidi Rubri, ʒj.
Terebinthinæ Venetæ, ʒj.
Fiat unguentum.

This preparation is applied to indolent sores, and is thought by some practitioners to be a specific in itch.

Simple Iodine Ointment.

R. Iodinii, gr. iij.
Adipis, ʒij.
Fiat unguentum.

Used as the preceding preparation.

Ointment of Iodide of Potassium.

R. Potassii Iodidi, ʒj.
Cerati Simplicis, ʒj.
Fiat ceratum.

Applied to scrofulous swellings, when the skin is unbroken. It may also be used for tetter, and kindred eruptions.

Lugol's Ointment of Ioduretted Iodide of Potassium.

R. Iodinii, gr. xij.
Potassii Iodidi, ʒiv.
Adipis, ʒij.
Fiat unguentum.—Used in scrofulous ophthalmia, ulcers, &c.

Ointment of Iodide of Zinc.

R. Zinci Iodidi, ʒj.
Adipis, ʒj.
Fiat unguentum.—A dram to be rubbed on twice a day. In tumors.

Dr. Ure recommends the above as a substitute for the ointment of iodide of potassium in glandular swellings.

Ointment of Iodide of Barium.

R. Barii Iodidi, gr. iv.
Adipis, ʒj.
Fiat unguentum.
For scrofulous swellings.

BIETT.

Ointment of Iodide of Iron.

R. Ferri Iodidi, ℥iss.

Adipis, ʒj.

Fiat unguentum.

Ointment of Iodide of Sulphur.

R. Sulphuris Iodidi, gr. xxv.

Adipis, ʒj.

Fiat unguentum.—In diseases of the skin.

MAGENDIE.

Ointment of Iodide of Arsenic.

R. Arsenici Iodidi, gr. iij.

Adipis, ʒj.

Fiat unguentum.

In lepra, impetigo, and other cutaneous affections.

BIETT.

Ointment of Iodine and Calomel.

R. Iodinii, gr. x.

Hydrargyri Chloridi Mitis, gr. xv.

Adipis, ʒj.

The iodine to be first united to the lard, and then the calomel added.

MM. Planck and Soubeiran, by whom this ointment is proposed, state that these are the only proportions that can be used, without leading to decompositions that will defeat the intention.

Ointment of Iodide of Mercury, &c.

R. Hydrargyri Iodidi, gr. vj.

Morphiæ Acetatis, gr. viij.

Adipis, ʒj.

Fiat unguentum.

Recommended by the French physicians in glandular swellings. The morphia is supposed by them to prevent the unpleasant effects of iodine when used as an ointment.

PELLETAN.

Ointment of Red Iodide of Mercury.

R. Hydrargyri Iodidi Rubri, ʒj.

Adipis, ʒjss.

Fiat unguentum.—Used to accelerate the cicatrization of inveterate venereal ulcers.

An ointment may be made in the same proportions of the *deutiodide of mercury*; but should be used with more caution.

Ointment of Red Iodide of Mercury.

R. Hydrargyri Iodidi Rubri, gr. xv.

Adipis, ʒij.

Olei Limonis, gtt. xx.

Fiat unguentum.

Employed as a curative application to syphilitic tubercles, chronic ulceration, and other sequelæ of venereal. Its great activity requires caution in its use. RATIER.

Ointment of Ioduretted Iodide of Potassium with Opium.

R. Iodinii, gr. xvj.

Potassii Iodidi, ʒj.

Tincturæ Opii, fʒij.

Adipis, ʒij.

Fiat unguentum.—Applied to scrofulous and other ulcers.

LUGOL.

Ointment of Cyanuret of Mercury.

R. Hydrargyri Cyanureti, gr. xij. ad xvj.

Adipis, ʒj.

Fiat unguentum.—Used in herpes attended by inflammation and itching.

Mercurial Ointment.

R. Unguenti Hydrargyri, ʒj.

One dram of this ointment to be rubbed on the inside of the thighs, before the fire.

Some physicians recommend the warm bath before using this ointment, while others consider it useless. We think the practice occasionally beneficial. The hands of the person rubbing it on should be enveloped in a bladder, or gloves made of oiled silk.

Mercurial Ointment with Camphor.

R. Unguenti Hydrargyri, ʒj.

Camphoræ, ʒj.

Misce.—To be employed as the former.

White Precipitate Ointment.

R. Hydrargyri Ammoniati, gr. xv.

Potassæ Nitratis, ʒss.

Sulphuris, ʒj.

Bene terantur et adde—

Adipis, ʒij.

Fiat unguentum.

This ointment has long been celebrated in the cure of itch.

Anti-herpetic Ointment.

- R. Hydrargyri Sulphatis Flavi, ʒj.
 Tincturæ Opii, fʒj.
 Sulphuris sublimati, ʒss.
 Adipis, ʒj.

Fiat unguentum.—In herpes, alternated with the simple water bath.

Sulphur Ointment.

- R. Adipis, lbj.
 Sulphuris sublimati, ʒviij.

Fiat unguentum.—Used in friction, three to six drams at a time.

Mercurial Cerate.

- R. Unguenti Hydrargyri, ʒiv.
 Cerati Simplicis, ʒx.

Fiat ceratum.—A dressing for venereal ulcers. RATIER.

Ointment of Belladonna.

- R. Extracti Belladonnæ, ʒij.
 Aquæ destillatæ, fʒij.
 Adipis, ʒij.

Misce.—Employed by M. Chaussier as a direct application to the neck of the uterus, in cases of rigidity. RATIER.

Ointment of Alum, Calomel, &c.

- R. Hydrargyri Chloridi Mitis, ʒij.
 Aluminis exsiccati,
 Plumbi Oxidi Semivitrei, āā ʒss.
 Olei Terebinthinæ, fʒij.
 Unguenti Simplicis, ʒiss.

Fiat unguentum.

This ointment is particularly adapted to *tinea capitis*. It should be applied every night going to bed, and washed off in the morning: as soon as the child's head is dry, dust it over with powdered starch, which, in the evening, is again to give place to the ointment. The hair must be cut off previous to the application of the ointment.

Ointment of Galls.

- R. Pulveris Gallæ, ʒj. vel ʒij.
 Adipis, ʒj.

Fiat unguentum.—In hemorrhoidal affections.

Ointment of Oxide of Zinc.

R. Zinci Oxidi, ʒj.

Butyri recentis, ʒvj.

Fiat unguentum.—To be applied to the eyes in defluxions arising from relaxation or slight inflammation of the lids.

Kentish's Ointment.

R. Cerati Resinæ, ʒij.

Olei Terebinthinæ, fʒij. vel fʒss.

Fiat unguentum.

This preparation has long been celebrated in burns and scalds. Care is requisite to avoid touching the sound skin, for turpentine occasionally inflames it severely.

Ointment of Bromide of Potassium.

R. Potassii Bromidi, gr. xxxiv. ad ʒj.

Adipis, ʒj.

Fiat unguentum.

Ointment with Goulard's Cerate, Calomel, &c.

R. Cerati Plumbi Subacetatis,

——— Simplicis, āā ʒss.

Hydrargyri Chloridi Mitis,

Pulveris Opii, āā ʒj.

Misce.—Useful in various cutaneous eruptions.

JOS. PARRISH.

Ointment of Rhubarb, Opium, &c.

R. Pulveris Rhei, gr. x.

——— Opii, gr. v.

Unguenti Simplicis, ʒjss.

Fiat unguentum.—In indolent ulcers.

Ointment of Tar, &c.

R. Unguenti Picis Liquidæ, ʒj.

——— Hydrargyri Nitratis, ʒss.

Fiat unguentum.—In tinea capitis.

Ointment of Tar and Opium.

R. Pulveris Opii, ʒij.

Unguenti Picis Liquidæ, ʒj.

Fiat unguentum.

For hemorrhoids.

Kirkland's Neutral Cerate.

R. Emplastri Plumbi, ℥iv.

Olei Olivæ,

Cretæ Præparatæ,

Acidi Acetici, āā ℥ij.

Plumbi Acetatis, ℥jss.

The plaster and oil are first to be melted over a gentle fire. The prepared chalk is then to be briskly stirred in, and when the mixture is sufficiently cooled, the vinegar and superacetate of lead are to be added, and the whole stirred until nearly cold.

This cerate, which is not to be met with in the pharmacopœia, is well adapted to abraded irritable surfaces, and the later stages of burns.

Aromatic Sulphur Ointment.

R. Potassæ Carbonatis, ℥ss.

Aquæ Rosæ, f ℥j.

Hydrargyri Sulphureti Rubri, ℥j.

Olei Bergamii, f ℥ss.

Sulphuris sublimati et loti,

Adipis, āā ℥x.

Misce secundum artem.—In scabies. BATEMAN'S *Synopsis*.

Ointment with Carbonate of Ammonia, &c.

R. Ammoniæ Carbonatis, ℥ss.

Unguenti Simplicis, ℥ss.

Fiat unguentum.—Applied to indolent ulcers of a scrofulous character.

Ointment with Verdigris, &c.

R. Cupri Subacetatis,

Cupri Sulphatis, } āā ℥ij.

Hydrargyri Oxidi Rubri, }

———— Chloridi Corrosivi, ℥j.

Adipis, q. s.

Ut fiat unguentum.

This may be spread on lint, and one or two applications will be sufficient to destroy a very large mass of fungous granulations.

B. C. BRODIE.

Ointment of Ipecacuanha.

R. Pulveris Ipecacuanhæ, ʒij.
 Olei Olivæ, fʒij.
 Adipis, ʒss.

Misce, et fiat unguentum.

Useful as a counter-irritant.

TURNBULL.

Ointment of Calomel and Tar.

R. Hydrargyri Chloridi Mitis, ʒj.
 Unguenti Cetacei, Lond. ʒj.
 ——— Picis Liquidæ, ʒss.

Fiat unguentum.—Used in lepra.

A. T. THOMSON.

Ointment of Creasote and Charcoal.

R. Creasoti, fʒss.
 Alcoholis, fʒjss.
 Carbonis Animalis purificati, ʒj.
 Unguenti Cetacei, Lond. ʒjss.

Fiat unguentum.

Used in cases of burns and scalds. "For children under five years, the ointment should be weakened by mixing it with four times its bulk of spermaceti; for children from five to ten years, with twice its bulk of spermaceti; and so on, gradually increasing the strength of the ointment according to the age of the patient."

S. SUTRO.

Oxide of Silver Ointment.

R. Argenti Oxidi, ʒj.
 Unguenti Simplicis, ʒj.

Fiat unguentum.

Used for sore nipples.

Ointment of Slaked Lime, Zinc, &c.

R. Calcis, ʒj
 Camphoræ, ʒj.
 Unguenti Zinci Oxidi, ʒj.

Fiat unguentum.

This is the "topique contre acné" of the

HÔPITAL ST. LOUIS.

Citrine Ointment.

R. Unguenti Hydrargyri Nitratis, ʒj.

When well prepared, this is one of our most efficient ointments in a great variety of cutaneous diseases.

Depilatory Ointment.

R. Sodæ Carbonatis, ʒij.

Calcis, ʒj.

Adipis, ʒj.

Fiat unguentum.

Used in porrigo.

BIETT.

Veratria Ointment.

R. Veratriæ, gr. x.

Acidi Acetici, gtt. x.

Adipis, ʒj.

Fiat unguentum.

Veratria has been used externally in neuralgia, and with the most marked success.

Ointment of Nitrate of Silver.

R. Argenti Nitratis, ʒss.

Adipis loti, ʒss.

Fiat unguentum.

Used to smear over bougies in the treatment of gonorrhœa.

MACDONALD.

Ointment of Tannin.

R. Acidi Tannici, ʒj.

Adipis loti, ʒss.

Fiat unguentum.

Used for the same purposes as the astringents generally.

FRONMÜLLER.

POWDERS.

Sometimes the practitioner prefers dusting medicinal agents directly upon diseased surfaces, instead of incorporating them with lard in the form of an ointment. Hundreds of such formulæ might be given, but we limit our selection to two or three.

Powder of Verdigris and Calomel.

R. Cupri Subacetatis,

Hydrargyri Chloridi Mitis, āā ʒj.

Fiat pulvis subtilissimus.—For chancres and indolent ulcers.

Powdered Rhubarb, &c.

R. Pulveris Rhei,

———— Ipecacuanhæ, āā ʒj.

Fiat pulvis.—To be dusted on sarcomatous and watery tumors.

Rhubarb alone forms an excellent application to indolent ulcers.

Powder of Savin, &c.

R. Pulveris Sabinæ,

———— Cupri Subacetatis, āā ʒj.

Fiat pulvis.—To be dusted on sores accompanied with fungous excrescences.

F.—CATAPLASMS.

Cataplasm of Bread and Milk.

Remove the crust from a part of a loaf of stale bread, and crumble the bread into a bowl. Pour over it sufficient sweet milk to cover it, and simmer over a fire, occasionally stirring, until the bread is reduced to a pulpy consistence. The poultice is to be applied as warm as the patient can bear it.

Water may be substituted for the milk without much if any disadvantage.

Cataplasm of Flaxseed Meal and Bran.

Take equal parts of ground flaxseed and bran. Pour boiling water upon them, and stir constantly until the poultice has a pulpy consistence.

In the Northern and Middle States, it is customary to use the common flaxseed meal, which contains both the oily and mucilaginous principles; but in New York city and some other places, the ground oil-cake is preferred. Equal parts of the latter and bran form an elegant poultice.

Mush Cataplasm.

This is made by stirring Indian meal, in small quantities, into water kept boiling in a pan, until the whole has acquired the proper degree of consistence.

Cataplasm of Yeast.

R. Farinæ, *Ed.* ℥j.

Cerevisiæ Fermenti, *Lond.* Oss.

Misce.—This cataplasm is useful in stimulating foul ulcers and gangrenous wounds, and removing the unpleasant odor. It should be made with fresh yeast, and applied warm.

Cataplasm of Charcoal.

This cataplasm is made by stirring into the common bread and milk poultice as much finely powdered charcoal as it will bear. It is a very useful application to offensive ulcers; removing in a short time the fetor, and giving them a more healthy aspect.

Cataplasms of Slippery Elm.

This poultice is made from the inner bark of the slippery elm ground into a fine powder, and moistened with hot water.

Cataplasms are also made of carrots or turnips, by simply boiling them to softness, and beating them to a pulpy mass. Either furnishes a good poultice for cancerous and other painful sores.

Emollient Cataplasm.

R. Farinæ Seminum Lini,

————— Hordei, āā partes æquales.

Aquæ, vel Lactis, q. s. ut fiat cataplasma.

The milk or water may be warmed if desirable. Applied to inflammatory tumors.

Antiseptic Cataplasm.

R. Cataplasmatiss Emollientis, ℥viij.

Pulveris Kino,

————— Camphoræ, āā ℥ij.

Fiat cataplasma.—Applied to unhealthy wounds and ulcers.

HÔTEL DIEU.

Resolvent Cataplasm.

R. Cataplasmatiss Emollientis, ℥iv.

Plumbi Acetatis, ℥j.

Ammoniæ Muriatis, ℥ss.

Fiat cataplasma.—Applicable to tumors that require gentle excitement.

RATIER'S *Formulary*.

Cataplasm of Common Salt, &c.

R. Pulveris Lini,

Micæ Panis, āā partes æquales.

Liquoris Sodii Chloridi saturati, q. s. ut fiat cataplasma.

To be applied to indolent strumous swellings, and enlargements of the joints. When it produces irritation or inflammation, the common bread and milk poultice must be substituted; and the other reapplied when circumstances permit.

Iodine Cataplasm.

R. Tincturæ Iodinii, f℥ss.

Pulveris Lini, ℥j.

Avenæ Farinæ, ℥iij.

Aquæ q. s. ut fiat cataplasma.

Applied to indolent ulcers (especially those of a scrofulous character), and to tumors of the same class.

G.—B A T H S .

The ordinary application of cold, tepid, and shower baths, is too well known to require any particular notice. The hot bath should be of a temperature of 93 to 96 degrees of Fahrenheit. It is designed to alleviate general or local irritation, and to induce repose and sleep. It is especially adapted to fevers with deficient capillary circulation, rheumatism, herpetic eruptions, paralytic affections, &c. The lower temperature is best adapted to cutaneous complaints, and to bring on relaxation of the skin during febrile irritation; the warmer temperature is suited to paralysis; and, as a general rule, more heat is desirable to act upon deep-seated than on superficial diseases.

Vapor Baths are produced by the evaporation of boiling water in a suitable apparatus. They are used, like ordinary warm baths, in rheumatic and cutaneous affections, and in every case wherein profuse perspiration is desirable. They are avoided in persons laboring under pulmonary or cardiac disease, and in exhaustion from any cause.

Local Baths are admirably adapted to parts which are swelled and painful, and may consist of tepid water only, or their strength may be increased by the addition of saline, emollient, narcotic, or stimulant substances. Among these salt and mustard are preferable.

Sulphurous Water, or Artificial Bareges Bath.

R. Potassii Sulphureti, ℥j.

Aquæ, cong. xxx.

Mix for a bath. In cutaneous eruptions.

Sulphureo-gelatinous Bath.

R. Potassii Sulphureti, ʒij. vel ʒiv.

Aquæ, cong. xxx.

Add to this solution,

Ichthyocollæ, ℥j. vel ℥ij. in Aquæ bullientis solutæ ℥x.

This bath, according to Dr. Green (*Diseases of the Skin*, p. 240), "is preferable to the artificial Bareges bath, as it is neither irritating nor apt to occasion feverishness, which the common sulphureous water-bath is. A cheaper and not less efficacious gelatine may be procured by dissolving from a pound and a half to two pounds of parchment-clippings in water by long boiling, or by using a neat's or a calf's foot for the purpose."

Emollient Bath.

To an ordinary tepid water-bath, add a large cupful of thick gruel or paste, and mix it well with the water. These baths are suited to prurigo, eczema, lichen, and impetigo.

Nitro-muriatic Acid Bath.

R. Acidi Nitrici, fʒijj.

— Muriatici, fʒj.

Mix, and add to the water of a tepid bath, which should then be about as sour as distilled vinegar.

Corrosive Sublimate Bath.

R. Hydrargyri Chloridi Corrosivi, ʒij.

Aquæ tepidæ, cong. xxx.

Mix for a bath. Prescribed by the French physicians in syphilitic affections, when it is desirable to avoid the action of the medicine on the stomach. This plan is seldom resorted to, on account of the great risk of employing corrosive sublimate in such large quantity.

Alkaline Bath.

R. Potassæ Carbonatis, ʒiv. vel ʒviiij.

Aquæ tepidæ, cong. xxx.

Mix for a bath. For promoting desquamation from the skin, and to allay itching in various forms of prurigo.

Ioduretted Baths.

	No. 1.	No. 2.	No. 3.
R. Iodinii,	℥ij.	℥iij.	℥iv.
Potassii Iodidi,	℥iv.	℥v.	℥viij.
Aquæ destillatæ,	f℥vj.	f℥vj.	f℥vj.
Fiant solutiones.			

One of these solutions, according to the strength required, is added to a tepid bath in a wooden tub. For children, the drams may be changed for scruples. Used in scrofula.

Artificial Harrowgate Bath.

R. Sodii Chloridi, ℔ij.
 Magnesiae Sulphatis, ℥iij.
 Potassii Sulphureti, ℔j.
 Aquæ, cong. xxx.

Mix for a bath.

H.—F O M E N T A T I O N S .

Fomentation is a yet more restricted mode of applying fluids to the surface of the body, and is best accomplished by means of a sponge, flannel, or soft cloth. To be efficacious it should be often repeated, and not allowed to get cold.

Stimulant Fomentation.

R. Vini Rubri, Oij.
 Mellis, ℥iv.

Misce.—This is an active fomentation, adapted to parts in a low state of vitality. RATIER.

Saponaceous Fomentation.

R. Alcoholis, Oij.
 Saponis, ℥j.

Misce.—A resolvent in indolent tumors and in sprains.

RATIER.

Narcotic Fomentation.

R. Opii, ℥ij.
 Aquæ, Oj.

Solve.—For painful affections of the joints, as chronic rheumatism, &c. RATIER.

Narcotic Fomentation.

R. Dulcamaræ, ℥ij.
 Capsularum Papaveris, ℥ij.
 Aquæ ferventis, Oj.

Fiat infusum.—Employed as a soothing application to irritable surfaces.

Mustard Fomentation.

R. Farinæ Sinapis, ℥iv.
 Aquæ ferventis, Oj.

Misce.—By wetting cloths in this fluid and applying them to the skin, irritation may be prolonged to any requisite degree.

Tobacco Fomentation.

R. Tabaci, ℥ij.
 Aquæ ferventis, Oj.

Misce.—Employed in psora, but is apt to produce nausea and vomiting. Its use requires great care.

I.—FUMIGATIONS.

Fumigations are for the most part used to purify infected air by absorbing or otherwise counteracting deleterious gases. They are also employed in diseases of the skin, and may be sometimes substituted for a local bath. The ordinary process for fumigating apartments with chlorine, has been already described. (Page 109.)

Fumigations of sulphur, cinnabar, and mercury, are also in common use for medicinal purposes, and are readily applied with a proper apparatus.

Fumigating Powders.

R. Sulphuris, ℥iij.
 Hydrargyri Sulphureti Rubri, ℥ij.
 Iodinii, gr. x.
 Fiant pulveres vj.

One of the powders to be used in the following manner: "If the disease is seated on the limbs, a tin case, or even a common jar, should be procured. A heated iron is to be placed at the bottom of this apparatus, with a grating above it to protect the foot or hand. One of the powders being placed on this heated iron, the limb is to be instantly put into the bath, the top of which should be covered over, to prevent the vapor from escaping. The limb may be continued in the bath for from fifteen to twenty minutes, according to circumstances." Used in lepra.

T. H. BURGESS.

BOOK III.

OF

DIETETIC PREPARATIONS, AND POISONS.

DIETETIC PREPARATIONS.

THE importance of an acquaintance with those preparations of diet peculiarly designed for the patient during disease and convalescence, is so obvious that it requires little or no comment. It was remarked by the late Dr. Rush in his lectures, that a physician should spend six months in a kitchen before entering upon his practical career. As the medical attendant, therefore, is called upon frequently to give directions and rules for preparing the diet for the sick, we have inserted a few formulæ for those articles which are in daily use. However apparently trifling such a class may appear to some persons, the physician will find that an attention to it will place the patient more completely under his own control, and inspire greater confidence in the relations and attendants. The rules to be observed in giving nourishment to the sick, will generally present themselves; but there is one in particular which should not be overlooked. In all low diseases, the nurse should be directed never to give too much at once, but frequently to repeat a small proportion. The lapse of a very short time without food, will frequently, in cases of great prostration, reduce the patient very low, and may endanger his life.

The rule is not intended to be applied to those chronic derangements of the digestive functions, which are to be remedied principally by a regulation of diet, and in which the old adage, "a little and often," has too frequently, we are convinced, been perniciously advised.

Decoction of Barley; or, Barley Water.

Take of

Pearl Barley,	2 ounces,
Boiling Water	2 quarts.

Before adding the boiling water, let the barley be well washed.

Then boil it to one-half, and strain the liquor. A little lemon-juice and sugar may be added if desirable. To be taken *ad libitum* in inflammatory diseases.

Rice Water.

Take of

Rice, 2 ounces,
Let it be well washed, and add to it—

Water, 2 quarts.
Boil it for an hour and a half, and then add sugar and nutmeg
as much as may be required. To be taken *ad libitum*.

Rice, when boiled for a considerable time, assumes a gelatinous form, and, mixed with milk, is a very excellent diet for children. It possesses in some measure a constipating property, which may be increased by boiling the milk.

Infusion of Flaxseed.

As under Expectorants. (Page 104.)

Mucilage of Gum Arabic.

As an article of diet, the proper proportions are an ounce of Gum Arabic to a pint of boiling water. The solution is allowed to cool before it is used. Gum Arabic is very nutritive, and life can be sustained on it alone for some time.

Decoction of Bran.

Take of

Fresh Wheat Bran, 1 pint,
Water, 3 quarts.

Boil down one-third, strain off the liquor, and add sugar, honey, or molasses, according to the taste of the patient.

Bran tea may be made by using boiling water, and suffering the mixture to stand in a covered vessel for three or four hours.

Sage Tea.

Take of

The dried Leaves of Sage, half an ounce,
Boiling Water, 1 quart.

Infuse for half an hour, and then strain. Sugar and lemon-juice may be added in the proportion required by the patient. In the same manner may be made balm and other teas.

These infusions form very agreeable and useful drinks in fever, and their diaphoretic powers may be increased by the addition of sweet spirits of nitre or antimonial wine.

Infusion of Malt.

Take of

Ground Malt, 1 pint,
Scalding Water, 3 pints.

Infuse for two hours, and strain off the liquor, to which may be added sugar or lemon-juice, if required.

This was a favorite preparation with the late Dr. Jos. Parrish, in inflammatory fevers.

Lemonade.

Take of

Fresh Lemon Juice,	4 ounces,
Fresh and very thin Peel of Lemon,	half an ounce,
White Sugar,	4 ounces,
Boiling Water,	3 pints.

Let them stand until cold, then strain off for use. When employed in fevers, a little nitrate of potash or sweet spirits of nitre may be added. It can be farther diluted to the taste of the patient.

This preparation, when freely taken, sometimes produces pain in the bowels. It must therefore be used with some reserve as a daily drink.

Water Gruel.

Take of

Oatmeal,	2 ounces,
Water,	1 quart.

Boil for ten or fifteen minutes and strain, adding sugar and salt to make it agreeable.

Milk for Infants.

Take of

Cow's Milk,	1 part,
Water,	2 parts.

Loaf Sugar as much as may be agreeable.

It is necessary, when children are to be raised by the hand, to dilute the milk. These proportions may be altered as the child advances in age. The object is to make a diet as near the qualities of the maternal milk as possible.

Rice Gruel.

Take of

Ground Rice,	1 ounce,
Cinnamon,	1 dram,
Water,	1 quart.

Boil for forty minutes, adding the aromatic near the conclusion. Strain and sweeten. Wine may be added if necessary.

Panado.

Take of

Wheat Bread,	1 ounce,
Cinnamon,	1 dram,
Water,	1 pint.

Boil them until well mixed, then add a little grated nutmeg and sugar. Wine or butter may also be added if desirable.

Compound Salep Powders.

R. Salep,
 Tragacanth,
 Sago, of each ʒiv .
 Cochineal, ʒss .
 Prepared Oyster-shells, ʒj .

To be carefully mixed and divided into powders of ʒj . each. Stir one of these powders into a pint of milk, and boil them together for ten or fifteen minutes. To be drank freely in dysentery and diarrhoea.

These are the dysenteric powders of Dr. Castanello. A somewhat analogous preparation which bears the same name, is prepared as follows:—

R. Gum Arabic,
 Tragacanth,
 Arrowroot,
 Sago,
 Tapioca, of each ʒij .

Mix them well together, and boil in a pint of milk, flavored with nutmeg or cinnamon. Used as a diet in dysentery, diarrhoea, &c. &c.

Sago.

Take of

Sago,	2 tablespoonfuls,
Water,	1 pint.

Boil gently until it thickens, frequently stirring. Wine, sugar, and nutmeg may be added according to circumstances.

Tapioca Jelly.

Take of

Tapioca,	2 tablespoonfuls;
Water,	1 pint.

Boil it gently for an hour, or until it assumes a jelly-like appearance. Add sugar, wine, and nutmeg, with lemon-juice, to suit the taste of the patient and the nature of the case.

Jelly of Carrageen, or Irish Moss.

Take of

Carrageen,	half an ounce,
Fresh Milk,	a pint and a half.

Boil them down to a pint; remove any sediment by filtering or otherwise, and then add the requisite quantity of sugar, with lemon-juice, or peach water, to give it an agreeable flavor. To be used freely.

The moss, before being used, should be well washed in cold water to remove its saline taste.

Arrowroot.

Take of

Arrowroot,	1 tablespoonful,
Sweet Milk,	half a pint,
Boiling Water,	half a pint.

Boil them a few moments over the fire.

Sweetened with loaf sugar, it is one of the best aliments for children, especially when the bowels are irritable.

The farinaceous substance called *Canna* is prepared by simply boiling in milk, and adding sugar.

[Arrowroot is better prepared in the following manner:—

Take of

Arrowroot,	1 large tablespoonful,
Cold Water,	sufficient,
Boiling Water,	half a pint.

Mix the arrowroot with just sufficient cold water to soften the lumps, then pour on the boiling water, and stir continually until the solution is gelatinous and transparent. Sweeten to the taste. EDITOR.]

Boiled Flour.

Take of

Fine Flour,	a pound.
-------------	---	---	---	---	---	----------

Tie it up in a linen cloth as tight as possible, and after frequently dipping it in cold water, dredge the outside with flour till a crust is formed round it, which will prevent the water soaking into it while boiling. It is then boiled until it becomes a hard, dry mass.

Two or three spoonfuls of this may be grated and prepared in the same manner as arrowroot, for which it forms an excellent substitute.

Vegetable Soup.

Take one turnip, one potato, one onion. Let them be sliced, and boiled in one quart of water for an hour. Add as much salt as is agreeable, and pour the whole upon a piece of dry toast.

This preparation forms an agreeable substitute for animal food, and may be given when the latter is inadmissible.

Chocolate.

Chocolate should be suffered to stand until cold, after having been made in the usual way. The oily parts thus collect on the surface and should be taken off. The liquid should then be boiled again, and sugar, &c. added.

Beef Tea.

Take of

Lean Beef, cut into shreds, 1 pound,
Water, 1 quart.

Boil it for twenty minutes, taking off the scum as it rises. After it grows cold, strain the liquor.

This preparation is more nourishing than ordinary broths, and very palatable.

Another Plan for Beef Tea.

"One pound of lean beef, free of fat, and separated from the bones in the finely chopped state in which it is used for mince meat, is uniformly mixed with its own weight of cold water slowly heated to boiling, and the liquid, after boiling briskly for a minute or two, is strained through a towel, from the coagulated albumen and fibrine, now become hard and horny. Thus we obtain an equal weight of the most aromatic soup, of such strength as cannot be obtained even by boiling for hours from a piece of flesh." This is to be seasoned to taste.

DR. BENEKE.

Essence of Beef.

Take of

Lean Beef, sliced, 1 pound.

Put a sufficient quantity into a porter bottle to fill up its body, cork it *loosely* and, place it in a pot of cold water, attaching the neck by means of a string to the handle of the vessel. Boil this for an hour and a half or two hours, then decant the liquor and skim it.

To this preparation may be added spices, salt, wine, brandy, &c., according to the taste of the patient and nature of the disease.

Hartshorn Jelly.

Take of

Hartshorn Shavings,	4 ounces,
Water,	1 quart.

Boil over a gentle fire until one pint of the water be dissipated.

Strain, and add lemon-juice, sugar, and wine.

This forms, either with or without the last-named ingredients, a very nourishing diet for convalescents; and, when mixed with an equal portion of milk, is well adapted to the bowel complaints of children.

Isinglass or fish glue may be used instead of the hartshorn shavings, if preferred.

Calf's Feet Jelly.

Take two calves' feet and add to them one gallon of water, which reduce by boiling to one quart. Strain it, and when cold, skim the fat entirely off. Add to this the white of six or eight eggs well beaten, a pint of wine, half a pound of loaf sugar and the juice of four lemons, and let them be well mixed. Boil the whole for a few minutes, stirring constantly, and then pass it through a flannel strainer.

This forms a very nutritious article of diet for the sick and convalescent. The wine may be omitted or added at option.

Chicken Water.

Take half a chicken divested of all fat, and break the bones; add to this half a gallon of water, and boil for half an hour. Season with salt.

This water is directed by some physicians in the early stages of cholera. Taken warm it promotes vomiting and washes out the stomach.

Suet Ptisan.

Take of

Sheep's Suet,	2 ounces,
Milk,	1 pint,
Starch,	half an ounce.

Boil slowly for half an hour.

This may be used as a common drink in dysentery.

Mucilage of Starch, &c.

Take of

Starch,	1 ounce,
Powdered Cinnamon,	1 dram,
———— Gum Arabic,	1 ounce,
Boiling Water,	3 pints.

Boil until reduced one-third, and strain.

The above may be taken for a common drink in dysentery.

Wine and Mustard Whey.

As under Stimulants. (Page 131.)

Rennet Whey.

Take of

New Milk,	1 quart,
Rennet,	a large spoonful.

Heat the milk and then add the rennet. Boil until the curd separates and is all taken off.

To many persons this forms a very agreeable nutriment.

Vinegar Whey.

Take of

Milk,	1 pint,
Vinegar,	half an ounce.

Boil for a few minutes, and separate the curd.

Tartar Whey.

Take of

Milk,	1 quart,
Cream of Tartar,	half an ounce.

Boil until the curd separates.

OF POISONS.

CASES of poisoning are so sudden and so awful that no time is to be lost in administering relief. To do this with judgment and effect, it is requisite to be familiar with all the usual antidotes; and of these, we have accordingly selected some which possess an acknowledged efficacy, omitting at the same time many others that are unworthy of confidence. For the purpose of convenience we have arranged the *Poisons* alphabetically, with the appropriate *antidotes* attached to each. But, as has been remarked when speaking of Emetics, our chief reliance in these cases is upon free vomiting, induced by those articles which are most speedy in their effect. Dr. Beck prefers sulphate of zinc, whenever it can be procured, and adds, "but if vomiting be present, we may aid it by diluents or a vegetable emetic. Tartrate of antimony should never be given." Yet, even from this treatment, little can be expected when the deleterious substance has remained long in the stomach; and the stomach-tube and syringe will be indispensable.

To protect the stomach and intestines from the direct operation of poisons, it is generally recommended to give milk, lime-water, soap, or drinks sweetened with sugar or honey. Oils and other fatty substances are found to be injurious.

When the vomiting has ceased, and retching and pain continue, Mr. Hume advises the free administration of carbonate of magnesia with tincture of opium, suspended in water.

ALCOHOL.

Large quantities of alcoholic liquors, taken by persons unaccustomed to their use, often produce fatal effects.

The treatment in such cases is to evacuate the stomach as soon as possible; but from the rapidity of absorption this may not always be effectual. The patient must be kept roused and

cold affusion plentifully used. Large draughts of tea or coffee ought to be taken until the stomach is thoroughly evacuated.

ALKALIES.

Ammonia, potash, soda, and their carbonates, are seldom taken in poisonous quantities; when it so happens, the best antidote is common vinegar; it is always at hand, and its affinity for these substances is sufficiently strong to neutralize them. Also, a *fixed oil*, as olive oil, should be given in large quantity—several pounds being required where a large portion of the alkali has been taken; it acts by rendering the vomiting easy and free, and partly by converting the alkali into a soap.

ANTIMONY.

The preparation of this mineral by which poisoning is commonly produced is *tartar emetic*. This of itself produces violent vomiting, and therefore our first efforts should be directed to keeping up the emesis by means of mild diluents and mucilaginous preparations. Its antidotes are the alkalies, tannin, bitter and astringent infusions, such as common tea, Peruvian bark, especially the yellow bark and its decoction—the administration of galls in *powder*, &c. &c. These decompose the antimonial salt. When the vomiting is excessive, opium by injection will assist in allaying it.

ARSENIC.

Until lately, there has been no known antidote to this deadly poison, and our sole dependence has been upon mild diluents, albumen (such as the whites of eggs suspended in milk) or warm water and sugar. Various substances have been proposed as antidotes, but without success. It is best first (if vomiting has been delayed) to give an emetic of the sulphate of zinc, and milk should be drunk liberally both before and after vomiting has begun. The patient should never be allowed to exhaust his strength in retching, without some milk or other fluid in his stomach to act on.

THE HYDRATED PEROXIDE OF IRON, freshly prepared, was announced in the year 1834, by Dr. Bunsen, of Göttingen, as an antidote for poisoning by arsenic; and its efficacy has been confirmed by MM. Orfila, Lesueur, Soubeiran, and Miguel, and also in this city by various chemists, and by experiments upon the inferior animals. The antidote is now officinal, and should always be kept on hand.

The following direction will enable the most ignorant person to make use of this antidote until medical advice can be obtained. It must be administered as soon as possible after the discovery that arsenic has been taken, and as it causes no bad effect itself, should be given EVERY FIVE OR TEN MINUTES, UNTIL ENTIRE RELIEF IS OBTAINED. It is said that twelve parts of it are necessary to neutralize one of arsenic. The dose for a grown person is a tablespoonful; for children a teaspoonful. The bottle to BE WELL SHAKEN BEFORE EACH DOSE. IN CASES OF EMERGENCY, IT MAY BE PREPARED BY SIMPLY PRECIPITATING THE MURIATED TINCTURE OF IRON by ammonia, and washing the precipitate.

Should the patient survive (and the probability is in his favor if the antidote be faithfully given), the inflammatory symptoms must be combated on the general principles for treating inflammation. The diet should consist exclusively of milk, gruel, cream, rice, and similar bland articles.

In a case of poisoning in this city, where half an ounce of the powdered arsenic had been taken and was followed by extreme thirst and vomiting, all the symptoms were allayed by freely taking the hydrated peroxide of iron. Only a slight gastritis followed, which was readily subdued. The oxide forms with the arsenious acid a proto-arseniate of iron.

In place of the hydrated oxide, the rust of iron and precipitated carbonate should be resorted to in cases of emergency, possessing as they do, to a certain extent, the power of antidotes. Magnesia in large doses acts also as an antidote, and in several cases lately reported was used with complete success.

When the poison has been taken in the form of Fowler's solution, Dr. Dunglison recommends copious draughts of lime-water to be used.

BARYTA.

The carbonate and muriate of this earth are poisonous, and the latter has been taken by mistake for Glauber's salts. In cases of poisoning with the carbonate, the alkaline sulphates should be given with a mixture of vinegar, and resort had to the stomach pump and emetics. The true antidote is sulphate of magnesia or soda; with these a sulphate of baryta is formed, which is perfectly insoluble, and may be brought off by emetics and copious diluents, oleaginous or albuminous.

COPPER.

The sulphate (blue vitriol), and subacetate (verdigris), are the salts of this metal from which poisoning most frequently re-

sults; the latter often from the careless use of culinary vessels or mineral water fountains. Producing violent emesis, this should be encouraged, by the free use of tepid water, milk, and mucilages.

According to Orfila, albumen in the form of white of eggs is the best antidote, and sugar, formerly in much repute, acts only as an emollient. In the absence of eggs, milk or wheaten flour should be employed. Vinegar in all cases must be avoided. The French physicians have tried iron-filings, in consequence of the power possessed by this metal of reducing the salts of copper; and the results they obtained, in some experiments performed on animals, are such as to justify considerable confidence in the antidote.

GOLD.

The nitro-muriate of gold is a powerful poison. Its antidotes are sulphate of iron (which throws down the gold in a metallic state), the free use of mucilaginous drinks, and albuminous substances.

IODINE AND IODIDE OF POTASSIUM.

Ten or twenty grains of Iodine sometimes, though not always, produce constriction of the fauces, nausea, vomiting, and severe colic pain. The supposed antidote for these preparations is starch freely given, or, what answers the same purpose, flour mixed with water. Magnesia is also recommended.

LAUDANUM.

Laudanum, when taken in large quantity, must be withdrawn from the stomach by active emetics, or by tickling the fauces. The stomach-tube should then be used to remove whatever remains of the poison. The free use of vegetable astringents and strong coffee will then be serviceable. An important part of the after treatment is the maintenance of artificial respiration, besides which active stimulants are often required. Iodine, bromine, and chlorine are stated by M. Donné, of Paris, to be antidotes for poisoning by *nux vomica* and opium. M. Orfila thinks that the decoction of galls may with propriety be used as an imperfect antidote, until the narcotic can be evacuated from the stomach.

LEAD.

The acetate is the only preparation which is resorted to as a poison. The sulphate of magnesia or soda is its true antidote, as it converts the poison into a sulphate, which is inert; also the alkaline carbonates and particularly the bicarbonates. Phosphate of soda is also a good antidote. If the patient does not vomit, give an emetic of the sulphate of zinc, which will not only empty the stomach, but convert any remaining soluble lead into an insoluble sulphate.

Milk or albumen should also be given in large quantities, as it has been found that the albuminous principle of milk is a very effectual precipitant of the oxide of lead; a compound which, though not entirely inert, is far less active than the acetate, and tends to prevent the action of the poison as a corrosive on the stomach.

MERCURY.

All the preparations of this metal are poisonous, and corrosive sublimate in particular. Our chief and only dependence, according to M. Orfila, is upon the albumen of eggs, copiously swallowed, or introduced through the *stomach tube*, which should be resorted to when the patient cannot swallow or vomit. Ene-mata of the same should also be used.

If the albumen be taken in sufficient quantities, it decomposes the sublimate, forming a triple compound of albumen, muriatic acid, and calomel. If the remedy excites fresh vomiting, so much the better, and if symptoms of gastritis appear, they are to be treated in the usual way. M. Orfila has also established that a fatal dose of corrosive sublimate may be swallowed by animals with impunity, provided the white of egg, or any other kind of fluid albumen, be introduced shortly before or after into the stomach. Cases are also upon record which establish that it possesses an equal efficacy in man.

The gluten of wheat flour is also strongly recommended on the same principle as an antidote to corrosive sublimate. To administer it, nothing more is requisite than to mix the flour with water. The quantity taken must be very considerable. It is said to reduce the muriate to the state of calomel.

Gluten of wheat flour is also given in the form of an emulsion with soft soap, by mixing partly in a mortar and partly with the hand, 5 or 6 parts of fresh gluten with 50 parts of soft soap.

When neither albumen nor gluten is at hand, milk forms a good substitute.

The only other antidotes deserving of mention here are iron filings, meconic acid, and the hydrated proto-sulphuret of iron. The former was recommended by Dumas, and must obviously operate by bringing the mercury to the metallic state. The meconic acid, particularly when combined with an alkaline base, affords, with corrosive sublimate, an insoluble and inert mercurate of mercury. The meconates of the ordinary alkalies, however, can scarcely be obtained in sufficient quantity to be applied to such a purpose; and from the smallness of the saturating power of meconic acid, it would be hazardous to administer the quantity of opium which would be adequate to the decomposition of a few grains of corrosive sublimate, inasmuch as the muriate of morphia, which would be contemporaneously formed, is a narcotic medicine of very considerable energy. [The hydrated proto-sulphuret of iron was proposed by Mialhe, and confirmed by Orfila, who says that, if it be taken *immediately* after the ingestion of the poison, it completely destroys the poisonous quality of corrosive sublimate.]

MINERAL ACIDS.

The first, because the most ready method of counteracting the action of the mineral acids, is the copious use of water, warm or cold, milk, oil, or mucilage, which, by simply diluting them, renders them less acrid, and at the same time assists the stomach in rejecting them. This course should be followed until the proper antidotes are prepared. These are the alkalies, alkaline earths and soaps; common pearlash or carbonate of potash; carbonate of magnesia; chalk; and soap in solution. Calcined magnesia, though advised by some authors, is objectionable, inasmuch as the rapidity with which it would combine with acid in the stomach, might alone produce inflammation of that organ.

The smallest quantity of nitric acid which is known to have destroyed life is two drams; the smallest fatal dose of muriatic acid, was one ounce.

NARCOTICS.

The remedies employed in all cases of poisoning from this class of substances, are nearly the same, viz. the most active emetics, as sulphate of zinc in large doses, alum and ipecacuanha. In all cases where it is possible, the stomach-tube should be employed. After these evacuations, strong decoctions of coffee or the vegetable astringents, with cold affusion, are proper. Electro-magnetism is also serviceable, together with powerful stimulants.

NITRE.

This salt is often taken in mistake for Epsom or Glauber's salts. When this happens, vomiting should be excited as speedily as possible, followed by the free use of mucilaginous drinks, hot gin-toddy, brandy and water, &c.

As there is no antidote to this poison, in case these means should fail we must resort to the stomach-pump.

OPIUM.

We cannot do better in this place than transcribe what Dr. T. R. Beck, of Albany, says of the antidotes to opium.

"The experiments of Orfila have demonstrated that the administration of vinegar, or rather vegetable acids, previous to the evacuation of the poison by vomiting, will accelerate and aggravate the action of the opium; but that when the opium has previously been expelled, water acidulated with vinegar, or any other vegetable acid, will tend to diminish and correct the effects. Coffee, when prepared in strong decoction or infusion, rapidly lessens the symptoms, but cannot be considered as an antidote. The result of his experiments led him to give the following directions for the treatment of a person poisoned by opium. Induce vomiting, if possible, with sulphate of zinc, or sulphate of copper; but the administration of watery liquors with the view of promoting the vomiting, should be forbidden, as they dissolve the opium and promote its absorption. In order to insure the action of the emetic, the patient must be kept *roused* by discharging cold water over the head and breast; yet cold affusion must be avoided in the advanced stage, when the body is cold, and the breathing scarcely perceptible. To insure success, in every case the patient must be kept awake, either by dragging him up and down between two men, or by flagellation on the back and soles of the feet. Artificial respiration may be used in desperate cases with propriety. Bleed the patient immediately after the rejection of the poison, and repeat this if necessary. Afterwards administer alternately water acidulated with any vegetable acid and a strong infusion of coffee warmed—and in ten or twelve hours exhibit an enema, and let the arms and legs of the patient be well rubbed. Purgative clysters should be continued, if it be suspected that any of the opium still remains in the large intestines."—Beck, *Med. Jurisprudence*, vol. ii. p. 403. These remarks are applicable to the poisonous effects induced by other articles of the Narcotic class, as belladonna, hyoscyamus, stramonium, &c. Yet as

these are not often taken with suicidal intentions, the quantity is seldom inordinately large. Emetics, if given soon after the poison has been swallowed, will at least remove a part of it; after which the practitioner must necessarily be governed in the treatment by the nature of the peculiar poison, the amount taken and the patient's constitution. Stimulants are occasionally necessary from the first; in other instances, depletion; but more frequently, each is in turn called for. S. G. M.

OXALIC ACID.

Death from this poison is generally sudden. Lime-water or a mixture of chalk and water is recommended by Dr. Thompson, for the purpose of converting the acid into oxalate of lime. The carbonate of magnesia, also, has been found by experiment to counteract the poison; but the other alkalies must be avoided, as they act with an energy little inferior to the acid itself.

It is advisable to avoid giving *warm water* with a view to accelerate vomiting; for dilution promotes the absorption of the poison.

PHOSPHORUS.

This substance acts so immediately and so powerfully in decomposing the animal tissues, that no available antidote has hitherto been suggested. A small portion is sufficient to destroy life. Nevertheless, large quantities of mild demulcent liquids should be exhibited to shield the alimentary canal, and magnesia to neutralize the phosphoric acid.

PRUSSIC ACID.

This deadly poison, in its purity, acts with such rapidity as to leave no time for the administration of antidotes. When taken in small quantities, much diluted, its action is less rapid, producing symptoms of excessive prostration, convulsions and coma. Cold affusion to the head has been found very effectual in such cases, and when applied before the convulsive stage has passed, has often succeeded. The head should be kept elevated, and the water poured from a height upon the occiput and spine. The inhalation of ammonia is the antidote most to be relied on, many experiments having proved its efficacy. It should not be used of too great strength, which would endanger inflammation of the air-passages and stomach; the strong aqua ammonia, diluted with twelve parts of water, is recommended by Orfila.

Chlorine also possesses strong claims as an antidote, and from

the results of his experiments, is believed by Orfila to be the best which has yet been proposed.

Ether and chloroform have been recommended, and no doubt will eventually prove of great value from the rapidity with which they act.

Cases of poisoning with the oil of bitter almonds, cherry laurel water, &c., containing Prussic acid, are not unfrequent, particularly among children, and after evacuating the stomach, require the same treatment as for Prussic acid.

SILVER.

The nitrate of this metal is sometimes taken by mistake; its antidote is common salt, *largely given*, with copious draughts of mucilaginous drinks to promote vomiting. The salt produces an insoluble chloride which has no power on the system.

BECK.

TIN.

When any of the salts of tin are taken, albumen of eggs, flour, or milk given freely, is the appropriate remedy. Milk forms a coagulum with the muriate of tin (*Dyer's spirit*), and renders it innoxious.

ZINC.

The sulphate of this metal is occasionally taken in an overdose. Warm water and emollient drinks are proper in such cases; but milk is particularly useful, from its power of decomposing the sulphate; also albumen, magnesia, &c.

The crystallized sulphate of zinc very closely resembles the sulphate of magnesia, and *has been mistaken* for it; a fact to be borne in mind by apothecaries and physicians.

The following works have been consulted in arranging the preceding remarks on Poisons: A Treatise on Poisons, by Robert Christison, M. D.—Turner's Chemistry.—Manuel de Médecine Légale, par A. Brierre de Boismont.—Orfila, Toxicologie Générale.—Beck and Taylor's Medical Jurisprudences.—Henry, Thompson, &c. &c.—Costill's Treatise on Poisons.¹

¹ A Practical Treatise on Poisons, their Symptoms, Antidotes, and Mode of Treatment. Dr. Costill's work embraces clear and concise views of the subject of which it treats, and constitutes an excellent *manual* for all practical purposes.

APPENDIX NO. I.

ON THE ENDERMIC USE OF MEDICINES.

"THE science of medicine," says Dr. Gerhard,¹ "has recently received an important auxiliary to its resources, by the discovery that remedial substances when applied to the cutis, after the removal of the epidermis, produce their peculiar effects with nearly the same energy as if introduced into the alimentary canal. This mode of exhibition, which has received the name of the *endermic method*, or *endermic medication*, is always found much more efficacious than the iatatreptic method of Christien, or the administration of medicines by means of frictions on the cuticle."

The endermic method requires the previous separation of the cuticle, which is best effected by a blister about four inches square. The cuticle should then be punctured or removed in very small patches, so that the medicine may come in contact with the vesicated surface. Substances used in this way must be applied in two or three times the quantity requisite when given by the mouth. "The remains of the substance applied are much less than the original quantity, and sometimes no trace of it can be perceived; the whole appearing to be absorbed or mingled with the secretions."

We avail ourselves of Dr. Gerhard's valuable paper for the following facts in relation to those medicines which act most effectually in the endermic method.

Sulphate of Quinia.—A blister to be applied to the epigastrium during the apyrexia; and if the paroxysm returns before the quinia can be used, defer it until the next apyrexia. Apply four grains of powdered quinia (either alone, or with an equal quantity of starch), four times a day, for two successive days. A less irritating plan, however, is to dissolve sixteen grains of quinia in two ounces of water, with which the blis-

¹ North Amer. Med. and Surg. Journ. April, 1830.

tered surface is to be repeatedly wet during the day by means of a soft linen cloth. There is no occasion to *remove* the loose epidermis; it is all-sufficient to *puncture* it in several places to allow the solution to touch the denuded surface.

Morphia.—The acetate and sulphate act readily and effectually; the latter is preferable, because more soluble and less irritating. It may be applied in powder, in solution, or in the form of cerate. The dose varies from half a grain to two grains, and its effects are speedily perceptible.

Digitalis.—Dose, two grains three times a day.

Aloes.—This medicine acts on the bowels with nearly the same certainty as if taken internally. Dose, ten grains, to be repeated, if necessary, in a few hours.

Gamboge, owing to its solubility, purges freely in the endermic method. Dose, from five to eight grains.

Rhubarb, *jalap*, *elaterium* and *croton oil*, contrary to what might be expected, do not produce catharsis, although they excite severe local irritation.

Squills.—This medicine in powder, sprinkled on a blistered surface three times a day, in the quantity of four grains at each application, causes copious diuresis. Its power is increased by combination with *digitalis*, in the proportion of four grains of the former to one of the latter.

Emetics from the *vegetable* kingdom all produce their effect in the endermic method; and tobacco does so without any previous irritation of the skin. The same remark is applicable to mercury and iodine, which last is best applied in the form of ointment, made by rubbing a dram with an ounce of cerate.

Emetics derived from the mineral kingdom exert no specific action.

APPENDIX NO. II.

ON THE USE OF ETHER AND CHLOROFORM.

THE employment of anæsthetic agents has now become so general, and their great value so fully attested by many eminent physicians and surgeons in all parts of the world, that we shall here give a brief account of their uses and mode of application.

Indeed, the discovery of a means by which to give complete insensibility to pain during the performance of tedious surgical and dental operations, as well as to mitigate the spontaneous sufferings attendant on many diseases, must be considered one of the most important in the annals of medicine. That pain is useful in indicating the seat and progress of disease, cannot be denied; yet, on the other hand, its severity may exhaust life; and to allay it is often the most difficult and necessary part the physician is called upon to perform. The inhalation of ethereal substances now gives the practitioner more or less control over these "ills that flesh is heir to," enabling him to alleviate some, and wholly to prevent others.

Notwithstanding the benefit to be derived from its judicious use, this mode of inhalation is also capable of producing much evil in the hands of the unskilful and ignorant; and it should, therefore, be employed by those only who are acquainted with its diversified operation.

Ether and chloroform produce essentially the same effects, but the latter is much the more powerful and rapid in its action, and has the disadvantage of being a preparation of unequal strength.

The progressive stages of etherization seem to be pretty much as stated by Mr. Sibson, viz:—

1. Cerebral excitement.
2. Cerebral derangement, staggering intoxication.

3. Cerebral sleep (sopor); pupils contracted; dreams; reflex functions of the brain and spinal marrow still active.

4. Cerebral death (coma); reflex functions of the medulla and spinal marrow still active.

5. Death of spinal marrow; cessation of respiration (heart's action generally still present).

6. Cessation of heart's action.

He also says: "It is very important to be able to tell easily when the stage of safety—sopor, is about to merge into that of danger—coma. The action of the pupils is the key to this knowledge. Inhalation should not be continued one instant after the pupils, previously contracted, have begun to dilate.

"If unconsciousness can be secured by sopor, the inhalation should not be carried on to produce coma. If complete muscular relaxation be sought for, as in hernia, to facilitate taxis; in dislocation, to make reduction easy; and in tetanus, then it will be needful, in general, to urge the patient from sopor into coma; but as soon as the muscular relaxation is secured, the inhalation should cease. When chloroform or ether is employed in chorea, delirium tremens, or other affections, it should never be urged beyond sopor. In neuralgia, it is not usually needful to produce unconsciousness. Whenever the pain disappears the inhalation ought to cease."

According to Dr. Snow, a necessary precaution for watching the effects of ether and chloroform is, that they possess an accumulative property; their influence sometimes increasing for twenty seconds after the inhalation is discontinued. It is seldom necessary to carry the process farther than the production of sopor; in this dreamy state insensibility to pain is perfect, and in some persons much earlier. Many patients seem to be perfectly awake to all mental impressions, and even take an interest in the progress of the operation, without feeling the slightest pain, and, indeed, in some instances, experiencing agreeable sensations.

The influence of these agents being transmitted by the blood, pervades the whole system simultaneously, and particularly excites the capillary circulation as evinced by the injected conjunctiva, increased lachrymal and salivary secretions, and turgidity of the face and neck, the redness of which, however, soon gives place to pallor.

Although used heretofore only to produce general effects, it has lately been discovered by Mr. T. Nunneley, of Leeds, that local anæsthesia may be caused by these agents; and that the brain and other parts of the body not subjected to their influence remain unaffected. After an extensive series of experiments on different animals, he finds that, by immersion in a

small quantity of the fluid, or by applying the vapor locally for a limited period, a limb may be rendered perfectly motionless and senseless, and, what may be an additional advantage, fixed in any desired position. The benefits likely to be derived from this new and valuable discovery promise to be very great.

As regards the mode of applying these vapors, the principal thing to be observed is, that they be freely diluted with atmospheric air. Many instruments for inhalation have been invented, but most of them have given place to a simple sponge; and in the case of chloroform, a towel or handkerchief is perhaps as good as anything yet proposed. The degree of etherization does not depend so much upon the amount respired, as upon the peculiar constitution of the individual, and the physician is thus better able to regulate the amount, and judge of its effect. A dram of *chloroform*, or even half a dram, poured upon a handkerchief, and gradually brought close to the mouth and nose, will, in about two minutes, more or less, produce the required effect. Of *ether*, half an ounce may be poured upon a sponge, and more added from time to time if required. The full effect of ether is not generally obtained in less than four or five minutes. As regards their comparative merits, it may be said of chloroform that it is less pungent, and requiring a less amount is more easily inhaled; but it is also said to have a greater tendency to produce involuntary muscular contraction, and exerts a more direct and powerful action on the heart than ether. As far as present experience goes, ether has been found the milder and safer of the two; and in the case of children, from their greater susceptibility, should always be given in preference to chloroform; etherization, in them, occurring in about two minutes.

According to Dr. Snow, the immunity from pain does not correspond exactly with the degree of narcotism, but is greater as the patient is recovering from the effects of the vapor, than in the corresponding degree, as he is getting under its influence; this is more particularly the case with ether, which he considers to possess superior anæsthetic effects to chloroform in proportion to the narcotism produced. When ether is inhaled, the patient sometimes completely recovers his mental faculties, although insensibility to the operation still continues.

The use of ether or chloroform is contra-indicated by the existence of any disease of the heart, tendency to engorgement of the lungs, or aneurism of any of the great vessels; nor should it ever be used either during the existence of shock resulting from a severe injury.

The existence of extreme debility is not always a contra-indication; for, acting as a stimulant, the inhaled vapor seems to

possess a sustaining power, and has been successfully used in a state of great collapse, as well as for its sedative influence in low cases of typhus fever.

Perhaps one of the most important uses of ether and chloroform is that for the relief of stricture, and to render easy the introduction of the catheter; here in old and sensitive cases as well as in retention of urine and the passage of calculi, they are invaluable.

The physician in using these vapors should bear in mind that he is administering an agent of great power, of which he must carefully observe the effect; for the life of the patient is in his hands. In some constitutions six or seven inspirations of chloroform have produced complete insensibility, and we should always be on our guard lest alarming symptoms suddenly supervene. These are marked by suspended respiration; livid skin; widely dilated pupils; the eyes fixed, and the iris immovable. The patient should be immediately laid upon the floor, cold water dashed upon the face, and the vapor of ammonia applied to the nostrils. These will generally restore animation; if, however, respiration has entirely ceased, it will be necessary to bring it about by artificial means. The abstraction of a few ounces of blood from the arm, or from the jugular vein, will also assist the renewal of the heart's action, by relieving distension.

We should be careful, also, not to endeavor to make the patient swallow anything while in this state of insensibility, as cough might induce suffocation and hasten death.

Besides sulphuric ether and chloroform, various other preparations have been found to possess anæsthetic properties; as Aldehyde, Benzoin, Acetic Ether, Bisulphuret of Carbon, &c., but sufficient experiments have not yet been made to determine their relative value.

As the action of chloroform depends upon its comparative purity, this may be tested by dropping it into the following mixture, viz. Sulphuric acid at 66° and distilled water, equal parts. When cool, this mixture will have a density by the areometer of 40°. (Sp. gr. 1.38.) Pure chloroform dropped into this liquid will fall to the bottom.¹ S. G. MORTON.

¹ French Journal of Pharmacy, 1848.

INDEX.

- A.
- Absorbents and antacids, 198
- Acacia, mucilage of, 262
- Acetate of ammonia, lotion of, 242
 and potash, 73
 and assafetida, 102
 and camphor, 226
 and opium, 226
- of lead, gargle of, 235
 solution of, 197
 enema of, 197
 and calomel, 190, 192
 and ipecacuanha, 191
 and opium, 192
- of morphia and tolu, 107
- of potash and ammonia, 73
 and buchu, 73
 digitalis, &c., 71
- of strychnia, solution of, 175, 186
 tincture of, 186
- of zinc, solution of, 231
 collyrium of, 224
- Acetic acid, gargle of, 235
- Acid nitrate of mercury, 207
- Aconite and Dover's powder, 137
 and wine of colchicum, 85
 and valerian, 86
- Acorn coffee, 172
- Alcohol as a poison, 269
- Alkalies as poisons, 270
- Alkaline bath, 256
- Aloes, decoction of, 59
 enema of, 123
 pills of, 51
 tincture of, 61
 compound tincture of, 116
 and assafetida, 51
 and canella, 110
 and gentian, 51, 174
 and guaiacum, 52
 and myrrh, 51
 myrrh, and rhubarb, 51
 sulphate of iron, &c., 112
- Almond emulsion, 99
- Alteratives, 177
- Alum, burnt, 206
 collyrium of, 223
 curd, 224
 gargle of, 234
 powder of, 37
 compound powder of, 190
 whey, 196
 and calomel ointment, 248
 and catechu, 192
 and extract of bark, 193, 196
 and opium, 191
 sage tea, and honey, 237
 and sulphate of zinc, 39
 and sweet spirits of nitre, 234
- Amber, cloves, &c., 150
 salt of, and castor, 147
- American hellebore, tincture of, 39
- Ammonia, liquor, injection of, 114
 lotion of, 217
 and tartar emetic, 241
- Ammonia, fetid spirits of, 128
 aromatic spirits of, 114, 128
 and olive oil, 122
 and glycerin, 243
 and magnesia, 200
 and salt of tartar, 200
 and valerian, 149
- Ammoniated iron, 166
- Ammoniuret of copper, 166
 injection of, 233
- Ammoniac, mixture of, 102, 106
 and nitric acid, 103
 seneka, &c., 99
 squill, &c., 94
- Angustura bark, powder of, 165
 decoction of, 171
- Animal oil, 129
- Anodyne collyrium, 227
 draught, 137
- Antacids, 198
- Antacid tincture, 189
- Anthelmintics, 117
- Anti-bilious pills, 50
- Anti-emetic mixture, 201
- Anti-gout pill, 48
- Anti-herpetic ointment, 248

- Anti-intermittents, 155
 Antilithics, 76
 Antimonial powder, 80
 wine, 39
 and spirits of Min-
 dereri, 83
 and sweet spirits of
 nitre, 84
 and tolu, 85
 Antimony, sulphuret of, 81
 as a poison, 270
 Antiseptic cataplasm, 254
 pills, 125
 Antispasmodics, 147
 Antisyphilitic pills, 151
 Aperient pills, 51
 Apocynum, decoction of, 70
 Aqua sappharina, 228
 Armenian bole, 232
 Arnica, mixture of, 86
 tincture of, 129
 Aromatic waters, mode of preparing, 243
 pills, 125
 Arrowroot, 265
 Arsenic as an alterative, 179
 as a poison, 270
 Arsenical mixture, 163
 Arsenious acid, 158
 solution of, 207
 ointment of, 208
 compound powder of,
 208
 and sal ammoniac, 158
 Arseniate of iron, pills of, 179
 of soda, solution of, 185
 Arsenite of potash, solution of, 163
 Asiatic pills, 179
 Assafetida, enema of, 150
 mixture of, 101, 149
 pills of, 147
 sirup of, 97
 and aloes, 51
 and acetate of ammonia, 102
 and camphor, 136
 castor, &c., 149
 magnesia, &c., 141
 Astringents, 190
 Astringent fomentation, 239
 Atropia, solution of, 143

 B.
 Balsam of Peru, mixture of, 103
 and ox gall, 229
 Bareges bath, 256
 Barley water, 261
 Bates's camphorated water, 226
 alum water, 228
 Baths, 255
 Baryta as a poison, 271
 Bebeerine, sulphate of, 159
 Beef tea, 266
 Belladonna, ointment of, 144, 248
 solution of, 141, 143
 cherry laurel, &c., 145
 quinia, and rhubarb, 165
 and tolu, 104
 Bicarbonate of soda, mixture of, 77, 101
 and calomel, 44
 magnesia, &c., 199
 quassia, &c., 77
 and uva ursi, 65
 of potash, solution of, 77
 Bitartrate of potash. See Cream of
 Tartar
 Black hellebore, tincture of, 115
 myrrh, &c., 115
 Black drop, 138
 wash, 240
 Blistering cerate, 212
 Blister issue, 210
 Bloodroot, tincture of, 101
 Blue mass and aloes, 47
 and colocynth, 48
 and extract of dandelion, 67
 digitalis and squill, 67
 and magnesia, 56
 and quinia, 178
 and soda, 48
 Boneset, infusion of, 86, 160
 sage, &c., 170
 Borax, gargle of, 234
 and glycerin, 243
 and morphia, 244
 and savine, 111
 Brandy, gargle of, 233
 mixture, 131
 Bran, decoction of, 262
 and flaxseed poultice, 253
 Bread and milk poultice, 253
 Bromide of potassium, solution of, 185
 ointment, 249
 Bromide of iron, pills of, 180
 Bromine, 184
 solution of, 185
 Brown mixture, 98
 Brucia, 180
 mixture of, 186
 pills of, 181
 tincture of, 186
 Buchu and acetate of potash, 73
 and iodide of potassium, 91
 and uva ursi, 74
 Burdock, decoction of, 88
 Burnt alum, 206
 sponge, 181
 Butternut and jalap, 49

 C.
 Cainca, decoction of, 69
 Calf's-foot jelly, 267

- Calomel, pills of, 151, 177
 injection of, 232
 and acetate of lead, 190, 192
 and alum ointment, 248
 and camphor, 153
 and colocynth, 48
 and conium, 136
 digitalis, &c., 67
 digitalis and opium, 66
 and gamboge, 117
 Goulard's cerate, &c., 249
 and iodine ointment, 246
 and ipecacuanha, 38, 191
 and jalap, 44
 jalap, and rhubarb, 44
 opium, and ipecacuanha, 81
 and ox gall, 180
 and pink-root, 117
 quinia, and opium, 178
 and rhubarb, 44
 semen contra, &c., 121
 and soda, 44
 and squill, 66, 94
 sulphuret of antimony, &c., 81
 and tar ointment, 251
 turpentine, &c., 119
 and verdigris, 252
- Camphor, enema of, 123, 128
 in milk, 127
 mixture of, 114, 127, 194
 pills of, 135
 Julep, 127
 assafetida, &c., 136
 and acetate of ammonia, 226
 and calomel, 153
 and cantharides, 124
 and carbonate of ammonia, 241
 and chloroform, 128, 146
 and cinchona, 166
 and Hoffman's anodyne, 140
 injection for ear, 228
 laudanum, and nitric acid, 194
 liniment, 241
 magnesia, &c., 200
 and mercurial ointment, 247, 216
 musk, &c., 147, 148
 and myrrh, 127
 and opium, collyrium of, 226
 and opium, plaster of, 145
- Camphorated oil, 241
- Canella, aloes, &c., 110
- Canquoin's paste, 207
- Cantharides, extract of, 214
 liniment of, 220, 241
 mixture of, 71
 tincture of, 116
 camphor, &c., 124
- Cantharides, hartshorn, &c., 114
 and savine, 111
 and oil of turpentine, 220
- Cantharidin, 213
- Carbonate of magnesia, mixture of, 56, 78
 of potash, mixture of, 100, 199
 ammonia, &c., 200
 and lime-water, 77
 squill, &c., 73
 of soda, injection of, 78
 mixture of, 201
 pills of, 77
 and calomel, 82
 rhubarb, &c., 199
 of ammonia, mixture of, 126
 ointment of, 250
 and musk, 148
 and sulphate of quinia, 157
 and valerian, 126
 of iron, pills of, 166
- Carrageen, 265
- Castor, assafetida, &c., 149
 and salt of amber, 147
- Cataplasms, 253
- Catechu, injection of, 229
 electuary of, 196
 mixture of, 196
 and alum, 192
- Cathartics, 43
- Caustics, 205
- Caustic potash, see Potash.
 paste of Canquoin, 207
- Cayenne pepper, gargle of, 236
 infusion of, 130
 pills of, 125
 tincture of, 219
 iron and aloes, 167
- Chalk mixture, 194, 199
 and galls, 195
 and opium, 198
- Chamomile and orange peel, 171
- Charcoal, poultice of, 254
 and creasote ointment, 251
- Cherry laurel, belladonna, &c., 145
- Chicken water, 267
- Chlorine fumigation, 109
 liniment, 240
- Chloroform, as an anæsthetic, 280
 and camphor, 128, 146
- Chlorinated lime, 130
 gargle of, 236
 mouth wash, 236
- Chloride of iron, tincture of, 174
 of gold, 169

- Chloride of zinc, as caustic, 207
 injection of, 231
 lotion of, 238
 solution of, 150
- Chocolate, 266
- Chromic acid, 208
- Cider mixture, 70
- Cinchona, enema of, 164
 extract of, 159
 decoction of, 159, 160
 infusion of, 160
 mixture of, 163
 powder of, 155
 and alum, 193, 196
 and camphor, 166
 and citrate of potash, 162
 and cloves, 156
 and Epsom salts, 156
 and lime-water, 160, 196
 and port wine, 163
 and rust of iron, 169
 sal ammoniac, &c., 166
 and serpentaria, 156
 uva ursi, and opium, 76
 and valerian, 172, 173
- Cinchonia, sulphate of, 157, 162
- Citrate of magnesia, 57
 of potash, solution of, 82
 and cinchona, 162
 of ammonia, solution of, 84
- Citrine ointment, 251
- Cloves, infusion of, 130
 plaster of, 216
 and bark, 156
- Cochineal, mixture of, 101
 and common salt, 118
- Cod-liver oil, 186
 mixture of, 187
 and iodine, 187
- Codeia, 139
- Colchicum, wine of root, 60
 and aconite, 85
 and colocynth, 48
 and digitalis, 73
 and elaterium, 72
 and magnesia, 57
 squill, and tobacco, 72
- Collodium vesicans, 213
- Collyria, 222
- Colocynth and blue mass, 48
 and calomel, 48
 and colchicum, 48
 and croton oil, 50
 jalap, &c, 49
 and manna, enema of, 62
- Colombo, mixture of, 200
 tincture of, 174
 ginger, &c., 164, 170
 magnesia, &c., 199
 rhubarb, &c., 170
 and tartrate of iron, 165
- Confection of senna, 52
 of sulphur, 53
 of sulphur and guaiac, 53
- Conia, collyrium of, 227
- Conium, infusion of, 144
 extract of, 135
 extract of, and calomel, 186
 mixture of, 99, 140
 suppository of, 143
 and corrosive sublimate, 152
 and iodine inhalation, 108
 ipecacuanha, &c., 93
- Conradi's collyrium, 226
- Copaiba, mixture of, 74
 compound mixture of, 75
 pills of, 67
 cubeb, &c., 75
 and guaiacum, 115
 and tolu, 102
- Copper, sulphate of, 38
 as a poison, 271
- Corrosive sublimate, bath of, 256
 collyrium of, 225
 gargle of, 235
 injection of, 231
 lotion of, 239
 mixture of, 188
 pills of, 151, 179
 solution of, 91
 tincture of, 189
 and hemlock, 152
 and tartar emetic, 217
- Cowhage electuary, 121
 ointment, 221
- Coxe's hive sirup, 41, 97
- Crab's claws and rhubarb, 198
- Cream of tartar and juniper berries, 68
 and jalap, 45
 and squill, 66
 and sulphur, 46
- Creasote, mixture of, 196
 injection of, 229
 liniment of, 240
 lotion of, 240
 pills of, 125, 192
 solution of, 196
 and charcoal ointment, 251
- Croton oil, liniment of, 220
 mixture of, 55
 pills of, 50
 saponaceous solution of, 55
 tincture of, 60
 and colocynth, 50
 and potash, 221
- Cubeb, fluid extract of, 75
 lozenges of, 96
 mixture of, 130
 and bismuth, 130
 copaiba, &c., 75
 and ergot, 66

Cuprum ammoniatum, 166
 Cyanuret of mercury, ointment of, 247
 solution of, 153
 of potassium, 95
 lotion of, 240
 mixture of, 106

D.

Dandelion, infusion of, 69
 and blue mass, 67
 Depilatory ointment, 252
 Dewees's carminative, 141
 tincture, 115
 Diaphoretics, 79
 alterative, 87
 Dietetic preparations, 261
 Digitalis, infusion of, 68
 tincture of, 101, 140
 acetate of potash, &c., 71
 calomel, &c., 67
 calomel and opium, 66
 and colchicum, 73
 and myrrh, 112
 squill and blue mass, 67
 and opium, 136
 squill and juniper, 67
 and tolu, 102
 Digitalin, pills of, 137
 solution of, 70
 Dippel's animal oil, 129
 Disinfecting mouth wash, 236
 Diuretics, 65
 Diuretic drops, 74
 Dolichos, see Cowhage.
 Donovan's solution, 184
 Dogwood bark, decoction of, 172
 Dover's powder, 80, 92
 and aconite, 137
 Dubois's arsenical powder, 208
 Dupuytren's lotion, 242
 Dyspeptic lye, 201

E.

Effervescing draught, 83
 Elaterium, pills of, 50
 and colchicum, 72
 Elaterin, tincture of, 60
 Electuary of senna and figs, 53
 of jalap and saltpetre, 53
 Elixir of vitriol and sulphate of iron, 173
 Emetics, 35
 Emetia, lozenges of, 42, 95
 solution of, 40
 Emmenagogues, 110
 Emmenagogue suppository, 116
 Emollient bath, 256
 cataplasm, 254
 collyrium, 226
 Endermic use of medicines, 278
 Enemata, 61, 123, 142, 150, 164

Enema, common, 62
 aloetic, 123
 of camphor and olive oil, 123
 ox gall, 62
 soap, 62
 sulphate of magnesia, 62
 turpentine, 63
 Epispastics, 211
 Epispastic ointment, 212
 Epsom salts, see Sulphate of Magnesia.
 Ergot, infusion of, 113
 powder of, 111
 sirup of, 114
 tincture of, 114
 and cubebs, 66
 Erigeron, infusion of, 68
 Escharotics, 205
 Essence of beef, 266
 Ether, oil of turpentine, &c., 122
 and laudanum, 150
 ioduretted, 184
 phosphorated, 129
 as an anæsthetic, 280
 cantharidalis, 213
 Eupatorium, see Boneset.
 Expectorants, 92

F.

Febrifuge bolus, 166
 Fennel-seed, magnesia, &c., 199
 Fern, oil of, 119
 Ferriar's hydragogue draught, 72
 Fetid spirits of ammonia, 128
 Figs and senna, electuary of, 53
 Firing, 205
 Flaxseed, infusion of, 104
 and bran poultice, 253
 Flour, boiled, 265
 Fluid extract of cubebs, 75
 of spigelia and senna, 122
 of senna, 58
 of rhubarb, 58
 Fomentations, 257
 Fothergill's pills, 49
 Fowler's solution, 163
 and iodine, 185
 Fumigations, 258
 Fumigating powders, 258

G.

Gall, see Ox gall.
 Galls, gargle of, 233
 and brandy, 197
 infusion of, 195
 ointment of, 248
 and chalk, 195
 Gallic acid, pills of, 193
 Gamboge and calomel, 117
 and jalap, 45
 Gargles, 233

- Garlic, cataplasm of, 215
 sirup of, 96
- Gentian, compound infusion of, 171
 tincture of, 173
 acidulated tincture of, 174
 and aloes, 51, 174
 and iron, 168
- Ginger, essences of, 132
 infusion of, 132
 carbonate of ammonia, &c., 126
 colombo, &c., 164, 170
- Glycerin, 243
 and ammonia, 243
 and borax, 243
- Gold, see Nitromuriate of Gold.
 as a poison, 272
- Gondret's pommade, 219
- Goulard's extract, lotion of, 238
 and rose water, 228
 cerate and calomel, 249
- Granville's lotion, 217
- Griffith's myrrh mixture, 172
 pills, 47
- Guaiacum, mixture of, 85
 tincture of, 115
 and aloes, 52
 and copaiba, 115
 nitre, &c., 82
 and paregoric, 128
 and prussiate of iron, 156
 and sulphur, 53
 and turpentine, 124
- Gum-Arabic mucilage, 262
- Guthrie's ointment, 227
- H.
- Harrowgate bath, 257
 water, artificial, 189
- Hartshorn jelly, 266
- Hemp, tincture of, 141
- Hemlock, see Conium.
- Henbane, extract of, 135
 infusion of, 144
 and ipecacuanha, 134
 opium, &c., 135
- Hiera picra, 110
 tincture of, 116
- Hoffman's anodyne and camphor, 140
 and laudanum, 149
 and valerian, 149
- Hops, infusion of, 171
 tincture of, 140
 and tar, infusion of, 107
- Hope's mixture, 194
- Hooper's pills, 112
- Hufeland's diuretic drops, 74
- Huxham's tincture, 174
- Hydrocyanic acid, 105
 as a poison, 276
- Hydrocyanic acid, lotion of, 239
 mixture of, 105
 and lobelia, 106
- Hydrocyanate of potash, 106
- Hydriodate of potash, 183
- Hyposulphite of soda, lotion of, 244
- Hyosciamus, see Henbane.
- I.
- Iceland moss, decoction of, 104
- Indian hemp, decoction of, 70
- Inhalation, 107
- Injections, 228
 for the ear, 228
 for the urethra, 230
- Instantaneous vesication, 213
- Iodide of arsenic, ointment of, 246
 pills of, 179
 of barium, ointment of, 245
 of iron, injection of, 232
 ointment of, 246
 sirup of, 176
 solution of, 183
 and manganese, 176
 of manganese, pills of, 168
 of mercury, ointment of, 246
 pills of, 152, 153
 compound pills of, 152
 (red) of mercury, ointment of, 246, 247
 pills of, 179
 of potassium, collyrium of, 224
 mixture of, 182, 183
 ointment of, 245
 solution of, 113, 183
 and buchu, 91
 and opium, ointment of, 247
 as a poison, 272
 of silver, pills of, 180
 of sulphur, ointment of, 246
 of zinc, ointment of, 245
 solution of, 236
- Iodine, cataplasm of, 255
 compound tincture of, 182
 ethereal solution of, 221
 ethereal tincture of, 182
 inhalation, 108
 liniment of, 240
 ointment, 245
 paint, 221
 preparations of, 181
 tincture of, 115, 182, 183
 waters, 184
 as a poison, 272
 and arsenic, 185
 and calomel ointment, 246
 and cod-liver oil, 187
 and hemlock inhalation, 108

- Iodine, and iodide of potassium, 182
and opium gargle, 236
Iodo-hydrargyrate of potash, 184
Ioduretted baths, 257
ether, 184
Ipecacuanha, infusion of, 40
ointment of, 251
powder of, 36
sirup of, 41
and calomel, 38
calomel and opium, 81
conium, &c., 93
and henbane, 134
iron, &c., 165
and mercury with chalk, 154
myrrh, &c., 92
and opium, 80, 92
and rhubarb, 38, 253
squill and seneka, 97
and tartar emetic, 36, 38, 40
Irish moss, 265
Iron, pills of, 112
powder of, 165
rust of, 165
and gentian, 168
ipecacuanha, &c., 165
and muriate of baryta, 90
muriated tincture of, 116
protocarbonate of, 166
protonitrate of, 175
persesquinitrate of, 175
subcarbonate of, 119, 164
and valerian, 148
Issues and setons, 209
J.
Jackson's lozenges, 96
mixture, 195
pectoral sirup, 98
Jalap and calomel, 44
calomel and rhubarb, 44
and cream of tartar, 45
and extract of butternut, 49
and gamboge, 45
and sulphate of potash, 58
Juniper berries and squill, 71
and cream of tartar, 68
oil of, mixture, 71
squill and digitalis, 67
K.
Kentish's ointment, 249
Kino, opium, &c., 193
Kirkland's neutral cerate, 250
Kousso, infusion of, 120
Krameria, mixture of, 197
pills of, 192
suppository of, 143
L.
Labarraque's solution, 244
Lactate of iron, 166
Lactucarium, pills of, 134
Laffeteur's sirup, 88
Lapis divinus, 223
Laudanum, 137
as a poison, 272
Sydenham's, 138
enema of, 142
liniment, 242
compound liniment, 242
and ether, 150
and Hoffman's anodyne, 149
and iodine gargle, 236
and oil of turpentine, 139
and sweet spirits of nitre, 84
and tartar emetic, 100
Laxative suppository, 64
Lead as a poison, 273
Lemonade, 263
Lenitive electuary, 52
Liniments, 237
Linseed oil and lime-water, 238
Lime-water and alcohol, 238
and carbonate of potash, 77
and cinchona, 160, 196
and linseed oil, 238
and milk, 200
and opium, 144
and zinc, 237
Lime and zinc ointment, 251
Lithontriptics, 76
Lobelia, infusion of, 40
tincture of, 39
and prussic acid, 106
Logwood, extract of, 195
infusion of, 195
Lotions, 237
Lugol's ointment, 245
solution, 183
Lunar caustic, 206
Lupulin, pills of, 134
tincture of, 140
M.
Madder, decoction of, 113
powder of, 111
Magendie's solution, 183
Magnesia, calcined, 45
ammonia, &c., 200
assafetida, &c., 141
and blue pill, 56
and colchicum, 57
and colombo, 199
camphor, &c., 200
fennel-seed, &c., 199
and rhubarb, 45, 56
soda, &c., 199
and sulphur, 46

- Male fern, extract of, 119
 powder of, 118
 Malt, infusion of, 262
 May apple and podophyllin, 48
 Manganese and iron, iodide of, 176
 Manna and colocynth, enema of, 62
 and senna, 59
 Matico, infusion of, 194
 Mel ægyptiacum, 236
 Mercury as an alterative, 177
 as a poison, 273
 and chalk, with ipecacuanha, 154
 chloride of. See Calomel.
 cyanuret of, 153
 red oxide of, 153
 yellow sulphate of, 37
 red sulphuret of, 87
 Mercurial cerate, 248
 ointment, 247
 and camphor, 216, 247
 Mezereon, decoction of, 89
 Milk for infants, 263
 and lime-water, 200
 Mindereri, spirits of, 83
 Mineral acids as caustics, 207
 as poisons, 274
 Mitchell's pills, 49, 167
 Monesia, 193
 Mouth washes, 233
 Moxa, 209
 Muriate of ammonia, solution of, 208, 231
 and arsenic, 158
 of baryta and iron, 90
 Muriated tincture of iron, 174
 Muriatic acid, gargle of, 235
 linctus of, 237
 mixture of, 78
 Mush poultice, 253
 Musk, enema of, 150
 mixture of, 148
 camphor, &c., 147, 148
 and carbonate of ammonia, 148
 Mustard, 169
 fomentation, 258
 powder of, 37
 whey, 131
 plaster, 215
 pepper, &c., 215
 Myrrh, lotion of, 242
 mixture, 172
 black hellebore, &c., 115
 camphor, &c., 127
 digitalis, &c., 112
 ipecacuanha, &c., 92
 squill, &c., 95
 sulphate of iron, &c., 111
 and sulphate of zinc, 93
- N.
- Naphthaline, mixture of, 100
 Narcotics, 133
 external use of, 143
 as poisons, 274
 Narcotic fomentation, 257, 258
 Neutral mixture, 82
 Nitrate (acid) of mercury, 207
 of potash as a poison, 275
 solution of, 85
 guaiacum, &c., 82
 opium, &c., 81, 93
 and squill, 65
 and tartar emetic, 84
 of silver as a poison, 277
 collyrium of, 225
 mixture of, 174
 ointment of, 227, 230, 252
 pills of, 167
 solution of, 229, 230
 Nitric acid, mixture of, 188
 diluted, 188
 and ammoniac, 103
 camphor, and laudanum, 194
 and Epsom salts, 56
 Nitrous powders, 81
 Nitromuriatic acid, 188
 bath, 256
 Nitromuriate of gold, pills of, 169
 as a poison, 272
 Nux vomica, 136
 extract of, 136
 liniment of, 242
 tincture of extract of, 140
- O.
- Oak-bark, gargle of, 233
 Oil of amber, liniment of, 241
 and cloves, 150
 and tolu, 100
 of bitter almonds, mixture of, 142
 of cloves and amber, 150
 of turpentine, 78, 122
 enema of, 63
 linctus of, 130
 liniment of, 220
 mixture of, 54, 72, 126
 ether, &c., 122
 and laudanum, 139
 and Spanish flies, 220
 of juniper, mixture of, 71
 of valerian and hartshorn, 149
 of wormseed, mixture of, 121
 Ointments, 245
 Oleaginous mixture, 53, 54
 and oil of turpentine, 55

- Olive oil and ammonia, 122
- Opium as a poison, 275
 confection of, 135
 enema of, 142
 mixture of, 137
 pills of, 133
 plaster of, 145
 suppository of, 143
 vinegar of, 138
 wine of, 138
 and acetate of ammonia, 226
 and acetate of lead, 192
 and alum, 191
 bark and uva ursi, 76
 and calomel, collyrium of, 223
 calomel, and digitalis, 66
 and camphor, collyrium of, 224
 and camphor, 145, 226
 and catechu, 196
 and chalk, 198
 and digitalis, 136
 henbane, &c., 135
 and iodide of potassium, 247
 ipecacuanha, &c., 81, 93
 and calomel, 81
 and kino, pills of, 193
 and lime-water, 144
 and pellitory, 154
 and rhatany, suppository of, 143
 and rhubarb ointment, 249
 and sulphate of copper, 159
 and tar ointment, 249
 and tartar emetic, 93
- Orange-peel and chamomile, 171
 and gentian, 173
- Oxalic acid as a poison, 276
- Ox gall, enema of, 62
 pills of, 52
 and calomel, 180
 and Peruvian balsam, 229
- Oxide of silver, ointment of, 251
 pills of, 169
 of zinc, collyrium of, 222
 ointment of, 249
 pills of, 168
 and lime ointment, 251
 and lime water, 237
- P.
- Panado, 264
- Paregoric elixir, 138
- Pareira Brava, decoction of, 70
- Parrish's camphor mixture, 194
- Parsley, infusion of, 68
- Pea issue, 210
- Pearson's solution, 185
- Pellitory, masticatory of, 154
 and opium, 154
- Peristaltic persuaders, 50
- Persesquintrate of iron, 175
- Peruvian bark, see Cinchona.
- Petroleum, liniment of, 243
- Phosphoric acid, solution of, 239
- Phosphorus, as a poison, 276
 emulsion of, 129
- Phosphorated ether, 129
- Phosphate of ammonia, 185
- Pills, antibilious, 50
 antigout, 48
 antiseptic, 125
 aperient, 51
 aromatic, 125
 Asiatic, 179
 blue, 153
 Dupuytren's, 151
 Fothergill's, 49
 Griffiths's, 47
 Hooper's, 112
 Mitchell's tonic, 167
 purgative, 49
 Plummer's, 87
 Rufus's, 51
 soap, 77
- Pink-root, infusion of, 120
 fluid extract of, 122
 and calomel, 117
 and savine, 118
 senna, &c., 120
- Piperine, pills of, 158
- Pipsissewa, decoction of, 69
 infusion of, 86
 spirituous decoction of, 69
- Pleurisy root, infusion of, 87
- Plummer's pills, 87
- Podophyllin, pills of, 48
- Poisons, 269
- Pomegranate, decoction of, 120
 gargle of, 235
- Pommade ammoniacale, 219
- Poppies, collyrium of, 227
- Potash, caustic, 206
 solution of, 219
 and croton oil, 221
 and opium, 206
- Poultices, 253
- Powders, antimonial, 80
 Dover's, 80, 92
 nitrous, 81
 Seidlitz, 46
 soda, 46, 82
 Stevens's, 47
- Protonitrate of iron, 175
- Pyroligneous acid, injection of, 229
 ether, &c., 229
- Prunes and senna, 59
- Prussic acid, see Hydrocyanic Acid.
 as a poison, 276
 pectoral mixture, 105

Prussiate of iron and guaiac, 156
 Purgative suppository, 64
 Purgatives, 43

Q.

Quassia, infusion of, 171
 pills of, 168
 extract of, and tar, 94
 Quinia, ferrocyanate of, 161
 muriate of, belladonna, &c., 165
 sulphate of, gargle of, 234
 enema of, 164
 injection of, 232
 mixture of, 161
 pills of, 156
 sirup of, 161
 solution of, 161
 and blue mass, 178
 and carbonate of ammonia, 157
 opium and calomel, 178
 and sulphate of iron, 167
 and tannic acid, 162
 and tartar emetic, 42
 and tartaric acid, 162
 and Vallet's mass, 167

R.

Red oxide of mercury, collyrium of, 223
 ointment of, 227, 245
 pills of, 153

Rennet whey, 268

Resolvent cataplasm, 254

Rhatany, see Krameria.

Rhubarb, fluid extract of, 58
 suppository of, 64
 aloes and myrrh, 51
 and calomel, 44
 calomel and jalap, 44
 and carbonate of soda, 199
 colombo, &c., 170
 and crabs' claws, 198
 and ipecacuanha, 38, 50, 253
 and magnesia, 45, 56
 and opium ointment, 249
 quinia, &c., 165
 and senna, 59
 and sulphate of iron, 52

Rice gruel, 263

water, 262

Rubefacients, 215

Rufus's pills, 51

S.

Sage tea, 262

alum and honey, 237

boneset, &c., 170

Sago, 264

Salaëtatus, solution of, 77

Sal ammoniac, cinchona, &c., 166
 squills, &c., 100

Salep powders, 264

Salt, common, 37

cataplasm, 255

and cochineal, 118

Saltpetre, see Nitrate of Potash.

Salt of tartar, see Carbonate of Potash.

Salicine, pills of, 158

Santonine, powder of, 118

Sarsaparilla beer, 91

decoction of, 89

Savine, powder of, 111

and borax, 111

and cantharides, 111

and pink-root, 118

and verdigris, 253

Scabious, infusion of, 68

Scudamore's mixture, 57

Seidlitz powders, 46

Semen contra, calomel, &c., 121

Senna, confection of, 52

decoction of, 59

fluid extract of, 58

infusion and tincture of, 58

and figs, electuary of, 53

and manna, 59

pink-root, &c., 120

and rhubarb, 59

and salts, 58, 63

Seneka, decoction of, 107, 113

sirup of, 96

ammoniac, &c., 99

and squill, 97

squill, and tartar emetic, 99

Serpentaria, infusion of, 170

and bark, 156

Setons, 210

Sialagogues, 151

Sirup of Laffecteur, 88

Slippery elm poultice, 254

Soap enema, 62

fomentation of, 257

liniment, 221

pills, 77

Soda powders, 46, 82

Sodium, chloride of, 37

Spermaceti mixture, 104

Spice plaster, 216

Spigelia, see Pink-root.

Spirits of Mindereri, 83

and antimonial

wine, 83

- Spirits of Mindereri, and sweet spirits of nitre, 83
- Spirit of ether, and pyroligneous acid, 229
- Sponge, burnt, 181
- Squill, compound sirup of, 41, 97
 sirup of, 96, 98
 tincture of ipecac, &c., 40
 ammoniac, &c., 94
 and calomel, 66, 94
 carbonate of potash, &c., 73
 colchicum, and tobacco, 72
 and cream of tartar, 66
 digitalis, and blue mass, 67
 digitalis, and juniper, 67
 and juniper, 71
 myrrh, &c., 95
 and nitre, 65
 sal ammoniac, &c., 100
 seneka, and ipecacuanha, 97
 seneka, and tartar emetic, 99
- Starch mucilage, 268
- Stevens's saline powders, 47
- Stimulants, 124
- Stimulant fomentation, 257
- Stramonium, decoction of, 144
 extract of, 135
 solution of, 144
- Strengthening plaster, 216
- Strychnia, 185
 collyrium of, 225
 injection of, 233
 mixture of, 141
 pills of, 136
 tincture of, 141
 acetate of, 175
- Subacetate of lead, collyrium of, 224
 lotion of, 238
- Subcarbonate of iron, powders of, 119, 164
 and cinchona, 169
- Subnitrate of bismuth, pills of, 168
 powders of, 191
 cubebs, &c., 130
- Suet ptisan, 267
- Sulphur, confection of, 53
 ointment of, 248, 250
 and cream of tartar, 46
 and guaiac, 53
 and magnesia, 46
- Sulphate of cadmium, collyrium of, 225
 solution of, 230
 of copper, collyrium of, 223, 225
 injection of, 232
 pills of, 159
 powder of, 38
 solution of, 197
 wash of, 237
 and opium, 159
 as a poison, 271
- Sulphate of iron, collyrium of, 224
 injection of, 232
 pills of, 168
 solution of, 244
 aloes, &c., 112
 and elixir of vitriol, 173
 myrrh, &c., 111
 and quinia, 167
 red pepper, &c., 167
 and rhubarb, 52
 and sulphuric acid, 173
- of magnesia, enema of, 62
 and bark, 156
 and nitric acid, 56
 and senna, 58
 and tartar emetic, 55
- of morphia, pills of, 134
 solution of, 139
 and borax, 244
 and tannin, 193
 tolu, &c., 107
- of potash and jalap, 58
 of quinia, see Quinia.
- of zinc, as a poison, 277
 gargle of, 234
 injection of, 231
 pills of, 94
 powder of, 38
 and alum, 39
 and myrrh, 93
- Sulphuret of antimony, calomel, &c., 81, 87
 (red) of mercury, 87
 of potassium, bath of, 256
 lotion of, 238, 239
- Sulphureo-gelatinous bath, 256
- Sulphuric acid, gargle of, 235
 mixture of, 173, 237
- Suppositories, 63, 143
- Sweet spirits of nitre, &c., 83
 and antimonial wine, 84
 and laudanum, 84
- Sydenham's laudanum, 138
- T.
- Tannic acid, ointment of, 252
 pills of, 193
 solution of, 233, 244
 and morphia, 193
 and quinia, 162
- Tapioca, 264
- Tar beer, 103
 ointment, 249
 pills, 94

- Tar water, 103
 wine of, 103
 and calomel, 231
 gentian and quassia, 94
 and infusion of hops, 107
 and opium, 249
- Tartar emetic, as a poison, 270
 enema of, 63
 plaster of, 216
 ointment of, 216
 solution of, 39, 243
 and ammonia, 241
 and corrosive sublimate, 217
 and ipecacuanha, 36, 38, 40
 and laudanum, 100
 and nitrate of potash, 84
 and opium, 93
 and quinia, 42
 and sulphate of magnesia, 55
 tolu, and morphia, 107
- Tartar whey, 268
- Tartaric acid and quinia, 162
- Tartrate of iron and colombo, 165
- Tea, injection of, 232
- Tin, as a poison, 277
 electuary of, 121
- Tobacco, fomentation of, 258
 infusion of, 41
 poultice of, 42
 colchicum, and squill, 72
- Tolu, inhalation of, 108
 and acetate of morphia, 107
 and belladonna, 104
 and copaiba, 102
 and digitalis, 102
 morphia, and antimony, 107
 and oil of amber, 100
 seneka, &c., 99
 and wine of antimony, 85
- Tonics, 155
- Turpentine, liniment of, 220
 pills of, 67
 calomel, &c., 119
 and guaiacum, 124
- Turpeth mineral, 37
- U.
- Urea, mixture of, 73
- Uva ursi, infusion of, 69
 bark, and opium, 76
 and buchu, 74
 and soda, 65
- V.
- Valerian, infusion of, 149
 and aconite, 86
 and carbonate of ammonia, 126
 and cinchona, 172, 173
 and Hoffman's anodyne, 149
 and rust of iron, 148
- Valerianate of quinia, 125
 of zinc, 148
- Vallet's mass, 166
 and quinia, 167
- Vegetable soup, 266
- Veratria, ointment of, 252
 tincture of, 60
 sulphate of, 57
- Veratrum viride, see American Hellebore.
- Vesication, instantaneous, 213
- Vienna paste, 206
- Verdigris, ointment of, 250
 and calomel, 252
 and savine, 253
- Vinegar draught, 173
 collyrium, 224
 whey, 268
 of opium, 138
- Volatile liniment, 220, 241
- W.
- Water gruel, 263
- Whey, alum, 196
 mustard, 131
 rennet, 268
 tartar, 268
 wine, 131
 vinegar, 268
- White linctus, 99
 precipitate ointment, 247
- Wild cherry bark, infusion of, 170
 sirup of, 97
- Wine mixture, 131
 whey, 131
 and bark, 163
 of antimony, 39
 of opium, 138
- Wistar's cough lozenges, 95
- Worm tea, 120
- Wormseed oil, mixture of, 122
- Y.
- Yeast mixture, 128
 poultice, 254
- Yellow wash, 239
- Z.
- Zittman's decoction, 89
- Zinc, as a poison, 277
 pills of oxide, 168
 powder of sulphate, 38

BLANCHARD & LEA'S MEDICAL AND SURGICAL PUBLICATIONS.

TO THE MEDICAL PROFESSION.

In the present catalogue we have affixed prices to our publications, in obedience to the repeated requests of numerous members of the profession. While books, like all other articles, must necessarily vary somewhat in cost throughout the extended territories of this country, yet our publications will generally be furnished at these rates by booksellers throughout the Union, who can readily and speedily procure any which they may not have on hand.

To accommodate those physicians who have not convenient access to bookstores, or who cannot order through merchants visiting the larger cities, we will forward our works by mail, *free of postage*, on receipt of the printed prices in current funds or postage stamps. As our business is wholesale, and we open accounts with booksellers only, the amount must in every case, without exception, accompany the order, and we can assume no risks of the mail, either on the money or the books; and as we only sell our own publications, we can supply no others. Physicians will, therefore, see the convenience and advantage of making their purchases, whenever practicable, from the nearest bookseller.

We can only add that no exertions are spared to merit a continuance of the gratifying confidence hitherto manifested by the profession in all works bearing our imprint.

BLANCHARD & LEA.

PHILADELPHIA, March, 1857.

* * A copy of our new Illustrated Catalogue, 80 pages, octavo, will be sent *free* to any address on receipt of nine cents in postage stamps. Catalogues of our numerous publications in miscellaneous and educational literature forwarded on application.

TWO MEDICAL PERIODICALS, FREE OF POSTAGE, FOR FIVE DOLLARS PER ANNUM.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, subject to postage, when not paid for in advance, - - - - -	\$5 00
THE MEDICAL NEWS AND LIBRARY, invariably in advance, - - -	1 00
or, BOTH PERIODICALS furnished, FREE OF POSTAGE, for Five Dollars remitted in advance.	

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, EDITED BY ISAAC HAYS, M. D.,

is published Quarterly, on the first of January, April, July, and October. Each number contains at least two hundred and eighty large octavo pages, handsomely and appropriately illustrated, wherever necessary. It has now been issued regularly for more than THIRTY-FIVE years, and it has been under the control of the present editor for more than a quarter of a century. Throughout this long period, it has maintained its position in the highest rank of medical periodicals both at home and abroad, and has received the cordial support of the entire profession in this country. Its list of Collaborators will be found to contain a large number of the most distinguished names of the profession in every section of the United States, rendering the department devoted to

ORIGINAL COMMUNICATIONS

full of varied and important matter, of great interest to all practitioners.

As the aim of the Journal, however, is to combine the advantages presented by all the different varieties of periodicals, in its

REVIEW DEPARTMENT

will be found extended and impartial reviews of all important new works, presenting subjects of novelty and interest, together with very numerous

BIBLIOGRAPHICAL NOTICES,

including nearly all the medical publications of the day, both in this country and Great Britain, with a choice selection of the more important continental works. This is followed by the

QUARTERLY SUMMARY,

being a very full and complete abstract, methodically arranged, of the

IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES.

This department of the Journal, so important to the practising physician, is the object of especial care on the part of the editor. It is classified and arranged under different heads, thus facilitating the researches of the reader in pursuit of particular subjects, and will be found to present a very full and accurate digest of all observations, discoveries, and inventions recorded in every branch of medical science. The very extensive arrangements of the publishers are such as to afford to the editor complete materials for this purpose, as he not only regularly receives

ALL THE AMERICAN MEDICAL AND SCIENTIFIC PERIODICALS,

but also twenty or thirty of the more important Journals issued in Great Britain and on the Continent, thus enabling him to present in a convenient compass a thorough and complete abstract of everything interesting or important to the physician occurring in any part of the civilized world.

An evidence of the success which has attended these efforts may be found in the constant and steady increase in the subscription list, which renders it advisable for gentlemen desiring the Journal, to make known their wishes at an early day, in order to secure a year's set with certainty, the publishers having frequently been unable to supply copies when ordered late in the year. To their old subscribers, many of whom have been on their list for twenty or thirty years, the publishers feel that no promises are necessary; but those who may desire for the first time to subscribe, can rest assured that no exertion will be spared to maintain the Journal in the high position which it has occupied for so long a period.

By reference to the terms it will be seen that, in addition to this large amount of valuable and practical information on every branch of medical science, the subscriber, by paying in advance, becomes entitled, without further charge, to

THE MEDICAL NEWS AND LIBRARY,

a monthly periodical of thirty-two large octavo pages. Its "NEWS DEPARTMENT" presents the current information of the day, while the "LIBRARY DEPARTMENT" is devoted to presenting standard works on various branches of medicine. Within a few years, subscribers have thus received, without expense, the following works which have passed through its columns:—

WATSON'S LECTURES ON THE PRACTICE OF PHYSIC.

BRODIE'S CLINICAL LECTURES ON SURGERY.

TODD AND BOWMAN'S PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN.

724 pages, with numerous wood-cuts, being all that has yet appeared in England.

WEST'S LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD.

MALGAIGNE'S OPERATIVE SURGERY, with wood-cuts.

SIMON'S LECTURES ON GENERAL PATHOLOGY.

BENNETT ON PULMONARY TUBERCULOSIS, with wood-cuts,

WEST ON ULCERATION OF THE OS UTERI, and

BROWN ON THE SURGICAL DISEASES OF FEMALES, with wood-cuts.

While in the number for July, 1856, is commenced a new and highly important work,

WEST'S LECTURES ON THE DISEASES OF WOMEN.

PART I.—DISEASES OF THE UTERUS.

The very favorable reception accorded by the profession to the valuable "LECTURES ON THE DISEASES OF CHILDREN," by the same author, which likewise appeared in this periodical, has induced the publishers to secure the present work for their subscribers, from advance sheets, supplied by the author. The very high reputation of Dr. West, and the unusual clinical advantages which he has enjoyed, sufficiently indicate the practical value of a systematic work from his pen on so important a subject. The publishers, therefore, trust that its appearance in the "News" will afford entire satisfaction to their numerous subscribers, who will thus receive it free of all expense.

For a more extended advertisement, see p. 32.

It will thus be seen that for the small sum of FIVE DOLLARS, paid in advance, the subscriber will obtain a Quarterly and a Monthly periodical,

EMBRACING ABOUT FIFTEEN HUNDRED LARGE OCTAVO PAGES,

mailed to any part of the United States, free of postage.

These very favorable terms are now presented by the publishers with the view of removing all difficulties and objections to a full and extended circulation of the Medical Journal to the office of every member of the profession throughout the United States. The rapid extension of mail facilities will now place the numbers before subscribers with a certainty and dispatch not heretofore attainable; while by the system now proposed, every subscriber throughout the Union is placed upon an equal footing, at the very reasonable price of Five Dollars for two periodicals, without further expense.

Those subscribers who do not pay in advance will bear in mind that their subscription of Five Dollars will entitle them to the Journal only, without the News, and that they will be at the expense of their own postage on the receipt of each number. The advantage of a remittance when ordering the Journal will thus be apparent.

As the Medical News and Library is in no case sent without advance payment, its subscribers will always receive it free of postage.

Remittances of subscriptions can be mailed at our risk, when a certificate is taken from the Postmaster that the money is duly inclosed and forwarded.

Address

BLANCHARD & LEA, PHILADELPHIA.


ALLEN (J. M.), M. D.,

Professor of Anatomy in the Pennsylvania Medical College, &c.

THE PRACTICAL ANATOMIST; or, The Student's Guide in the Dissecting-ROOM. With 266 illustrations. In one handsome royal 12mo. volume, of over 600 pages, leather. \$2 25. (*Now Ready.*)

In the arrangement of this work, the author has endeavored to present a complete and thorough course of dissections in a clearer and more available form for practical use, than has as yet been accomplished. The chapters follow each other in the order in which dissections are usually conducted in this country, and as each region is taken up, every detail regarding it is fully described and illustrated, so that the student is not interrupted in his labors, by the necessity of referring from one portion of the volume to another.

ANALYTICAL COMPENDIUM

OF MEDICAL SCIENCE, containing Anatomy, Physiology, Surgery, Midwifery, Chemistry, Materia Medica, Therapeutics, and Practice of Medicine. By JOHN NEILL, M. D., and F. G. SMITH, M. D. New and enlarged edition, one thick volume royal 12mo. of over 1000 pages, with 374 illustrations.  See NEILL, p. 24.

ABEL (F. A.), F. C. S. AND C. L. BLOXAM.

HANDBOOK OF CHEMISTRY, Theoretical, Practical, and Technical; with a Recommendatory Preface by Dr. HOFMANN. In one large octavo volume, extra cloth, of 662 pages, with illustrations. \$3 25.

It must be understood that this is a work fitted for the earnest student, who resolves to pursue for himself a steady search into the chemical mysteries of creation. For such a student the 'Handbook' will prove an excellent guide, since he will find in it, not merely the approved modes of analytical investigation, but most descriptions of the apparatus necessary, with such manipulatory details as rendered Faraday's 'Chemical Manipulations' so valuable at the time of its publication. Beyond this, the importance of the work is increased by the introduction of much of the technical chemistry of the manufactory.—*Dr. Hofmann's Preface.*

ASHWELL (SAMUEL), M. D.,

Obstetric Physician and Lecturer to Guy's Hospital, London.

A PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN.

Illustrated by Cases derived from Hospital and Private Practice. Third American, from the Third and revised London edition. In one octavo volume, extra cloth, of 528 pages. (*Lately Published.*) \$3 00.

The most useful practical work on the subject in the English language.—*Boston Med. and Surg. Journal.*

The most able, and certainly the most standard and practical, work on female diseases that we have yet seen.—*Medico-Chirurgical Review.*

The young practitioner will find it invaluable, while those who have had most experience will yet

find something to learn, and much to commend, in a book which shows so much patient observation, practical skill, and sound sense.—*British and Foreign Med. Review.*

We commend it to our readers as the best practical treatise on the subject which has yet appeared.—*London Lancet.*

ARNOTT (NEILL), M. D.

ELEMENTS OF PHYSICS; or Natural Philosophy, General and Medical.

Written for universal use, in plain or non-technical language. A new edition, by ISAAC HAYS, M. D. Complete in one octavo volume, leather, of 484 pages, with about two hundred illustrations. \$2 50.

BENNETT (HENRY), M. D.

A PRACTICAL TREATISE ON INFLAMMATION OF THE UTERUS,

ITS CERVIX AND APPENDAGES, and on its connection with Uterine Disease. Fourth American, from the third and revised London edition. To which is added (*July, 1856*), a REVIEW OF THE PRESENT STATE OF UTERINE PATHOLOGY. In one neat octavo volume, extra cloth, of 500 pages, with wood-cuts. \$2 00.

The addition of the "REVIEW" presents the most recent aspects of the questions discussed in this well-known work, bringing it down to the latest moment.

This edition has been carefully revised and altered, and various additions have been made, which render it more complete, and, if possible, more worthy of the high appreciation in which it is held by the medical profession throughout the world. A copy should be in the possession of every physician.—*Charleston Med. Journal and Review.*

We are firmly of opinion that in proportion as a knowledge of uterine diseases becomes more appreciated, this work will be proportionably established as a text-book in the profession.—*The Lancet.*

Also, just ready, by the same author, and for sale separate,

A REVIEW OF THE PRESENT STATE OF UTERINE PATHOLOGY.

1 small vol. 8vo. 50 cents, in flexible cloth.

In this little work, which can be had either in connection with the "Practical Treatise," or separate, the author presents his latest views with regard to the various doctrines which have recently been brought forward on this interesting question, under the following heads:—

CHAP. I. Preliminary. II. Sketch of Uterine Pathology. III. Objections. IV. The Leucorrhœa Theory—the Syphilis Theory—the Ovarian Theory. V. The Displacement Theory. VI. Summary.

BROWN (ISAAC BAKER),

Surgeon-Accoucheur to St. Mary's Hospital, &c.

ON SOME DISEASES OF WOMEN ADMITTING OF SURGICAL TREATMENT. With handsome illustrations. One vol. 8vo., extra cloth. (*Now Ready.*) \$1 60.

Mr. Brown has earned for himself a high reputation in the operative treatment of sundry diseases and injuries to which females are peculiarly subject. We can truly say of his work that it is an important addition to obstetrical literature. The operative suggestions and contrivances which Mr. Brown describes, exhibit much practical sagacity and skill,

and merit the careful attention of every surgeon-accoucheur.—*Association Journal.*

We have no hesitation in recommending this book to the careful attention of all surgeons who make female complaints a part of their study and practice.—*Dublin Quarterly Journal.*

BENNETT (J. HUGHES), M. D., F. R. S. E.,

Professor of Clinical Medicine in the University of Edinburgh, &c.

THE PATHOLOGY AND TREATMENT OF PULMONARY TUBERCULOSIS, and on the Local Medication of Pharyngeal and Laryngeal Diseases frequently mistaken for or associated with, Phthisis. In one handsome octavo volume, extra cloth, with beautiful wood-cuts. pp. 130. (*Lately Issued.*) \$1 25.

BUDD (GEORGE), M. D., F. R. S.,

Professor of Medicine in King's College, London.

ON DISEASES OF THE LIVER. Second American, from the second and enlarged London edition. In one very handsome octavo volume, extra cloth, with four beautifully colored plates, and numerous wood-cuts. pp. 468. \$3 00.

For many years, Dr. Budd's work must be the authority of the great mass of British practitioners on the hepatic diseases; and it is satisfactory that

the subject has been taken up by so able and experienced a physician.—*British and Foreign Medico-Chirurgical Review.*

BY THE SAME AUTHOR. (*Now Ready.*)

ON THE ORGANIC DISEASES AND FUNCTIONAL DISORDERS OF THE STOMACH. In one neat octavo volume, extra cloth. \$1 50. A new work, 1856.

While special treatises have been multiplying upon almost all the different classes of diseases, there has long been felt the want of an authoritative work on the disorders of the stomach, which constitute, perhaps, a larger proportion of the daily practice of the physician than any other class of maladies. To supply this want has been the object of the author, and his reputation is an ample guarantee of the value of his labors.

From the high position occupied by Dr. Budd as a teacher, a writer, and a practitioner, it is almost needless to state that the present book may be consulted with great advantage. It is written in an easy

style, the subjects are well arranged, and the practical precepts, both of diagnosis and treatment, denote the character of a thoughtful and experienced physician.—*London Med. Times and Gazette*, Dec. 1855.

BIRD (GOLDING), A. M., M. D., &c.

URINARY DEPOSITS: THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS. A new and enlarged American, from the last improved London edition. With over sixty illustrations. In one royal 12mo. vol, extra cloth. pp. 372. \$1 30.

It can scarcely be necessary for us to say anything of the merits of this well-known Treatise, which so admirably brings into practical application the results of those microscopical and chemical researches regarding the physiology and pathology of the urinary secretion, which have contributed so much to the increase of our diagnostic powers, and to the

extension and satisfactory employment of our therapeutic resources. In the preparation of this new edition of his work, it is obvious that Dr. Golding Bird has spared no pains to render it a faithful representation of the present state of scientific knowledge on the subject it embraces.—*The British and Foreign Medico-Chirurgical Review.*

BY THE SAME AUTHOR.

ELEMENTS OF NATURAL PHILOSOPHY; being an Experimental Introduction to the Physical Sciences. Illustrated with nearly four hundred wood-cuts. From the third London edition. In one neat volume, royal 12mo., extra cloth. pp. 402. \$1 25.

BILLING'S PRINCIPLES OF MEDICINE.—Second American, from the Fifth and Improved London edition. In one handsome octavo volume, extra cloth. 250 pages. \$1 25.

BLAKISTON'S PRACTICAL OBSERVATIONS ON CERTAIN DISEASES OF THE CHEST, and on the Principles of Auscultation. In one vol., cloth, 8vo pp. 384. \$1 25.

BURROWS ON DISORDERS OF THE CEREBRAL CIRCULATION, and on the Connection between the Affections of the Brain and Diseases of the Heart. In one 8vo. vol., extra cloth, with colored plates. pp. 216. \$1 25.

BEALE ON THE LAWS OF HEALTH IN RELATION TO MIND AND BODY. A Series of Letters from an old Practitioner to a Patient. In one volume, royal 12mo., extra cloth. pp. 296. 60 cents.

BUSHNAN'S PHYSIOLOGY OF ANIMAL AND VEGETABLE LIFE; a Popular Treatise on the Functions and Phenomena of Organic Life. In one handsome royal 12mo. volume, extra cloth, with over 100 illustrations. pp. 234. 80 cents.

BUCKLER ON THE ETIOLOGY, PATHOLOGY, AND TREATMENT OF FIBRO-BRONCHITIS AND RHEUMATIC PNEUMONIA. In one 8vo. volume, extra cloth. pp. 150. \$1 25.

BLOOD AND URINE (MANUALS ON). BY JOHN WILLIAM GRIFFITH, G. OWEN REESE, AND ALFRED MARKWICK. One thick volume, royal 12mo., extra cloth, with plates. pp. 460. \$1 25.

BRODIE'S CLINICAL LECTURES ON SURGERY. 1 vol. 8vo., cloth. 350 pp. \$1 25.

BARLOW (GEORGE H.), M. D.

Physician to Guy's Hospital, London, &c.

A MANUAL OF THE PRACTICE OF MEDICINE. With Additions by D.F. CONDIE, M. D., author of "A Practical Treatise on Diseases of Children," &c. In one handsome octavo volume, leather, of over 600 pages. (*A new work, just ready, 1856.*) \$2 75.

The position of the author as physician to Guy's Hospital and other large public institutions, is a sufficient guarantee of the extent and value of the experience which is here systematically recorded and condensed. His aim throughout has been to produce a *practical* work, on which the student can rely as a guide, and to which the practitioner can refer with confidence. The additions by the editor comprise chapters on Cerebro-spinal Meningitis, Cholera Infantum, and Yellow Fever, besides numerous notes wherever the diseases or practice of this country seemed to render them necessary or desirable.

We most emphatically commend it to the attention of the profession, as deserving their confidence—a depository of practical knowledge, from which they may draw with great benefit.—*Cincinnati Med. Observer*, Mar. 1856.

The student has long been in want of a good elementary work on the Practice of Medicine. In Dr. Barlow's Manual that want is supplied; and we have no question that it will at once be installed as the favorite text-book in all Medical Schools.—*Medical Times and Gazette*.

We recommend Dr. Barlow's Manual in the warmest manner as a most valuable *vade-mecum*. We have had frequent occasion to consult it, and have found it clear, concise, practical, and sound. It is eminently a practical work, containing all that is essential, and avoiding useless theoretical discus-

sion. The work supplies what has been for some time wanting, a manual of practice based upon modern discoveries in pathology and rational views of treatment of disease. It is especially intended for the use of students and junior practitioners, but it will be found hardly less useful to the experienced physician. The American editor has added to the work three chapters—on Cholera Infantum, Yellow Fever, and Cerebro-spinal Meningitis. These additions, the two first of which are indispensable to a work on practice destined for the profession in this country, are executed with great judgment and fidelity, by Dr. Condie, who has also succeeded happily in imitating the conciseness and clearness of style which are such agreeable characteristics of the original book.—*Boston Med. and Surg. Journal*, Feb. 1856.

BARTLETT (ELISHA), M. D.**THE HISTORY, DIAGNOSIS, AND TREATMENT OF THE FEVERS**

OF THE UNITED STATES. A new and revised edition. By ALONZO CLARK, M. D., Prof. of Pathology and Practical Medicine in the N. Y. College of Physicians and Surgeons, &c. In one octavo volume, of six hundred pages, extra cloth. (*Now Ready.*) Price \$3 00.

The position which this work has obtained as one of our medical classics, renders unnecessary any remark further than to say that the editor, in executing the task assigned to him by the late author, has endeavored to render the work a faithful exposition of the subject in its most advanced condition. To effect this, a considerable amount of matter has been introduced, but by a slight enlargement of the page it has been accommodated without unduly increasing the bulk of the volume. The reputation of the editor as an accurate observer and philosophical writer is sufficient guarantee that, in his hands, the work will fully maintain its former character.

The masterly and elegant treatise by Dr. Bartlett is invaluable to the American student and practitioner.—*Dr. Holmes's Report to the Nat. Med. Association*.

We regard it, from the examination we have made of it, the best work on fevers extant in our language,

and as such cordially recommend it to the medical public.—*St. Louis Medical and Surgical Journal*.

Take it altogether, it is the most complete history of our fevers which has yet been published, and every practitioner should avail himself of its contents.—*The Western Lancet*.

BOWMAN (JOHN E.), M. D.

PRACTICAL HANDBOOK OF MEDICAL CHEMISTRY. Second American, from the third and revised English Edition. In one neat volume, royal 12mo., extra cloth, with numerous illustrations. pp. 288. (*Now Ready, 1856.*) \$1 25.

Presenting, in a condensed and convenient form, at a very low price, the applications of Chemistry to the practical purposes of Clinical Medicine, this work supplies a want which has long been felt by the physician. The numerous editions which have been called for both in England and this country, sufficiently attest the success with which the author has carried out his plan.

BY THE SAME AUTHOR. (*Now Ready.*)

INTRODUCTION TO PRACTICAL CHEMISTRY, INCLUDING ANALYSIS.

Second American, from the second and revised London edition. With numerous illustrations. In one neat vol., royal 12mo., extra cloth. pp. 350. \$1 25.

CURLING (T. B.), F. R. S.,

Surgeon to the London Hospital, President of the Hunterian Society, &c.

A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM.

Second American, from the second and enlarged English edition. In one handsome octavo volume, extra cloth, with numerous illustrations. pp. 420. (*Now Ready, 1856.*) \$2 00.

In the revised English edition, of which this is a reprint, the author, for want of space, omitted the Anatomical Introduction. By a more condensed style of printing, room has been found in the present volume to retain this important portion without rendering the work inconveniently large. Some of the notes of the former American editor have also been incorporated, and a number of new illustrations introduced. With these improvements, and the thorough revision which it has enjoyed at the hands of the author, it will be found fully worthy to retain the authoritative position which it has acquired with regard to this class of affections.

CARPENTER (WILLIAM B.), M. D., F. R. S., &c.,
 Examiner in Physiology and Comparative Anatomy in the University of London.

PRINCIPLES OF HUMAN PHYSIOLOGY; with their chief applications to Psychology, Pathology, Therapeutics, Hygiene, and Forensic Medicine. A new American, from the last and revised London edition. With nearly three hundred illustrations. Edited, with additions, by FRANCIS GURNEY SMITH, M. D., Professor of the Institutes of Medicine in the Pennsylvania Medical College, &c. In one very large and beautiful octavo volume, of about nine hundred large pages, handsomely printed and strongly bound in leather, with raised bands. (*Just Issued*, 1856.) \$4 25.

In the preparation of this new edition, the author has spared no labor to render it, as heretofore, a complete and lucid exposition of the most advanced condition of its important subject. The amount of the additions required to effect this object thoroughly, joined to the former large size of the volume, presenting objections arising from the unwieldy bulk of the work, he has omitted all those portions not bearing directly upon HUMAN PHYSIOLOGY, designing to incorporate them in his forthcoming Treatise on GENERAL PHYSIOLOGY. As a full and accurate text-book on the Physiology of Man, the work in its present condition therefore presents even greater claims upon the student and physician than those which have heretofore won for it the very wide and distinguished favor which it has so long enjoyed. The additions of Prof. Smith will be found to supply whatever may have been wanting to the American student, while the introduction of many new illustrations, and the most careful mechanical execution, render the volume one of the most attractive as yet issued.

For upwards of thirteen years Dr. Carpenter's work has been considered by the profession generally, both in this country and England, as the most valuable compendium on the subject of physiology in our language. This distinction it owes to the high attainments and unwearied industry of its accomplished author. The present edition (which, like the last American one, was prepared by the author himself), is the result of such extensive revision, that it may almost be considered a new work. We need hardly say, in concluding this brief notice, that while the work is indispensable to every student of medicine in this country, it will amply repay the practitioner for its perusal by the interest and value of its contents.—*Boston Med. and Surg. Journal*.

This is a standard work—the text-book used by all medical students who read the English language. It has passed through several editions in order to keep pace with the rapidly growing science of Physiology. Nothing need be said in its praise, for its merits are universally known; we have nothing to say of its defects, for they only appear where the science of which it treats is incomplete.—*Western Lancet*.

The most complete exposition of physiology which any language can at present give.—*Brit. and For. Med.-Chirurg. Review*.

The greatest, the most reliable, and the best book on the subject which we know of in the English language.—*Stethoscope*.

To eulogize this great work would be superfluous. We should observe, however, that in this edition the author has remodelled a large portion of the former, and the editor has added much matter of interest, especially in the form of illustrations. We may confidently recommend it as the most complete work on Human Physiology in our language.—*Southern Med. and Surg. Journal*, December, 1855.

The most complete work on the science in our language.—*Am. Med. Journal*.

The most complete work now extant in our language.—*N. O. Med. Register*.

The best text-book in the language on this extensive subject.—*London Med. Times*.

A complete cyclopædia of this branch of science.—*N. Y. Med. Times*.

The profession of this country, and perhaps also of Europe, have anxiously and for some time awaited the announcement of this new edition of Carpenter's Human Physiology. His former editions have for many years been almost the only text-book on Physiology in all our medical schools, and its circulation among the profession has been unsurpassed by any work in any department of medical science.

It is quite unnecessary for us to speak of this work as its merits would justify. The mere announcement of its appearance will afford the highest pleasure to every student of Physiology, while its perusal will be of infinite service in advancing physiological science.—*Ohio Med. and Surg. Journ.*

BY THE SAME AUTHOR. (*Lately Issued*.)

PRINCIPLES OF COMPARATIVE PHYSIOLOGY. New American, from the Fourth and Revised London edition. In one large and handsome octavo volume, with over three hundred beautiful illustrations. pp. 752. Extra cloth, \$4 80; leather, raised bands, \$5 25.

The delay which has existed in the appearance of this work has been caused by the very thorough revision and remodelling which it has undergone at the hands of the author, and the large number of new illustrations which have been prepared for it. It will, therefore, be found almost a new work, and fully up to the day in every department of the subject, rendering it a reliable text-book for all students engaged in this branch of science. Every effort has been made to render its typographical finish and mechanical execution worthy of its exalted reputation, and creditable to the mechanical arts of this country.

This book should not only be read but thoroughly studied by every member of the profession. None are too wise or old, to be benefited thereby. But especially to the younger class would we cordially commend it as best fitted of any work in the English language to qualify them for the reception and comprehension of those truths which are daily being developed in physiology.—*Medical Counsellor*.

Without pretending to it, it is an encyclopedia of the subject, accurate and complete in all respects—a truthful reflection of the advanced state at which the science has now arrived.—*Dublin Quarterly Journal of Medical Science*.

A truly magnificent work—in itself a perfect physiological study.—*Ranking's Abstract*.

This work stands without its fellow. It is one few men in Europe could have undertaken; it is one

no man, we believe, could have brought to so successful an issue as Dr. Carpenter. It required for its production a physiologist at once deeply read in the labors of others, capable of taking a general, critical, and unprejudiced view of those labors, and of combining the varied, heterogeneous materials at his disposal, so as to form an harmonious whole. We feel that this abstract can give the reader a very imperfect idea of the fulness of this work, and no idea of its unity, of the admirable manner in which material has been brought, from the most various sources, to conduce to its completeness, of the lucidity of the reasoning it contains, or of the clearness of language in which the whole is clothed. Not the profession only, but the scientific world at large, must feel deeply indebted to Dr. Carpenter for this great work. It must, indeed, add largely even to his high reputation.—*Medical Times*.

CARPENTER (WILLIAM B.), M. D., F. R. S.,
Examiner in Physiology and Comparative Anatomy in the University of London.

(Now Ready, 1856.)

THE MICROSCOPE AND ITS REVELATIONS. With an Appendix containing the Applications of the Microscope to Clinical Medicine, &c. By F. G. SMITH, M. D. Illustrated by four hundred and thirty-four beautiful engravings on wood. In one large and very handsome octavo volume, of 724 pages, extra cloth, \$4 00; leather, \$4 50.

Dr. Carpenter's position as a microscopist and physiologist, and his great experience as a teacher, eminently qualify him to produce what has long been wanted—a good text-book on the practical use of the microscope. In the present volume his object has been, as stated in his Preface, "to combine, within a moderate compass, that information with regard to the use of his 'tools,' which is most essential to the working microscopist, with such an account of the objects best fitted for his study, as might qualify him to comprehend what he observes, and might thus prepare him to benefit science, whilst expanding and refreshing his own mind." That he has succeeded in accomplishing this, no one acquainted with his previous labors can doubt.

The great importance of the microscope as a means of diagnosis, and the number of microscopists who are also physicians, have induced the American publishers, with the author's approval, to add an Appendix, carefully prepared by Professor Smith, on the applications of the instrument to clinical medicine, together with an account of American Microscopes, their modifications and accessories. This portion of the work is illustrated with nearly one hundred wood-cuts, and, it is hoped, will adapt the volume more particularly to the use of the American student.

Every care has been taken in the mechanical execution of the work, which is confidently presented as in no respect inferior to the choicest productions of the London press.

The mode in which the author has executed his intentions may be gathered from the following condensed synopsis of the

CONTENTS.

INTRODUCTION—History of the Microscope. CHAP. I. Optical Principles of the Microscope. CHAP. II. Construction of the Microscope. CHAP. III. Accessory Apparatus. CHAP. IV. Management of the Microscope. CHAP. V. Preparation, Mounting, and Collection of Objects. CHAP. VI. Microscopic Forms of Vegetable Life—Protophytes. CHAP. VII. Higher Cryptogamia. CHAP. VIII. Phanerogamic Plants. CHAP. IX. Microscopic Forms of Animal Life—Protozoa—Animalcules. CHAP. X. Foraminifera, Polycystina, and Sponges. CHAP. XI. Zoophytes. CHAP. XII. Echinodermata. CHAP. XIII. Polyzoa and Compound Tunicata. CHAP. XIV. Molluscan Animals Generally. CHAP. XV. Annulosa. CHAP. XVI. Crustacea. CHAP. XVII. Insects and Arachnida. CHAP. XVIII. Vertebrated Animals. CHAP. XIX. Applications of the Microscope to Geology. CHAP. XX. Inorganic or Mineral Kingdom—Polarization. APPENDIX. Microscope as a means of Diagnosis—Injections—Microscopes of American Manufacture.

Those who are acquainted with Dr. Carpenter's previous writings on Animal and Vegetable Physiology, will fully understand how vast a store of knowledge he is able to bring to bear upon so comprehensive a subject as the revelations of the microscope;

and even those who have no previous acquaintance with the construction or uses of this instrument, will find abundance of information conveyed in clear and simple language.—*Med. Times and Gazette.*

BY THE SAME AUTHOR.

ELEMENTS (OR MANUAL) OF PHYSIOLOGY, INCLUDING PHYSIOLOGICAL ANATOMY. Second American, from a new and revised London edition. With one hundred and ninety illustrations. In one very handsome octavo volume, leather. pp. 566. \$3 00.

In publishing the first edition of this work, its title was altered from that of the London volume, by the substitution of the word "Elements" for that of "Manual," and with the author's sanction the title of "Elements" is still retained as being more expressive of the scope of the treatise.

To say that it is the best manual of Physiology now before the public, would not do sufficient justice to the author.—*Buffalo Medical Journal.*

In his former works it would seem that he had exhausted the subject of Physiology. In the present, he gives the essence, as it were, of the whole.—*N. Y. Journal of Medicine.*

Those who have occasion for an elementary treatise on Physiology, cannot do better than to possess themselves of the manual of Dr. Carpenter.—*Medical Examiner.*

The best and most complete exposé of modern Physiology, in one volume, extant in the English language.—*St. Louis Medical Journal.*

BY THE SAME AUTHOR. (Preparing.)

PRINCIPLES OF GENERAL PHYSIOLOGY, INCLUDING ORGANIC CHEMISTRY AND HISTOLOGY. With a General Sketch of the Vegetable and Animal Kingdom. In one large and very handsome octavo volume, with several hundred illustrations.

The subject of general physiology having been omitted in the last editions of the author's "Comparative Physiology" and "Human Physiology," he has undertaken to prepare a volume which shall present it more thoroughly and fully than has yet been attempted, and which may be regarded as an introduction to his other works.

BY THE SAME AUTHOR.

A PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New edition, with a Preface by D. F. CONDIE, M. D., and explanations of scientific words. In one neat 12mo. volume, extra cloth. pp. 178. (Just Issued.) 50 cents.

CHELIUS (J. M.), M. D.,
Professor of Surgery in the University of Heidelberg, &c.

A SYSTEM OF SURGERY. Translated from the German, and accompanied with additional Notes and References, by JOHN F. SOUTH. Complete in three very large octavo volumes, of nearly 2200 pages, strongly bound, with raised bands and double titles. \$10 00.

CONDIE (D. F.), M. D., &c.

A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Fourth edition, revised and augmented. In one large volume, 8vo., leather, of nearly 750 pages. \$3 00.

FROM THE AUTHOR'S PREFACE.

The demand for another edition has afforded the author an opportunity of again subjecting the entire treatise to a careful revision, and of incorporating in it every important observation recorded since the appearance of the last edition, in reference to the pathology and therapeutics of the several diseases of which it treats.

In the preparation of the present edition, as in those which have preceded, while the author has appropriated to his use every important fact that he has found recorded in the works of others, having a direct bearing upon either of the subjects of which he treats, and the numerous valuable observations—pathological as well as practical—dispersed throughout the pages of the medical journals of Europe and America, he has, nevertheless, relied chiefly upon his own observations and experience, acquired during a long and somewhat extensive practice, and under circumstances peculiarly well adapted for the clinical study of the diseases of early life.

Every species of hypothetical reasoning has, as much as possible, been avoided. The author has endeavored throughout the work to confine himself to a simple statement of well-ascertained pathological facts, and plain therapeutical directions—his chief desire being to render it what its title imports it to be, **A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.**

Dr. Condie's scholarship, acumen, industry, and practical sense are manifested in this, as in all his numerous contributions to science.—*Dr. Holmes's Report to the American Medical Association.*

Taken as a whole, in our judgment, Dr. Condie's Treatise is one from the perusal of which the practitioner in this country will rise with the greatest satisfaction.—*Western Journal of Medicine and Surgery.*

One of the best works upon the Diseases of Children in the English language.—*Western Lancet.*

Perhaps the most full and complete work now before the profession of the United States; indeed, we may say in the English language. It is vastly superior to most of its predecessors.—*Transylvania Med. Journal.*

We feel assured from actual experience that no physician's library can be complete without a copy of this work.—*N. Y. Journal of Medicine.*

A veritable pædiatric encyclopædia, and an honor to American medical literature.—*Ohio Medical and Surgical Journal.*

We feel persuaded that the American medical profession will soon regard it not only as a very good, but as the VERY BEST "Practical Treatise on the Diseases of Children."—*American Medical Journal.*

We pronounced the first edition to be the best work on the diseases of children in the English language, and, notwithstanding all that has been published, we still regard it in that light.—*Medical Examiner.*

CHRISTISON (ROBERT), M. D., V. P. R. S. E., &c.

A DISPENSATORY; or, Commentary on the Pharmacopœias of Great Britain and the United States; comprising the Natural History, Description, Chemistry, Pharmacy, Actions, Uses, and Doses of the Articles of the Materia Medica. Second edition, revised and improved, with a Supplement containing the most important New Remedies. With copious Additions, and two hundred and thirteen large wood-engravings. By R. EGLESFELD GRIFFITH, M. D. In one very large and handsome octavo volume, leather, raised bands, of over 1000 pages. \$3 50.

It is not needful that we should compare it with the other pharmacopœias extant, which enjoy and merit the confidence of the profession: it is enough to say that it appears to us as perfect as a Dispensatory, in the present state of pharmaceutical science, could be made. If it omits any details pertaining to

this branch of knowledge which the student has a right to expect in such a work, we confess the omission has escaped our scrutiny. We cordially recommend this work to such of our readers as are in need of a Dispensatory. They cannot make choice of a better.—*Western Journ. of Medicine and Surgery.*

COOPER (BRANSBY B.), F. R. S.

LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY.

In one very large octavo volume, extra cloth, of 750 pages. \$3 00.

COOPER ON DISLOCATIONS AND FRACTURES OF THE JOINTS.—Edited by BRANSBY B. COOPER, F. R. S., &c. With additional Observations by Prof. J. C. WARREN. A new American edition. In one handsome octavo volume, extra cloth, of about 500 pages, with numerous illustrations on wood. \$3 25.

COOPER ON THE ANATOMY AND TREATMENT OF ABDOMINAL HERNIA. One large volume, imperial 8vo., extra cloth, with over 130 lithographic figures. \$2 50.

COOPER ON THE ANATOMY AND DISEASES OF THE BREAST, with twenty-five Miscellaneous and Surgical Papers. One large volume, imperial 8vo., extra cloth, with 252 figures, on 36 plates. \$2 50.

COOPER ON THE STRUCTURE AND DIS-

EASES OF THE TESTIS, AND ON THE THYMUS GLAND. One vol. imperial 8vo., extra cloth, with 177 figures on 29 plates. \$2 00.

COPLAND ON THE CAUSES, NATURE, AND TREATMENT OF PALSY AND APOPLEXY. In one volume, royal 12mo., extra cloth. pp. 326. 80 cents.

CLYMER ON FEVERS; THEIR DIAGNOSIS, PATHOLOGY, AND TREATMENT In one octavo volume, leather, of 600 pages. \$1 50.

COLOMBAT DE L'ISERE ON THE DISEASES OF FEMALES, and on the special Hygiene of their Sex. Translated, with many Notes and Additions, by C. D. MEIGS, M. D. Second edition, revised and improved. In one large volume, octavo, leather, with numerous wood-cuts. pp. 720. \$3 50.

CARSON (JOSEPH), M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

SYNOPSIS OF THE COURSE OF LECTURES ON MATERIA MEDICA AND PHARMACY, delivered in the University of Pennsylvania. Second and revised edition. In one very neat octavo volume, extra cloth, of 208 pages. (*Now Ready.*) \$1 50.

CHURCHILL (FLEETWOOD), M. D., M. R. I. A.

ON THE THEORY AND PRACTICE OF MIDWIFERY. A new American, from the last and improved English edition. Edited, with Notes and Additions, by D. FRANCIS CONDIE, M. D., author of a "Practical Treatise on the Diseases of Children," &c. With 139 illustrations. In one very handsome octavo volume, leather. pp. 510. \$3 00.

To bestow praise on a book that has received such marked approbation would be superfluous. We need only say, therefore, that if the first edition was thought worthy of a favorable reception by the medical public, we can confidently affirm that this will be found much more so. The lecturer, the practitioner, and the student, may all have recourse to its pages, and derive from their perusal much interest and instruction in everything relating to theoretical and practical midwifery.—*Dublin Quarterly Journal of Medical Science.*

A work of very great merit, and such as we can confidently recommend to the study of every obstetric practitioner.—*London Medical Gazette.*

This is certainly the most perfect system extant. It is the best adapted for the purposes of a text-book, and that which he whose necessities confine him to one book, should select in preference to all others.—*Southern Medical and Surgical Journal.*

The most popular work on midwifery ever issued from the American press.—*Charleston Med. Journal.*

Were we reduced to the necessity of having but one work on midwifery, and permitted to choose, we would unhesitatingly take Churchill.—*Western Med. and Surg. Journal.*

It is impossible to conceive a more useful and elegant manual than Dr. Churchill's Practice of Midwifery.—*Provincial Medical Journal.*

Certainly, in our opinion, the very best work on the subject which exists.—*N. Y. Annalist.*

No work holds a higher position, or is more deserving of being placed in the hands of the tyro, the advanced student, or the practitioner.—*Medical Examiner.*

Previous editions, under the editorial supervision of Prof. R. M. Huston, have been received with marked favor, and they deserved it; but this, reprinted from a very late Dublin edition, carefully revised and brought up by the author to the present time, does present an unusually accurate and able exposition of every important particular embraced in the department of midwifery. * * The clearness, directness, and precision of its teachings, together with the great amount of statistical research which its text exhibits, have served to place it already in the foremost rank of works in this department of remedial science.—*N. O. Med. and Surg. Journal.*

In our opinion, it forms one of the best if not the very best text-book and epitome of obstetric science which we at present possess in the English language.—*Monthly Journal of Medical Science.*

The clearness and precision of style in which it is written, and the great amount of statistical research which it contains, have served to place it in the first rank of works in this department of medical science.—*N. Y. Journal of Medicine.*

Few treatises will be found better adapted as a text-book for the student, or as a manual for the frequent consultation of the young practitioner.—*American Medical Journal.*

BY THE SAME AUTHOR. (Now Ready, 1856.)

ON THE DISEASES OF INFANTS AND CHILDREN. Second American

Edition, revised and enlarged by the author. Edited, with Notes, by W. V. KEATING, M. D. In one large and handsome volume, extra cloth, of over 700 pages. \$3 00, or in leather, \$3 25.

In preparing this work a second time for the American profession, the author has spared no labor in giving it a very thorough revision, introducing several new chapters, and rewriting others, while every portion of the volume has been subjected to a severe scrutiny. The efforts of the American editor have been directed to supplying such information relative to matters peculiar to this country as might have escaped the attention of the author, and the whole may, therefore, be safely pronounced one of the most complete works on the subject accessible to the American Profession. By an alteration in the size of the page, these very extensive additions have been accommodated without unduly increasing the size of the work.

A few notices of the former edition are subjoined:—

We regard this volume as possessing more claims to completeness than any other of the kind with which we are acquainted. Most cordially and earnestly, therefore, do we commend it to our professional brethren, and we feel assured that the stamp of their approbation will in due time be impressed upon it. After an attentive perusal of its contents, we hesitate not to say, that it is one of the most comprehensive ever written upon the diseases of children, and that, for copiousness of reference, extent of research, and perspicuity of detail, it is scarcely to be equalled, and not to be excelled, in any language.—*Dublin Quarterly Journal.*

After this meagre, and we know, very imperfect notice of Dr. Churchill's work, we shall conclude by saying, that it is one that cannot fail from its copiousness, extensive research, and general accuracy, to exalt still higher the reputation of the author in this country. The American reader will be particularly pleased to find that Dr. Churchill has done full justice throughout his work to the various American authors on this subject. The names of Dewees, Eberle, Condie, and Stewart, occur on nearly every page, and these authors are constantly referred to by the author in terms of the highest praise, and with the most liberal courtesy.—*The Medical Examiner.*

The present volume will sustain the reputation acquired by the author from his previous works. The reader will find in it full and judicious directions for the management of infants at birth, and a compendious, but clear account of the diseases to which children are liable, and the most successful mode of treating them. We must not close this notice without calling attention to the author's style, which is perspicuous and polished to a degree, we regret to say, not generally characteristic of medical works. We recommend the work of Dr. Churchill most cordially, both to students and practitioners, as a valuable and reliable guide in the treatment of the diseases of children.—*Am. Journ. of the Med. Sciences.*

We know of no work on this department of Practical Medicine which presents so candid and unprejudiced a statement or posting up of our actual knowledge as this.—*N. Y. Journal of Medicine.*

Its claims to merit both as a scientific and practical work, are of the highest order. Whilst we would not elevate it above every other treatise on the same subject, we certainly believe that very few are equal to it, and none superior.—*Southern Med. and Surgical Journal.*

BY THE SAME AUTHOR.

ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN. Selected from the writings of British Authors previous to the close of the Eighteenth Century. In one neat octavo volume, extra cloth, of about 450 pages. \$2 50.

CHURCHILL (FLEETWOOD), M. D., M. R. I. A., &c.

ON THE DISEASES OF WOMEN; including those of Pregnancy and Child-bed. A new American edition, revised by the Author. With Notes and Additions, by D. FRANCIS CONDIE, M. D., author of "A Practical Treatise on the Diseases of Children." With numerous illustrations. In one large and handsome octavo volume. (*Nearly Ready.*)

This edition of Dr. Churchill's very popular treatise may almost be termed a new work, so thoroughly has he revised it in every portion. It will be found greatly enlarged, and thoroughly brought up to the most recent condition of the subject, while the very handsome series of illustrations introduced, representing such pathological conditions as can be accurately portrayed, present a novel feature, and afford valuable assistance to the young practitioner. Such additions as appeared desirable for the American student have been made by the editor, Dr. Condie, while a marked improvement in the mechanical execution keeps pace with the advance in all other respects which the volume has undergone.

A few notices of the former edition are subjoined:—

We now regretfully take leave of Dr. Churchill's book. Had our typographical limits permitted, we should gladly have borrowed more from its richly stored pages. In conclusion, we heartily recommend it to the profession, and would at the same time express our firm conviction that it will not only add to the reputation of its author, but will prove a work of great and extensive utility to obstetric practitioners.—*Dublin Medical Press.*

Former editions of this work have been noticed in previous numbers of the Journal. The sentiments of high commendation expressed in those notices, have only to be repeated in this; not from the fact that the profession at large are not aware of the high merits which this work really possesses, but from a desire to see the principles and doctrines therein contained more generally recognized, and more universally carried out in practice.—*N. Y. Journal of Medicine.*

We know of no author who deserves that approbation, on "the diseases of females," to the same extent that Dr. Churchill does. His, indeed, is the only thorough treatise we know of on the subject; and it may be commended to practitioners and students as a masterpiece in its particular department. The former editions of this work have been commended strongly in this journal, and they have won their way to an extended, and a well-deserved popu-

larity. This fifth edition, before us, is well calculated to maintain Dr. Churchill's high reputation. It was revised and enlarged by the author, for his American publishers, and it seems to us that there is scarcely any species of desirable information on its subjects that may not be found in this work.—*The Western Journal of Medicine and Surgery.*

We are gratified to announce a new and revised edition of Dr. Churchill's valuable work on the diseases of females. We have ever regarded it as one of the very best works on the subjects embraced within its scope, in the English language; and the present edition, enlarged and revised by the author, renders it still more entitled to the confidence of the profession. The valuable notes of Prof. Huston have been retained, and contribute, in no small degree, to enhance the value of the work. It is a source of congratulation that the publishers have permitted the author to be, in this instance, his own editor, thus securing all the revision which an author alone is capable of making.—*The Western Lancet.*

As a comprehensive manual for students, or a work of reference for practitioners, we only speak with common justice when we say that it surpasses any other that has ever issued on the same subject from the British press.—*The Dublin Quarterly Journal.*

DICKSON (S. H.), M. D.,

Professor of Institutes and Practice of Medicine in the Medical College of South Carolina.

ELEMENTS OF MEDICINE; a Compendious View of Pathology and Therapeutics, or the History and Treatment of Diseases. In one large and handsome octavo volume, of 750 pages, leather (*Now Ready.*) \$3 75.

As an American text-book on the Practice of Medicine for the student, and as a condensed work of reference for the practitioner, this volume will have strong claims on the attention of the profession. Few physicians have had wider opportunities than the author for observation and experience, and few perhaps have used them better. As the result of a life of study and practice, therefore, the present volume will doubtless be received with the welcome it deserves.

This book is eminently what it professes to be; a distinguished merit in these days. Designed for "Teachers and Students of Medicine," and admirably suited to their wants, we think it will be received, on its own merits, with a hearty welcome.—*Boston Med. and Surg. Journal.*

Indited by one of the most accomplished writers of our country, as well as by one who has long held a high position among teachers and practitioners of medicine, this work is entitled to patronage and careful study. The learned author has endeavored to condense in this volume most of the practical matter contained in his former productions, so as to adapt it to the use of those who have not time to devote to more extensive works.—*Southern Med. and Surg. Journal.*

We can strongly recommend Dr. Dickson's work to our readers as one of interest and practical utility, well deserving of a place in their libraries as a book of reference; and we especially commend the first part as presenting an admirable outline of the principles of medicine.—*Dublin Quarterly Journal*, Feb. 1856

This volume, while as its title denotes it is a compendious view, is also a comprehensive system of practice, perspicuously and pleasantly written, and admirably suited to engage the interest, and instruct the reader.—*Peninsular Journal of Medicine*, Jan. 1856.

Prof. Dickson's work supplies, to a great extent, a desideratum long felt in American medicine.—*N. O. Med. and Surg. Journal.*

Estimating this work according to the purpose for which it is designed, we must think highly of its merits, and we have no hesitation in predicting for it a favorable reception by both students and teachers.

Not professing to be a complete and comprehensive treatise, it will not be found full in detail, nor filled with discussions of theories and opinions, but embracing all that is essential in theory and practice, it is admirably adapted to the wants of the American student. Avoiding all that is uncertain, it presents more clearly to the mind of the reader that which is established and verified by experience. The varied and extensive reading of the author is conspicuously apparent, and all the recent improvements and discoveries in therapeutics and pathology are chronicled in its pages.—*Charleston Med. Journal.*

In the first part of the work the subject of general pathology is presented in outline, giving a beautiful picture of its distinguishing features, and throughout the succeeding chapters we find that he has kept scrupulously within the bounds of sound reasoning and legitimate deduction. Upon the whole, we do not hesitate to pronounce it a superior work in its class, and that Dr. Dickson merits a place in the first rank of American writers.—*Western Lancet.*

DRUITT (ROBERT), M. R. C. S., &c.

THE PRINCIPLES AND PRACTICE OF MODERN SURGERY. A new American, from the improved London edition. Edited by F. W. SARGENT, M. D., author of "Minor Surgery," &c. Illustrated with one hundred and ninety-three wood-engravings. In one very handsomely printed octavo volume, leather, of 576 large pages. \$3 00.

Dr. Druiitt's researches into the literature of his subject have been not only extensive, but well directed; the most discordant authors are fairly and impartially quoted, and, while due credit is given to each, their respective merits are weighed with an unprejudiced hand. The grain of wheat is preserved, and the chaff is unmercifully stripped off. The arrangement is simple and philosophical, and the style, though clear and interesting, is so precise, that the book contains more information condensed into a few words than any other surgical work with which we are acquainted.—*London Medical Times and Gazette.*

No work, in our opinion, equals it in presenting so much valuable surgical matter in so small a compass.—*St. Louis Med. and Surgical Journal.*

Druiitt's Surgery is too well known to the American medical profession to require its announcement anywhere. Probably no work of the kind has ever been more cordially received and extensively circulated than this. The fact that it comprehends in a comparatively small compass, all the essential elements of theoretical and practical Surgery—that it is found to contain reliable and authentic information on the nature and treatment of nearly all surgical affections—is a sufficient reason for the liberal patronage it has obtained. The editor, Dr. F. W. Sargent, has contributed much to enhance the value of the work, by such American improvements as are calculated more perfectly to adapt it to our own views and practice in this country. It abounds everywhere with spirited and life-like illustrations, which to the young surgeon, especially, are of no minor consideration. Every medical man frequently needs just such a work as this, for immediate reference in moments of sudden emergency, when he has not time to consult more elaborate treatises.—*The Ohio Medical and Surgical Journal.*

The author has evidently ransacked every standard treatise of ancient and modern times, and all that

is really practically useful at the bedside will be found in a form at once clear, distinct, and interesting.—*Edinburgh Monthly Medical Journal.*

Druiitt's work, condensed, systematic, lucid, and practical as it is, beyond most works on Surgery accessible to the American student, has had much currency in this country, and under its present auspices promises to rise to yet higher favor.—*The Western Journal of Medicine and Surgery.*

The most accurate and ample resumé of the present state of Surgery that we are acquainted with.—*Dublin Medical Journal.*

A better book on the principles and practice of Surgery as now understood in England and America, has not been given to the profession.—*Boston Medical and Surgical Journal.*

An unsurpassable compendium, not only of Surgical, but of Medical Practice.—*London Medical Gazette.*

This work merits our warmest commendations, and we strongly recommend it to young surgeons as an admirable digest of the principles and practice of modern Surgery.—*Medical Gazette.*

It may be said with truth that the work of Mr. Druiitt affords a complete, though brief and condensed view, of the entire field of modern surgery. We know of no work on the same subject having the appearance of a manual, which includes so many topics of interest to the surgeon; and the terse manner in which each has been treated evinces a most enviable quality of mind on the part of the author, who seems to have an innate power of searching out and grasping the leading facts and features of the most elaborate productions of the pen. It is a useful handbook for the practitioner, and we should deem a teacher of surgery unpardonable who did not recommend it to his pupils. In our own opinion, it is admirably adapted to the wants of the student.—*Provincial Medical and Surgical Journal.*

DUNGLISON, FORBES, TWEEDIE, AND CONOLLY.

THE CYCLOPÆDIA OF PRACTICAL MEDICINE: comprising Treatises on the Nature and Treatment of Diseases, Materia Medica, and Therapeutics, Diseases of Women and Children, Medical Jurisprudence, &c. &c. In four large super-royal octavo volumes, of 3254 double-columned pages, strongly and handsomely bound, with raised bands. \$12 00.

. This work contains no less than four hundred and eighteen distinct treatises, contributed by sixty-eight distinguished physicians, rendering it a complete library of reference for the country practitioner.

The most complete work on Practical Medicine extant; or, at least, in our language.—*Buffalo Medical and Surgical Journal.*

For reference, it is above all price to every practitioner.—*Western Lancet.*

One of the most valuable medical publications of the day—as a work of reference it is invaluable.—*Western Journal of Medicine and Surgery.*

It has been to us, both as learner and teacher, a work for ready and frequent reference, one in which modern English medicine is exhibited in the most advantageous light.—*Medical Examiner.*

We rejoice that this work is to be placed within the reach of the profession in this country, it being unquestionably one of very great value to the prac-

titioner. This estimate of it has not been formed from a hasty examination, but after an intimate acquaintance derived from frequent consultation of it during the past nine or ten years. The editors are practitioners of established reputation, and the list of contributors embraces many of the most eminent professors and teachers of London, Edinburgh, Dublin, and Glasgow. It is, indeed, the great merit of this work that the principal articles have been furnished by practitioners who have not only devoted especial attention to the diseases about which they have written, but have also enjoyed opportunities for an extensive practical acquaintance with them, and whose reputation carries the assurance of their competency justly to appreciate the opinions of others, while it stamps their own doctrines with high and just authority.—*American Medical Journ.*

DEWEES'S COMPREHENSIVE SYSTEM OF MIDWIFERY. Illustrated by occasional cases and many engravings. Twelfth edition, with the author's last improvements and corrections. In one octavo volume, extra cloth, of 600 pages. \$3 20.

DEWEES'S TREATISE ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN. Tenth edition. In one volume, octavo, extra cloth, 518 pages. \$2 80.

DEWEES'S TREATISE ON THE DISEASES OF FEMALES. Tenth edition. In one volume, octavo, extra cloth, 532 pages, with plates. \$3 00.

DANA ON ZOOPHYTES AND CORALS. In one volume, imperial quarto, extra cloth, with woodcuts. \$15 00. Also, **AN ATLAS**, in one volume, imperial folio, with sixty-one magnificent colored plates. Bound in half morocco. \$30 00.

DE LA BECHE'S GEOLOGICAL OBSERVER. In one very large and handsome octavo volume, extra cloth, of 700 pages, with 300 woodcuts. \$4 00.

FRICK ON RENAL AFFECTIONS; their Diagnosis and Pathology. With illustrations. One volume, royal 12mo., extra cloth. 75 cents.

DUNGLISON (ROBLEY), M. D.,

Professor of Institutes of Medicine in the Jefferson Medical College, Philadelphia.

MEDICAL LEXICON; a Dictionary of Medical Science, containing a concise

Explanation of the various Subjects and Terms of Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Obstetrics, Medical Jurisprudence, &c. With the French and other Synonyms; Notices of Climate and of celebrated Mineral Waters; Formulæ for various Official, Empirical, and Dietetic Preparations, etc. THIRTEENTH EDITION, revised, is now ready. In one very thick octavo volume, of over nine hundred large double-columned pages, strongly bound in leather, with raised bands. \$4 00.

Every successive edition of this work bears the marks of the industry of the author, and of his determination to keep it fully on a level with the most advanced state of medical science. Thus nearly FIFTEEN THOUSAND WORDS have been added to it within the last few years. As a complete Medical Dictionary, therefore, embracing over FIFTY THOUSAND DEFINITIONS, in all the branches of the science, it is presented as meriting a continuance of the great favor and popularity which have carried it, within no very long space of time, to a thirteenth edition.

Every precaution has been taken in the preparation of the present volume, to render its mechanical execution and typographical accuracy worthy of its extended reputation and universal use. The very extensive additions have been accommodated, without materially increasing the bulk of the volume by the employment of a small but exceedingly clear type, cast for this purpose. The press has been watched with great care, and every effort used to insure the verbal accuracy so necessary to a work of this nature. The whole is printed on fine white paper; and, while thus exhibiting in every respect so great an improvement over former issues, it is presented at the original exceedingly low price.

We welcome it cordially; it is an admirable work, and indispensable to all literary medical men. The labor which has been bestowed upon it is something prodigious. The work, however, has now been done, and we are happy in the thought that no human being will have again to undertake the same gigantic task. Revised and corrected from time to time, Dr. Dunglison's "Medical Lexicon" will last for centuries.—*British and Foreign Med.-Chirurg. Review.*

The fact that this excellent and learned work has passed through eight editions, and that a ninth is rendered necessary by the demands of the public, affords a sufficient evidence of the general appreciation of Dr. Dunglison's labors by the medical profession in England and America. It is a book which will be of great service to the student, in teaching him the meaning of all the technical terms used in medicine, and will be of no less use to the practitioner who desires to keep himself on a level with the advance of medical science.—*London Medical Times and Gazette.*

In taking leave of our author, we feel compelled to confess that his work bears evidence of almost incredible labor having been bestowed upon its composition.—*Edinburgh Journal of Med. Science.*

A miracle of labor and industry in one who has written able and voluminous works on nearly every branch of medical science. There could be no more useful book to the student or practitioner, in the present advancing age, than one in which would be found, in addition to the ordinary meaning and derivation of medical terms—so many of which are of modern introduction—concise descriptions of their explanation and employment; and all this and much more is contained in the volume before us. It is therefore almost as indispensable to the other learned professions as to our own. In fact, to all who may have occasion to ascertain the meaning of any word belonging to the many branches of medicine. From a careful examination of the present edition, we can vouch for its accuracy, and for its being brought quite up to the date of publication; the author states in his preface that he has added to it about four thousand terms, which are not to be found in the preceding ones.—*Dublin Quarterly Journal of Medical Science.*

On the appearance of the last edition of this valuable work, we directed the attention of our

readers to its peculiar merits; and we need do little more than state, in reference to the present issue, that, notwithstanding the large additions previously made to it, no fewer than four thousand terms, not to be found in the preceding edition, are contained in the volume before us.—Whilst it is a wonderful monument of its author's erudition and industry, it is also a work of great practical utility, as we can testify from our own experience; for we keep it constantly within our reach, and make very frequent reference to it, nearly always finding in it the information we seek.—*British and Foreign Med.-Chirurg. Review.*

It has the rare merit that it certainly has no rival in the English language for accuracy and extent of references. The terms generally include short physiological and pathological descriptions, so that, as the author justly observes, the reader does not possess in this work a mere dictionary, but a book, which, while it instructs him in medical etymology, furnishes him with a large amount of useful information. The author's labors have been properly appreciated by his own countrymen; and we can only confirm their judgment, by recommending this most useful volume to the notice of our cisatlantic readers. No medical library will be complete without it.—*London Med. Gazette.*

It is certainly more complete and comprehensive than any with which we are acquainted in the English language. Few, in fact, could be found better qualified than Dr. Dunglison for the production of such a work. Learned, industrious, persevering, and accurate, he brings to the task all the peculiar talents necessary for its successful performance; while, at the same time, his familiarity with the writings of the ancient and modern "masters of our art," renders him skilful to note the exact usage of the several terms of science, and the various modifications which medical terminology has undergone with the change of theories or the progress of improvement.—*American Journal of the Medical Sciences.*

One of the most complete and copious known to the cultivators of medical science.—*Boston Med. Journal.*

The most comprehensive and best English Dictionary of medical terms extant.—*Buffalo Medical Journal.*

BY THE SAME AUTHOR.

THE PRACTICE OF MEDICINE. A Treatise on Special Pathology and Therapeutics. Third Edition. In two large octavo volumes, leather, of 1,500 pages. \$6 25.

Upon every topic embraced in the work the latest information will be found carefully posted up.—*Medical Examiner.*

The student of medicine will find, in these two elegant volumes, a mine of facts, a gathering of precepts and advice from the world of experience, that will nerve him with courage, and faithfully direct him in his efforts to relieve the physical suf-

ferings of the race.—*Boston Medical and Surgical Journal.*

It is certainly the most complete treatise of which we have any knowledge.—*Western Journal of Medicine and Surgery.*

One of the most elaborate treatises of the kind we have.—*Southern Med. and Surg. Journal.*

DUNGLISON (ROBLEY), M. D.,

Professor of Institutes of Medicine in the Jefferson Medical College, Philadelphia.

HUMAN PHYSIOLOGY. Eighth edition. Thoroughly revised and extensively modified and enlarged, with five hundred and thirty-two illustrations. In two large and handsomely printed octavo volumes, leather, of about 1500 pages. (*Now Ready*, 1856.) \$7 00.

In revising this work for its eighth appearance, the author has spared no labor to render it worthy a continuance of the very great favor which has been extended to it by the profession. The whole contents have been rearranged, and to a great extent remodelled; the investigations which of late years have been so numerous and so important, have been carefully examined and incorporated, and the work in every respect has been brought up to a level with the present state of the subject. The object of the author has been to render it a concise but comprehensive treatise, containing the whole body of physiological science, to which the student and man of science can at all times refer with the certainty of finding whatever they are in search of, fully presented in all its aspects; and on no former edition has the author bestowed more labor to secure this result.

A similar improvement will be found in the typographical execution of the volumes, which, in this respect, are superior to their predecessors. A large number of additional wood-cuts have been introduced, and the series of illustrations has been greatly modified by the substitution of many new ones for such as were not deemed satisfactory. By an enlargement of the page, these very considerable additions have been accommodated without increasing the size of the volumes to an extent to render them unwieldy.

It has long since taken rank as one of the medical classics of our language. To say that it is by far the best text-book of physiology ever published in this country, is but echoing the general testimony of the profession.—*N. Y. Journal of Medicine*.

There is no single book we would recommend to the student or physician, with greater confidence than the present, because in it will be found a mirror of almost every standard physiological work of the day. We most cordially recommend the work to every member of the profession, and no student

should be without it. It is the completest work on Physiology in the English language, and is highly creditable to the author and publishers.—*Canadian Medical Journal*.

The most complete and satisfactory system of Physiology in the English language.—*Amer. Med. Journal*.

The best work of the kind in the English language.—*Silliman's Journal*.

The most full and complete system of Physiology in our language.—*Western Lancet*.

BY THE SAME AUTHOR.

GENERAL THERAPEUTICS AND MATERIA MEDICA; adapted for a Medical Text-book. Fifth edition, much improved. With one hundred and eighty-seven illustrations. In two large and handsomely printed octavo vols., leather, of about 1100 pages. \$6 00.

In this work of Dr. Dunglison, we recognize the same untiring industry in the collection and embodying of facts on the several subjects of which he treats, that has heretofore distinguished him, and we cheerfully point to these volumes, as two of the most interesting that we know of. In noticing the additions to this, the fourth edition, there is very little in the periodical or annual literature of the profession, published in the interval which has elapsed since the issue of the first, that has escaped the careful search of the author. As a book for reference, it is invaluable.—*Charleston Med. Journal and Review*.

It may be said to be the work now upon the subjects upon which it treats.—*Western Lancet*.

As a text-book for students, for whom it is particularly designed, we know of none superior to it.—*St. Louis Medical and Surgical Journal*.

It purports to be a new edition, but it is rather a new book, so greatly has it been improved, both in the amount and quality of the matter which it contains.—*N. O. Medical and Surgical Journal*.

We bespeak for this edition, from the profession, an increase of patronage over any of its former ones, on account of its increased merit.—*N. Y. Journal of Medicine*.

We consider this work unequalled.—*Boston Med. and Surg. Journal*.

BY THE SAME AUTHOR. (*A new Edition*.)

NEW REMEDIES, WITH FORMULÆ FOR THEIR PREPARATION AND ADMINISTRATION. Seventh edition, with extensive Additions. In one very large octavo volume, leather, of 770 pages. (*Now Ready*, May, 1856.) \$3 75.

Another edition of the "New Remedies" having been called for, the author has endeavored to add everything of moment that has appeared since the publication of the last edition.

The chief remedial means which have obtained a place, for the first time, in this volume, either owing to their having been recently introduced into pharmacology, or to their having received novel applications—and which, consequently, belong to the category of "New Remedies"—are the following:—

Apioi, Caffein, Carbazotic acid, Cauterization and catheterism of the larynx and trachea, Cedron, Cerium, Chloride of bromine, Chloride of iron, Chloride of sodium, Cinchonine, Cod-liver oil, Congelation, Eau de Pagliari, Galvanic cautery, Hydriodic ether, Hyposulphite of soda and silver, Inunction, Iodide of sodium, Nickel, Permanganate of potassa, Phosphate of lime, Pumpkin, Quinidia, Rennet, Saccharine carbonate of iron and manganese, Santonin, Tellurium, and Traumaticine.

The articles treated of in the former editions will be found to have undergone considerable expansion in this, in order that the author might be enabled to introduce, as far as practicable, the results of the subsequent experience of others, as well as of his own observation and reflection; and to make the work still more deserving of the extended circulation with which the preceding editions have been favored by the profession. By an enlargement of the page, the numerous additions have been incorporated without greatly increasing the bulk of the volume.—*Preface*.

One of the most useful of the author's works.—*Southern Medical and Surgical Journal*.

This elaborate and useful volume should be found in every medical library, for as a book of reference, for physicians, it is unsurpassed by any other work in existence, and the double index for diseases and for remedies, will be found greatly to enhance its value.—*New York Med. Gazette*.

The great learning of the author, and his remarkable industry in pushing his researches into every source whence information is derivable, has enabled him to throw together an extensive mass of facts and statements, accompanied by full reference to authorities; which last feature renders the work practically valuable to investigators who desire to examine the original papers.—*The American Journal of Pharmacy*.

ERICHSEN (JOHN),

Professor of Surgery in University College, London, &c.

THE SCIENCE AND ART OF SURGERY; BEING A TREATISE ON SURGICAL INJURIES, DISEASES, AND OPERATIONS. Edited by JOHN H. BRINTON, M. D. Illustrated with three hundred and eleven engravings on wood. In one large and handsome octavo volume, of over nine hundred closely printed pages, leather, raised bands. \$4 25.

It is, in our humble judgment, decidedly the best book of the kind in the English language. Strange that just such books are not oftener produced by public teachers of surgery in this country and Great Britain. Indeed, it is a matter of great astonishment, but no less true than astonishing, that of the many works on surgery republished in this country within the last fifteen or twenty years as text-books for medical students, this is the only one that even approximates to the fulfilment of the peculiar wants of young men just entering upon the study of this branch of the profession.—*Western Jour. of Med. and Surgery.*

Its value is greatly enhanced by a very copious well-arranged index. We regard this as one of the most valuable contributions to modern surgery. To one entering his novitiate of practice, we regard it the most serviceable guide which he can consult. He will find a fulness of detail leading him through every step of the operation, and not deserting him until the final issue of the case is decided. For the same reason we recommend it to those whose routine of practice lies in such parts of the country that they must

rarely encounter cases requiring surgical management.—*Stethoscope.*

Embracing, as will be perceived, the whole surgical domain, and each division of itself almost complete and perfect, each chapter full and explicit, each subject faithfully exhibited, we can only express our estimate of it in the aggregate. We consider it an excellent contribution to surgery, as probably the best single volume now extant on the subject, and with great pleasure we add it to our text-books.—*Nashville Journal of Medicine and Surgery.*

Prof. Erichsen's work, for its size, has not been surpassed; his nine hundred and eight pages, profusely illustrated, are rich in physiological, pathological, and operative suggestions, doctrines, details, and processes; and will prove a reliable resource for information, both to physician and surgeon, in the hour of peril.—*N. O. Med. and Surg. Journal.*

We are acquainted with no other work wherein so much good sense, sound principle, and practical inferences, stamp every page.—*American Lancet.*

ELLIS (BENJAMIN), M. D.

THE MEDICAL FORMULARY: being a Collection of Prescriptions, derived from the writings and practice of many of the most eminent physicians of America and Europe. Together with the usual Dietetic Preparations and Antidotes for Poisons. To which is added an Appendix, on the Endermic use of Medicines, and on the use of Ether and Chloroform. The whole accompanied with a few brief Pharmaceutical and Medical Observations. Tenth edition, revised and much extended by ROBERT P. THOMAS, M. D., Professor of Materia Medica in the Philadelphia College of Pharmacy. In one neat octavo volume, extra cloth, of 296 pages. (*Lately Issued.*) \$1 75.

After an examination of the new matter and the alterations, we believe the reputation of the work built up by the author, and the late distinguished editor, will continue to flourish under the auspices of the present editor, who has the industry and accuracy, and we would say, conscientiousness requisite for the responsible task.—*Am. Jour. of Pharm.*

It will prove particularly useful to students and young practitioners, as the most important prescriptions employed in modern practice, which lie scattered through our medical literature, are here collected and conveniently arranged for reference.—*Charleston Med. Journal and Review.*

FOWNES (GEORGE), PH. D., &c.

ELEMENTARY CHEMISTRY; Theoretical and Practical. With numerous illustrations. A new American, from the last and revised London edition. Edited, with Additions, by ROBERT BRIDGES, M. D. In one large royal 12mo. volume, of over 550 pages, with 181 wood-cuts. (*Lately Issued.*) In leather, \$1 50; extra cloth, \$1 35.

We know of no better text-book, especially in the difficult department of organic chemistry, upon which it is particularly full and satisfactory. We would recommend it to preceptors as a capital "office book" for their students who are beginners in Chemistry. It is copiously illustrated with excellent wood-cuts, and altogether admirably "got up."—*N. J. Medical Reporter.*

A standard manual, which has long enjoyed the reputation of embodying much knowledge in a small space. The author has achieved the difficult task of condensation with masterly tact. His book is concise without being dry, and brief without being too dogmatical or general.—*Virginia Med. and Surgical Journal.*

The work of Dr. Fownes has long been before the public, and its merits have been fully appreciated as the best text-book on chemistry now in existence. We do not, of course, place it in a rank superior to the works of Brande, Graham, Turner, Gregory, or Gmelin, but we say that, as a work for students, it is preferable to any of them.—*London Journal of Medicine.*

A work well adapted to the wants of the student. It is an excellent exposition of the chief doctrines and facts of modern chemistry. The size of the work, and still more the condensed yet perspicuous style in which it is written, absolve it from the charges very properly urged against most manuals termed popular.—*Edinburgh Journal of Medical Science.*

FERGUSON (WILLIAM), F. R. S.,

Professor of Surgery in King's College, London, &c.

A SYSTEM OF PRACTICAL SURGERY. Fourth American, from the third and enlarged London edition. In one large and beautifully printed octavo volume, of about 700 pages, with 393 handsome illustrations, leather. \$3 00.

The most important subjects in connection with practical surgery which have been more recently brought under the notice of, and discussed by, the surgeons of Great Britain, are fully and dispassionately considered by Mr. Ferguson, and that which was before wanting has now been supplied, so that we can now look upon it as a work on practical surgery instead of one on operative surgery alone. *Medical Times and Gazette.*

No work was ever written which more nearly comprehended the necessities of the student and

practitioner, and was more carefully arranged to that single purpose than this.—*N. Y. Med. Journal.*

The addition of many new pages makes this work more than ever indispensable to the student and practitioner.—*Ranking's Abstract.*

Among the numerous works upon surgery published of late years, we know of none we value more highly than the one before us. It is perhaps the very best we have for a text-book and for ordinary reference, being concise and eminently practical.—*Southern Med. and Surg. Journal.*

FLINT (AUSTIN), M. D.,

Professor of the Theory and Practice of Medicine in the University of Louisville, &c.

(An Important New Work.)

PHYSICAL EXPLORATION AND DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS. In one large and handsome octavo volume, extra cloth, 636 pages. (Now Ready.) \$3 00.

The author has aimed in the present work to supply a vacancy in medical literature, viz: "a work limited to diseases affecting the respiratory organs, treating *in extenso* and almost exclusively of the principles and practice of physical exploration, as applied to the diagnosis of those affections." The intricacy and importance of the subject demand a fuller and more detailed exposition than has been accorded to it in any volume as yet accessible to the American profession; while the high reputation which the author has acquired by his researches in kindred topics sufficiently manifests his ability to render the present work a text-book of great practical utility for the student, and a source to which the practitioner can at all times refer with certainty.

A very condensed summary of the contents is subjoined.

CONTENTS.

INTRODUCTION.

SECTION I. Preliminary points pertaining to the Anatomy and Physiology of the Respiratory Apparatus. SECTION II. Topographical Divisions of the Chest.

PART I.

PHYSICAL EXPLORATION OF THE CHEST.

CHAP. I. Definitions—Different Methods of Explorations—General Remarks. CHAP. II. Percussion. CHAP. III. Auscultation. CHAP. IV. Inspection. CHAP. V. Mensuration. CHAP. VI. Palpation. CHAP. VII. Succession. CHAP. VIII. Recapitulatory Enumeration of the Physical Signs furnished by the several methods of Exploration. CHAP. IX. Correlation of Physical Signs.

PART II.

DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS.

CHAP. I. Bronchitis, Pulmonary or Bronchial Catarrh. CHAP. II. Dilatation and Contraction of the Bronchial Tubes—Pertussis—Asthma. CHAP. III. Pneumonitis—Imperfect Expansion (Atelectasis) and Collapse. CHAP. IV. Emphysema. CHAP. V. Pulmonary Tuberculosis—Bronchial Phthisis. CHAP. VI. Pulmonary Edema—Gangrene of the Lungs—Pulmonary Apoplexy—Cancer of the Lungs—Cancer in the Mediastinum. CHAP. VII. Acute Pleuritis—Chronic Pleuritis—Empyema—Hydrothorax—Pneumothorax—Pneumo-hydrothorax—Pleuralgia—Diaphragmatic Hernia. CHAP. VIII. Diseases affecting the Trachea and Larynx—Foreign Bodies in the Air-passages. APPENDIX. On the Pitch of the Whispering Soudle over Pulmonary Excavations.

GRAHAM (THOMAS), F. R. S.,

Professor of Chemistry in University College, London, &c.

THE ELEMENTS OF CHEMISTRY. Including the application of the Science to the Arts. With numerous illustrations. With Notes and Additions, by ROBERT BRIDGES, M. D., &c. &c. Second American, from the second and enlarged London edition.

PART I. (Lately Issued) large 8vo., 430 pages, 185 illustrations. \$1 50.

PART II. (Preparing) to match.

GRIFFITH (ROBERT E.), M. D., &c.

A UNIVERSAL FORMULARY, containing the methods of Preparing and Administering Official and other Medicines. The whole adapted to Physicians and Pharmacæutists. SECOND EDITION, thoroughly revised, with numerous additions, by ROBERT P. THOMAS, M. D., Professor of Materia Medica in the Philadelphia College of Pharmacy. In one large and handsome octavo volume, extra cloth, of over 600 pages, double columns. (Just Issued.) \$3 00; or bound in sheep, \$3 25.

It was a work requiring much perseverance, and when published was looked upon as by far the best work of its kind that had issued from the American press. Prof Thomas has certainly "improved" as well as added to this Formulary, and has rendered it additionally deserving of the confidence of pharmacæutists and physicians.—*Am. Journal of Pharmacy*.

We are happy to announce a new and improved edition of this, one of the most valuable and useful works that have emanated from an American pen. It would do credit to any country, and will be found of daily usefulness to practitioners of medicine; it is better adapted to their purposes than the dispensatories.—*Southern Med. and Surg. Journal*.

A new edition of this well-known work, edited by R. P. Thomas, M. D., affords occasion for renewing our commendation of so useful a handbook, which ought to be universally studied by medical men of every class, and made use of by way of reference by office pupils, as a standard authority. It has been much enlarged, and now condenses a vast amount of useful and necessary knowledge in small compass. The more of such books the better for the profession and the public.—*N. Y. Med. Gazette*.

It is one of the most useful books a country practi-

tioner can possibly have in his possession.—*Medical Chronicle*.

The amount of useful, every-day matter, for a practicing physician, is really immense.—*Boston Med. and Surg. Journal*.

This is a work of six hundred and fifty-one pages, embracing all on the subject of preparing and administering medicines that can be desired by the physician and pharmacæutist.—*Western Lancet*.

In short, it is a full and complete work of the kind, and should be in the hands of every physician and apothecary. *O Med. and Surg Journal*

We predict a great sale for this work, and we especially recommend it to all medical teachers.—*Richmond Stethoscope*.

This edition of Dr Griffith's work has been greatly improved by the revision and ample additions of Dr. Thomas, and is now, we believe, one of the most complete works of its kind in any language. The additions amount to about seventy pages, and no effort has been spared to include in them all the recent improvements which have been published in medical journals, and systematic treatises. A work of this kind appears to us indispensable to the physician, and there is none we can more cordially recommend.—*N. Y. Journal of Medicine*.

BY THE SAME AUTHOR.

MEDICAL BOTANY; or, a Description of all the more important Plants used in Medicine, and of their Properties, Uses, and Modes of Administration. In one large octavo volume, extra cloth, of 704 pages, handsomely printed, with nearly 350 illustrations on wood. \$3 00.

GROSS (SAMUEL D.), M. D.,

Professor of Surgery in the Jefferson Medical College of Philadelphia, &c.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Second Edition, revised and much enlarged, with one hundred and eighty-four illustrations. In one large and very handsome octavo volume, of over nine hundred pages. (*Just Issued.*) In leather, raised bands, \$5 25; extra cloth, \$4 75.

The author has availed himself of the opportunity afforded by a call for a new edition of this work, to thoroughly revise and render it in every respect worthy, so far as in his power, of the very flattering reception which has been accorded to it by the profession. The new matter thus added amounts to almost one-third of the original work, while the number of illustrations has been nearly doubled. These additions pervade every portion of the work, which thus has rather the aspect of a new treatise than a new edition. In its present improved form, therefore, it may confidently be presented as a complete and reliable storehouse of information on this important class of diseases, and as in every way fitted to maintain the position which it has acquired in Europe and in this country, as the standard of authority on the subjects treated of.

A volume replete with truths and principles of the utmost value in the investigation of these diseases.—*American Medical Journal.*

On the appearance of the first edition of this work, the leading English medical review predicted that it would have a "permanent place in the literature of surgery worthy to rank with the best works of the present age." This prediction has been amply fulfilled. Dr. Gross's treatise has been found to supply completely the want which has been felt ever since the elevation of surgery to the rank of a science, of a good practical treatise on the diseases of the bladder and its accessory organs. Philosophical in its design, methodical in its arrangement, ample and sound in its practical details, it may in truth be said to leave scarcely anything to be desired on so important a subject, and with the additions and modifications resulting from future discoveries and im-

provements, it will probably remain one of the most valuable works on this subject so long as the science of medicine shall exist.—*Boston Med. and Surg. Journal.*

Dr. Gross has brought all his learning, experience, tact, and judgment to the task, and has produced a work worthy of his high reputation. We feel perfectly safe in recommending it to our readers as a monograph unequalled in interest and practical value by any other on the subject in our language.—*Western Journal of Med. and Surg.*

Whoever will peruse the vast amount of valuable practical information it contains, and which we have been unable even to notice, will, we think, agree with us, that there is no work in the English language which can make any just pretensions to be its equal.—*N. Y. Journal of Medicine.*

BY THE SAME AUTHOR. (*Just Issued.*)

A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR-PASSAGES. In one handsome octavo volume, extra cloth, with illustrations. pp. 468. \$2 75.

A very elaborate work. It is a complete summary of the whole subject, and will be a useful book of reference.—*British and Foreign Medico-Chirurg. Review.*

A highly valuable book of reference on a most important subject in the practice of medicine. We

conclude by recommending it to our readers, fully persuaded that its perusal will afford them much practical information well conveyed, evidently derived from considerable experience and deduced from an ample collection of facts.—*Dublin Quarterly Journal*, May, 1855.

BY THE SAME AUTHOR. (*Preparing.*)

A SYSTEM OF SURGERY; Diagnostic, Pathological, Therapeutic, and Operative. With very numerous engravings on wood.

BY THE SAME AUTHOR. (*Preparing.*)

ELEMENTS OF PATHOLOGICAL ANATOMY; illustrated by several hundred wood-cuts. New edition, thoroughly revised. In one very large and handsome octavo volume.

GLUGE (GOTTLIEB), M. D.,

Professor of Physiology and Pathological Anatomy in the University of Brussels, &c.

AN ATLAS OF PATHOLOGICAL HISTOLOGY. Translated, with Notes and Additions, by JOSEPH LEIDY, M. D., Professor of Anatomy in the University of Pennsylvania. In one volume, very large imperial quarto, extra cloth, with 320 figures, plain and colored, on twelve copperplates. \$5 00.

GARDNER'S MEDICAL CHEMISTRY, for the use of Students and the Profession. In one royal 12mo. vol., ex. cloth, pp. 396, with illustrations. \$1 00.

HARRISON'S ESSAY TOWARDS A CORRECT THEORY OF THE NERVOUS SYSTEM. In one octavo volume, leather, 292 pages. \$1 50.

HUGHES' CLINICAL INTRODUCTION TO THE PRACTICE OF AUSCULTATION AND OTHER MODES OF PHYSICAL DIAGNOSIS, IN DISEASES OF THE LUNGS AND HEART. Second American, from the second London edition. 1 vol. royal 12mo., ex. cloth, pp. 304. \$1 00.

HAMILTON (FRANK H.), M. D.,

Professor of Surgery, in Buffalo Medical College, &c.

A TREATISE ON FRACTURES AND DISLOCATIONS. In one handsome octavo volume, with numerous illustrations. (*Preparing.*)

The numerous improvements which this important branch of surgery has received from the skill and ingenuity of American surgeons, renders particularly appropriate and valuable a complete and systematic original work on the subject. The essays which Professor Hamilton has published on kindred topics are already widely and favorably known, and give earnest that his forthcoming work will prove indispensable, both as a text-book for the student, and as a guide for the practitioner.

HOBLYN (RICHARD D.), M. D.

A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. By RICHARD D. HOBLYN, A. M., &c. A new American from the last London edition. Revised, with numerous Additions, by ISAAC HAYS, M. D., editor of the "American Journal of the Medical Sciences." In one large royal 12mo. volume, leather, of over 500 double columned pages. (*Now Ready*, 1856.) \$1 50.

If the frequency with which we have referred to this volume since its reception from the publisher, two or three weeks ago, be any criterion for the future, the binding will soon have to be renewed, even with careful handling. We find that Dr. Hays has done the profession great service by his careful and industrious labors. The Dictionary has thus become eminently suited to our medical brethren in this country. The additions by Dr. Hays are in brackets, and we believe there is not a single page but bears these insignia; in every instance which we have thus far noticed, the additions are really needed and exceedingly valuable. We heartily commend the work to all who wish to be *au courant* in medical terminology.—*Boston Med. and Surg. Journal*.

To both practitioner and student, we recommend this dictionary as being convenient in size, accurate in definition, and sufficiently full and complete for ordinary consultation.—*Charleston Med. Journ. and Review*.

Admirably calculated to meet the wants of the practitioner or student, who has neither the means

nor desire to procure a larger work.—*American Lancet*.

Hoblyn has always been a favorite dictionary, and in its present enlarged and improved form will give greater satisfaction than ever. The American editor, Dr. Hays, has made many very valuable additions.—*N. J. Med. Reporter*.

To supply the want of the medical reader arising from this cause, we know of no dictionary better arranged and adapted than the one bearing the above title. It is not encumbered with the obsolete terms of a bygone age, but it contains all that are now in use; embracing every department of medical science down to the very latest date. The volume is of a convenient size to be used by the medical student, and yet large enough to make a respectable appearance in the library of a physician.—*Western Lancet*.

Hoblyn's Dictionary has long been a favorite with us. It is the best book of definitions we have, and ought always to be upon the student's table.—*Southern Med. and Surg. Journal*.

HOLLAND (SIR HENRY), BART., M. D., F. R. S.,


Physician in Ordinary to the Queen of England, &c.

MEDICAL NOTES AND REFLECTIONS. From the third London edition.

In one handsome octavo volume, extra cloth. (*Now Ready*.) \$3 00.

As the work of a thoughtful and observant physician, embodying the results of forty years' active professional experience, on topics of the highest interest, this volume is commended to the American practitioner as well worthy his attention. Few will rise from its perusal without feeling their convictions strengthened, and armed with new weapons for the daily struggle with disease.

HUNTER (JOHN).

TREATISE ON THE VENEREAL DISEASE. With copious Additions, by DR. PH. RICORD, Surgeon to the Venereal Hospital of Paris. Edited, with additional Notes, by F. J. BUMSTEAD, M. D. In one octavo volume, with plates. \$3 25.  See RICORD.

Also, HUNTER'S COMPLETE WORKS, with Memoir, Notes, &c. &c. In four neat octavo volumes, leather, with plates. \$10 00.

HORNER (WILLIAM E.), M. D.,

Professor of Anatomy in the University of Pennsylvania.

SPECIAL ANATOMY AND HISTOLOGY. Eighth edition. Extensively revised and modified. In two large octavo volumes, extra cloth, of more than one thousand pages, handsomely printed, with over three hundred illustrations. \$6 00.

This edition enjoyed a thorough and laborious revision on the part of the author shortly before his death, with the view of bringing it fully up to the existing state of knowledge on the subject of general and special anatomy. To adapt it more perfectly to the wants of the student, he introduced a large number of additional wood-engravings, illustrative of the objects described, while the publishers have endeavored to render the mechanical execution of the work worthy of its extended reputation.

JONES (T. WHARTON), F. R. S.,

Professor of Ophthalmic Medicine and Surgery in University College, London, &c.

THE PRINCIPLES AND PRACTICE OF OPHTHALMIC MEDICINE AND SURGERY. With one hundred and ten illustrations. Second American from the second and revised London edition, with additions by EDWARD HARTSHORNE, M. D., Surgeon to Wills' Hospital, &c. In one large, handsome royal 12mo. volume, extra cloth, of 500 pages. (*Now Ready*.) \$1 50.

We are confident that the reader will find, on perusal, that the execution of the work amply fulfils the promise of the preface, and sustains, in every point, the already high reputation of the author as an ophthalmic surgeon as well as a physiologist and pathologist. The book is evidently the result of much labor and research, and has been written with the greatest care and attention; it possesses that best quality which a general work, like a system or manual can show, viz: the quality of having all the materials whencesoever derived, so thorough-

ly wrought up, and digested in the author's mind, as to come forth with the freshness and impressiveness of an original production. We entertain little doubt that this book will become what its author hoped it might become, a manual for daily reference and consultation by the student and the general practitioner. The work is marked by that correctness, clearness, and precision of style which distinguish all the productions of the learned author.—*British and For. Med. Review*.

JONES (C. HANDFIELD), F. R. S., & EDWARD H. SIEVEKING, M. D.,
Assistant Physicians and Lecturers in St. Mary's Hospital, London.

A MANUAL OF PATHOLOGICAL ANATOMY. First American Edition, Revised. With three hundred and ninety-seven handsome wood engravings. In one large and beautiful octavo volume of nearly 750 pages, leather. (*Lately Issued.*) \$3 75.

As a concise text-book, containing, in a condensed form, a complete outline of what is known in the domain of Pathological Anatomy, it is perhaps the best work in the English language. Its great merit consists in its completeness and brevity, and in this respect it supplies a great desideratum in our literature. Heretofore the student of pathology was obliged to glean from a great number of monographs, and the field was so extensive that but few cultivated it with any degree of success. As a simple work of reference, therefore, it is of great value to the student of pathological anatomy, and should be in every physician's library.—*Western Lancet*.

We urge upon our readers and the profession generally the importance of informing themselves in regard to modern views of pathology, and recommend to them to procure the work before us as the best means of obtaining this information.—*Stethoscope*.

In offering the above titled work to the public, the

authors have not attempted to intrude new views on their professional brethren, but simply to lay before them, what has long been wanted, an outline of the present condition of pathological anatomy. In this they have been completely successful. The work is one of the best compilations which we have ever perused.—*Charleston Medical Journal and Review*.

We have no hesitation in recommending it as worthy of careful and thorough study by every member of the profession, old or young.—*N. W. Med. and Surg. Journal*.

From the casual examination we have given we are inclined to regard it as a text-book, plain, rational, and intelligible, such a book as the practical man needs for ready reference. For this reason it will be likely to be largely useful, as it suits itself to those busy men who have little time for minute investigation, and prefer a summary to an elaborate treatise.—*Buffalo Medical Journal*.

KIRKES (WILLIAM SENHOUSE), M. D.,
Demonstrator of Morbid Anatomy at St. Bartholomew's Hospital, &c.

A MANUAL OF PHYSIOLOGY. Second American, from the second and improved London edition. With one hundred and sixty-five illustrations. In one large and handsome royal 12mo. volume, leather. pp. 550. \$2 00.

In the present edition, the Manual of Physiology has been brought up to the actual condition of the science, and fully sustains the reputation which it has already so deservedly attained. We consider the work of MM. Kirkes and Paget to constitute one of the very best handbooks of Physiology we possess—presenting just such an outline of the science, comprising an account of its leading facts and generally admitted principles, as the student requires during his attendance upon a course of lectures, or for reference whilst preparing for examination.—*Am. Medical Journal*.

We need only say, that, without entering into discussions of unsettled questions, it contains all the recent improvements in this department of medical science. For the student beginning this study, and

the practitioner who has but leisure to refresh his memory, this book is invaluable, as it contains all that it is important to know, without special details, which are read with interest only by those who would make a specialty, or desire to possess a critical knowledge of the subject.—*Charleston Medical Journal*.

One of the best treatises that can be put into the hands of the student.—*London Medical Gazette*.

Particularly adapted to those who desire to possess a concise digest of the facts of Human Physiology.—*British and Foreign Med.-Chirurg. Review*.

We conscientiously recommend it as an admirable "Handbook of Physiology."—*London Journal of Medicine*.

KNAPP'S TECHNOLOGY; or, Chemistry applied to the Arts and to Manufactures. Edited, with numerous Notes and Additions, by Dr. EDMUND RONALDS and Dr. THOMAS RICHARDSON. First American edition, with Notes and Additions, by Prof. WALTER R. JOHNSON. In two handsome

octavo volumes, extra cloth, with about 500 wood engravings. \$6 00.

LAILLEMAND ON SPERMATORRHOEA. Translated and edited by HENRY J. McDOUGAL. In one volume, octavo, extra cloth, 320 pages. Second American edition. \$1 75.

LUDLOW (J. L.), M. D.

A MANUAL OF EXAMINATIONS upon Anatomy, Physiology, Surgery, Practice of Medicine, Obstetrics, Materia Medica, Pharmacy, and Therapeutics. To which is added a Medical Formulary. Designed for Students of Medicine throughout the United States. Third edition, thoroughly revised and greatly extended and enlarged. With over three hundred and fifty illustrations. In one large and handsome royal 12mo. volume, leather, of over 800 closely printed pages (*Just Ready.*) \$2 50.

The great popularity of this volume, and the numerous demands for it during the two years in which it has been out of print, have induced the author in its revision to spare no pains to render it a correct and accurate digest of the most recent condition of all the branches of medical science. In many respects it may, therefore, be regarded rather as a new book than a new edition, an entire section on Physiology having been added, as also one on Organic Chemistry, and many portions having been rewritten. A very complete series of illustrations has been introduced, and every care has been taken in the mechanical execution to render it a convenient and satisfactory book for study or reference.

The arrangement of the volume in the form of question and answer renders it especially suited for the office examination of students and for those preparing for graduation.

LARDNER (DIONYSIUS), D. C. L., &c.

HANDBOOKS OF NATURAL PHILOSOPHY AND ASTRONOMY.

Revised, with numerous Additions, by the American editor. **FIRST COURSE**, containing Mechanics, Hydrostatics, Hydraulics, Pneumatics, Sound, and Optics. In one large royal 12mo. volume, of 750 pages, with 424 wood-cuts. \$1 75. **SECOND COURSE**, containing Heat, Electricity, Magnetism, and Galvanism, one volume, large royal 12mo., of 450 pages, with 250 illustrations. \$1 25. **THIRD COURSE** (*now ready*), containing Meteorology and Astronomy, in one large volume, royal 12mo. of nearly eight hundred pages, with thirty-seven plates and two hundred wood-cuts. \$2 00. The whole complete in three volumes, of about two thousand large pages, with over one thousand figures on steel and wood. \$5 00. Any volume sold separate, strongly bound in leather.

LEHMANN (C. G.)

PHYSIOLOGICAL CHEMISTRY. Translated from the second edition by GEORGE E. DAY, M. D., F. R. S., &c., edited by R. E. ROGERS, M. D., Professor of Chemistry in the Medical Department of the University of Pennsylvania, with illustrations selected from Funke's Atlas of Physiological Chemistry, and an Appendix of plates. Complete in two large and handsome octavo volumes, extra cloth, containing 1200 pages, with nearly two hundred illustrations. (*Now Ready.*) \$6 00.

This great work, universally acknowledged as the most complete and authoritative exposition of the principles and details of Zoochemistry, in its passage through the press, has received from Professor Rogers such care as was necessary to present it in a correct and reliable form. To such a work additions were deemed superfluous, but several years having elapsed between the appearance in Germany of the first and last volume, the latter contained a supplement, embodying numerous corrections and additions resulting from the advance of the science. These have all been incorporated in the text in their appropriate places, while the subjects have been still further elucidated by the insertion of illustrations from the Atlas of Dr. Otto Funke. With the view of supplying the student with the means of convenient comparison, a large number of wood-cuts, from works on kindred subjects, have also been added in the form of an Appendix of Plates. The work is, therefore, presented as in every way worthy the attention of all who desire to be familiar with the modern facts and doctrines of Physiological Science.

The most important contribution as yet made to Physiological Chemistry.—*Am. Journal Med. Sciences*, Jan. 1856.

The present volumes belong to the small class of medical literature which comprises elaborate works of the highest order of merit.—*Montreal Med. Chronicle*, Jan. 1856.

The work of Lehmann stands unrivalled as the most comprehensive book of reference and information extant on every branch of the subject on which it treats.—*Edinburgh Monthly Journal of Medical Science*.

All teachers must possess it, and every intelligent physician ought to do likewise.—*Southern Med. and Surg. Journal*, Dec. 1855.

Already well known and appreciated by the scientific world, Professor Lehmann's great work requires no laudatory sentences, as, under a new garb, it is now presented to us. The little space at our command would ill suffice to set forth even a small portion of its excellences. To all whose studies or professional duties render the revelations of Physiological Chemistry at once interesting and essential, these volumes will be indispensable. Highly complimented by European reviewers, sought for with avidity by scholars of every nation, and admirably written throughout, it is sure to win a welcome and to be thoroughly studied.—*Boston Med. and Surg. Journal*, Dec. 1855.

BY THE SAME AUTHOR. (*Now Ready, 1856.*)

MANUAL OF CHEMICAL PHYSIOLOGY. Translated from the German, with Notes and Additions, by J. CHESTON MORRIS, M. D., with an Introductory Essay on Vital Force, by SAMUEL JACKSON, M. D., Professor of the Institutes of Medicine in the University of Pennsylvania. With illustrations on wood. In one very handsome octavo volume, extra cloth, of 363 pages. \$2 25.

The original of this work, though but lately issued by its distinguished author, has already assumed the highest position, as presenting in their latest development the modern doctrines and discoveries in the chemistry of life. The numerous additions by the translator, and the Introduction by Professor JACKSON will render its physiological aspect more complete than designed by the author, and will adapt it for use as a text-book of physiology, presenting more thoroughly than has yet been attempted, the modifications arising from the vast impulse which organic chemistry has received within a few years past.

From Prof. Jackson's Introductory Essay.

In adopting the handbook of Dr. Lehmann as a manual of Organic Chemistry for the use of the students of the University, and in recommending his original work of PHYSIOLOGICAL CHEMISTRY for their more mature studies, the high value of his researches, and the great weight of his authority in that important department of medical science are fully recognized.

The present volume will be a very convenient one for students, as offering a brief epitome of the more elaborate work, and as containing, in a very condensed form, the positive facts of Physiological Chemistry.—*Am. Journal Med. Sciences*, April, 1856.

LAWRENCE (W.), F. R. S., &c.

A TREATISE ON DISEASES OF THE EYE. A new edition, edited, with numerous additions, and 243 illustrations, by ISAAC HAYS, M. D., Surgeon to Will's Hospital, &c. In one very large and handsome octavo volume, of 950 pages, strongly bound in leather with raised bands. \$5 00.

This work is so universally recognized as the standard authority on the subject, that the publishers in presenting this new edition have only to remark that in its preparation the editor has carefully revised every portion, introducing additions and illustrations wherever the advance of science has rendered them necessary or desirable, constituting it a complete and thorough exponent of the most advanced state of the subject.

This admirable treatise—the safest guide and most comprehensive work of reference, which is within the reach of the profession.—*Stethoscope*.

This standard text-book on the department of which it treats, has not been superseded, by any or all of the numerous publications on the subject heretofore issued. Nor with the multiplied improvements of Dr. Hays, the American editor, is it at all likely that this great work will cease to merit the confidence and preference of students or practitioners. Its ample extent—nearly one thousand large

octavo pages—has enabled both author and editor to do justice to all the details of this subject, and condense in this single volume the present state of our knowledge of the whole science in this department, whereby its practical value cannot be excelled. We heartily commend it, especially as a book of reference, indispensable in every medical library. The additions of the American editor very greatly enhance the value of the work, exhibiting the learning and experience of Dr. Hays, in the light in which he ought to be held, as a standard authority on all subjects appertaining to this specialty.—*N. Y. Med. Gaz.*

LA ROCHE (R.), M. D., &c.

YELLOW FEVER, considered in its Historical, Pathological, Etiological, and Therapeutical Relations. Including a Sketch of the Disease as it has occurred in Philadelphia from 1699 to 1854, with an examination of the connections between it and the fevers known under the same name in other parts of temperate as well as in tropical regions. In two large and handsome octavo volumes of nearly 1500 pages, extra cloth. (*Just Ready.*) \$7 00.

From Professor S. H. Dickson, Charleston, S. C.,
September 18, 1855.

A monument of intelligent and well applied research, almost without example. It is, indeed, in itself, a large library, and is destined to constitute the special resort as a book of reference, in the subject of which it treats, to all future time.

We have not time at present, engaged as we are, by day and by night, in the work of combating this very disease, now prevailing in our city, to do more than give this cursory notice of what we consider as undoubtedly the most able and erudite medical publication our country has yet produced. But in view of the startling fact, that this, the most malignant and unmanageable disease of modern times, has for several years been prevailing in our country to a greater extent than ever before; that it is no longer confined to either large or small cities, but penetrates country villages, plantations, and farm-houses; that it is treated with scarcely better success now than thirty or forty years ago; that there is vast mischief done by ignorant pretenders to knowledge in regard to the disease, and in view of the probability that a majority of southern physicians will be called upon to treat the disease, we trust that this able and comprehensive treatise will be very generally read in the south.—*Memphis Med. Recorder.*

This is decidedly the great American medical work of the day—a full, complete, and systematic treatise, unequalled by any other upon the all-important subject of Yellow Fever. The laborious, indefatigable, and learned author has devoted to it many years of arduous research and careful study, and the result is such as will reflect the highest honor upon the author and our country.—*Southern Med. and Surg. Journal.*

The genius and scholarship of this great physician could not have been better employed than in the

erection of this towering monument to his own fame, and to the glory of the medical literature of his own country. It is destined to remain the great authority upon the subject of Yellow Fever. The student and physician will find in these volumes a *résumé* of the sum total of the knowledge of the world upon the awful scourge which they so elaborately discuss. The style is so soft and so pure as to refresh and invigorate the mind while absorbing the thoughts of the gifted author, while the publishers have succeeded in bringing the externals into a most felicitous harmony with the inspiration that dwells within. Take it all in all, it is a book we have often dreamed of, but dreamed not that it would ever meet our waking eye as a tangible reality.—*Nashville Journal of Medicine.*

We deem it fortunate that the splendid work of Dr. La Roche should have been issued from the press at this particular time. The want of a reliable digest of all that is known in relation to this frightful malady has long been felt—a want very satisfactorily met in the work before us. We deem it but faint praise to say that Dr. La Roche has succeeded in presenting the profession with an able and complete monograph, one which will find its way into every well ordered library.—*Va. Stethoscope.*

Although we have no doubt that controversial treatises on the mode of origin and propagation of the fever in question will, as heretofore, occasionally appear, yet it must be some time before another systematic work can arise in the face of so admirable and carefully executed a one as the present. It is a mine of information, quite an encyclopædia of references, and *résumé* of knowledge relative to what has been recorded upon the subject.—*London Lancet.*

A miracle of industry and research, constituting a complete library of reference on the disease of which it treats.—*Dublin Quarterly Journal.*

BY THE SAME AUTHOR.

PNEUMONIA; its Supposed Connection, Pathological and Etiological, with Autumnal Fevers, including an Inquiry into the Existence and Morbid Agency of Malaria. In one handsome octavo volume, extra cloth, of 500 pages. \$3 00.

A more simple, clear, and forcible exposition of the groundless nature and dangerous tendency of certain pathological and etiological heresies, has seldom been presented to our notice.—*N. Y. Journal of Medicine and Collateral Science.*

This work should be carefully studied by Southern physicians, embodying as it does the reflections of an original thinker and close observer on a subject peculiarly their own.—*Virginia Med. and Surg. Journal.*

LAYCOCK (THOMAS), M. D., F. R. S. E.,

Professor of Practical and Clinical Medicine in the University of Edinburgh, &c.

LECTURES ON THE PRINCIPLES AND METHODS OF MEDICAL OBSERVATION AND RESEARCH. For the Use of Advanced Students and Junior Practitioners. In one very neat royal 12mo. volume, extra cloth. Price \$1 00. (*Now Ready.*)

Though addressed more particularly to the student, this little volume will be found of much use to all members of the profession who aim at rendering their practice a science rather than a routine. The importance of the development and application of principles, and the exercise of correct and logical reasoning from clinical observations are beginning to be generally recognized, and in furtherance of this object, a wide circulation of the present volume cannot fail to exercise a most beneficial influence. The object and scope of the work may be gathered from the following condensed summary of the contents:—

LECTURE I. General Principles of Observation and Inquiry—Nature and Acquisition of Experience in Medicine—Combination of Experience with Theory and Observation, with illustrations of the Fallacious use of Theories. LECT. II. General Methods and Objects of Clinical Study. LECT. III. Methods of Clinical Examination—Clinical Observations of General or Constitutional Morbid States. LECT. IV. On Prognosis, and the Order of Succession of Morbid Phenomena. LECT. V. On the Due Estimate of Treatment—Management of the Case. LECT. VI. Numerical Method of Research in Medicine. LECT. VII. Analogical, Philosophical, or Purely Inductive Method of Research—Practical Examples of the Conduct of an Analogical Investigation—Examples of its Applications to Anatomy, Physiology, and Histology.

MULLER (PROFESSOR J.), M. D.

PRINCIPLES OF PHYSICS AND METEOROLOGY. Edited, with Additions, by R. EGLESFELD GRIFFITH, M. D. In one large and handsome octavo volume, extra cloth, with 550 wood-cuts, and two colored plates. pp. 636. \$3 50.

MEIGS (CHARLES D.), M. D.,

Professor of Obstetrics, &c. in the Jefferson Medical College, Philadelphia.

OBSTETRICS: THE SCIENCE AND THE ART. Third edition, revised and improved. With one hundred and twenty-nine illustrations. In one beautifully printed octavo volume, leather, of seven hundred and fifty-two large pages. \$3 75. (*Now Ready.*)

The rapid demand for another edition of this work is a sufficient expression of the favorable verdict of the profession. In thus preparing it a third time for the press, the author has endeavored to render it in every respect worthy of the favor which it has received. To accomplish this he has thoroughly revised it in every part. Some portions have been rewritten, others added, new illustrations have been in many instances substituted for such as were not deemed satisfactory, while, by an alteration in the typographical arrangement, the size of the work has not been increased, and the price remains unaltered. In its present improved form, it is, therefore, hoped that the work will continue to meet the wants of the American profession as a sound, practical, and extended **SYSTEM OF MIDWIFERY.**

BY THE SAME AUTHOR. (*Lately Issued.*)

WOMAN: HER DISEASES AND THEIR REMEDIES. A Series of Lectures to his Class. Third and Improved edition. In one large and beautifully printed octavo volume, leather. pp. 672. \$3 60.

The gratifying appreciation of his labors, as evinced by the exhaustion of two large impressions of this work within a few years, has not been lost upon the author, who has endeavored in every way to render it worthy of the favor with which it has been received. The opportunity thus afforded for a second revision has been improved, and the work is now presented as in every way superior to its predecessors, additions and alterations having been made whenever the advance of science has rendered them desirable. The typographical execution of the work will also be found to have undergone a similar improvement, and the work is now confidently presented as in every way worthy the position it has acquired as the standard American text-book on the Diseases of Females.

It contains a vast amount of practical knowledge, by one who has accurately observed and retained the experience of many years, and who tells the result in a free, familiar, and pleasant manner.—*Dublin Quarterly Journal.*

There is an off-hand fervor, a glow, and a warm-heartedness infecting the effort of Dr. Meigs, which is entirely captivating, and which absolutely hurries the reader through from beginning to end. Besides, the book teems with solid instruction, and it shows the very highest evidence of ability, viz., the clearness with which the information is presented. We know of no better test of one's understanding a subject than the evidence of the power of lucidly explaining it. The most elementary, as well as the obscurest subjects, under the pencil of Prof. Meigs, are isolated and made to stand out in

such bold relief, as to produce distinct impressions upon the mind and memory of the reader.—*The Charleston Med. Journal.*

Professor Meigs has enlarged and amended this great work, for such it unquestionably is, having passed the ordeal of criticism at home and abroad, but been improved thereby; for in this new edition the author has introduced real improvements, and increased the value and utility of the book immeasurably. It presents so many novel, bright, and sparkling thoughts; such an exuberance of new ideas on almost every page, that we confess ourselves to have become enamored with the book and its author; and cannot withhold our congratulations from our Philadelphia conferees, that such a teacher is in their service.—*N. Y. Med. Gazette.*

BY THE SAME AUTHOR. (*Lately Published.*)

ON THE NATURE, SIGNS, AND TREATMENT OF CHILD BED FEVER. In a Series of Letters addressed to the Students of his Class. In one handsome octavo volume, extra cloth, of 365 pages. \$2 50.

The instructive and interesting author of this work, whose previous labors in the department of medicine which he so sedulously cultivates, have placed his countrymen under deep and abiding obligations, again challenges their admiration in the fresh and vigorous, attractive and racy pages before us. It is a delectable book. * * * This treatise upon child-bed fevers will have an extensive sale, being destined, as it deserves, to find a place in the library of every practitioner who scorns to lag in the rear of his brethren.—*Nashville Journal of Medicine and Surgery.*

This book will add more to his fame than either of those which bear his name. Indeed we doubt whether any material improvement will be made on the teachings of this volume for a century to come, since it is so eminently practical, and based on profound knowledge of the science and consummate skill in the art of healing, and ratified by an ample and extensive experience, such as few men have the industry or good fortune to acquire.—*N. Y. Med. Gazette.*

BY THE SAME AUTHOR; WITH COLORED PLATES.

A TREATISE ON ACUTE AND CHRONIC DISEASES OF THE NECK OF THE UTERUS. With numerous plates, drawn and colored from nature in the highest style of art. In one handsome octavo volume, extra cloth. \$4 50.

BY THE SAME AUTHOR.

OBSERVATIONS ON CERTAIN OF THE DISEASES OF YOUNG CHILDREN. In one handsome octavo volume, extra cloth, of 214 pages. \$1 75

MAYNE'S DISPENSATORY AND THERAPEUTICAL REMEMBRANCER. Comprising the entire lists of Materia Medica, with every Practical Formula contained in the three British Pharmacopœias. Edited, with the addition of the Formulæ of the U. S. Pharmacopœia, by R. E. GRIFFITH, M. D. 12mo. vol. ex. cl., 300 pp. 75 c.

MALGAIGNE'S OPERATIVE SURGERY, based on Normal and Pathological Anatomy. Translated from the French by FREDERICK BRITTON, A. B., M. D. With numerous illustrations on wood. In one handsome octavo volume, extra cloth, of nearly six hundred pages. \$2 25.

MACLISE (JOSEPH), SURGEON.

SURGICAL ANATOMY. Forming one volume, very large imperial quarto.

With sixty-eight large and splendid Plates, drawn in the best style and beautifully colored. Containing one hundred and ninety Figures, many of them the size of life. Together with copious and explanatory letter-press. Strongly and handsomely bound in extra cloth, being one of the cheapest and best executed Surgical works as yet issued in this country. \$11 00.

* * The size of this work prevents its transmission through the post-office as a whole, but those who desire to have copies forwarded by mail, can receive them in five parts, done up in stout wrappers. Price \$9 00.

One of the greatest artistic triumphs of the age in Surgical Anatomy.—*British American Medical Journal*.

Too much cannot be said in its praise: indeed, we have not language to do it justice.—*Ohio Medical and Surgical Journal*.

The most admirable surgical atlas we have seen. To the practitioner deprived of demonstrative dissections upon the human subject, it is an invaluable companion.—*N. J. Medical Reporter*.

The most accurately engraved and beautifully colored plates we have ever seen in an American book—one of the best and cheapest surgical works ever published.—*Buffalo Medical Journal*.

It is very rare that so elegantly printed, so well illustrated, and so useful a work, is offered at so moderate a price.—*Charleston Medical Journal*.

Its plates can boast a superiority which places them almost beyond the reach of competition.—*Medical Examiner*.

Every practitioner, we think, should have a work of this kind within reach.—*Southern Medical and Surgical Journal*.

No such lithographic illustrations of surgical regions have hitherto, we think, been given.—*Boston Medical and Surgical Journal*.

As a surgical anatomist, Mr. MacLise has probably no superior.—*British and Foreign Medico-Chirurgical Review*.

Of great value to the student engaged in dissecting, and to the surgeon at a distance from the means

✎ The very low price at which this work is furnished, and the beauty of its execution, require an extended sale to compensate the publishers for the heavy expenses incurred.

MOHR (FRANCIS) PH. D., AND REDWOOD (THEOPHILUS).

PRACTICAL PHARMACY. Comprising the Arrangements, Apparatus, and Manipulations of the Pharmaceutical Shop and Laboratory. Edited, with extensive Additions, by Prof. WILLIAM PROCTER, of the Philadelphia College of Pharmacy. In one handsomely printed octavo volume, extra cloth, of 570 pages, with over 500 engravings on wood. \$2 75.

MACKENZIE (W.), M. D.,

Surgeon Oculist in Scotland in ordinary to Her Majesty, &c. &c.

A PRACTICAL TREATISE ON DISEASES AND INJURIES OF THE

EYE. To which is prefixed an Anatomical Introduction explanatory of a Horizontal Section of the Human Eyeball, by THOMAS WHARTON JONES, F. R. S. From the Fourth Revised and Enlarged London Edition. With Notes and Additions by ADDINELL HEWSON, M. D., Surgeon to Wills Hospital, &c. &c. In one very large and handsome octavo volume, leather, raised bands, with plates and numerous wood-cuts. (Now Ready.) \$5 25.

The treatise of Dr. Mackenzie indisputably holds the first place, and forms, in respect of learning and research, an Encyclopædia unequalled in extent by any other work of the kind, either English or foreign.—*Dixon on Diseases of the Eye*.

Few modern books on any department of medicine or surgery have met with such extended circulation, or have procured for their authors a like amount of European celebrity. The immense research which it displayed, the thorough acquaintance with the subject, practically as well as theoretically, and the able manner in which the author's stores of learning and experience were rendered available for general use, at once procured for the first edition, as well on the continent as in this country, that high position as a standard work which each successive edition has more firmly established, in spite of the attractions of several rivals of no mean ability. This, the fourth edition, has been in a great measure re-written; new matter, to the extent of one hundred and fifty pages, has been added, and in several instances formerly expressed opinions have been modified in

accordance with the advances in the science which have been made of late years. Nothing worthy of repetition upon any branch of the subject appears to have escaped the author's notice. We consider it the duty of every one who has the love of his profession and the welfare of his patient at heart, to make himself familiar with this the most complete work in the English language upon the diseases of the eye.—*Med. Times and Gazette*.

The mechanical execution cannot be excelled.—*Transylvania Medical Journal*.

A work which has no parallel in point of accuracy and cheapness in the English language.—*N. Y. Journal of Medicine*.

To all engaged in the study or practice of their profession, such a work is almost indispensable.—*Dublin Quarterly Medical Journal*.

No practitioner whose means will admit should fail to possess it.—*Ranking's Abstract*.

Country practitioners will find these plates of immense value.—*N. Y. Medical Gazette*.

We are extremely gratified to announce to the profession the completion of this truly magnificent work, which, as a whole, certainly stands unrivalled, both for accuracy of drawing, beauty of coloring, and all the requisite explanations of the subject in hand.—*The New Orleans Medical and Surgical Journal*.

This is by far the ablest work on Surgical Anatomy that has come under our observation. We know of no other work that would justify a student, in any degree, for neglect of actual dissection. In those sudden emergencies that so often arise, and which require the instantaneous command of minute anatomical knowledge, a work of this kind keeps the details of the dissecting-room perpetually fresh in the memory.—*The Western Journal of Medicine and Surgery*.

✎ The very low price at which this work is furnished, and the beauty of its execution, require an extended sale to compensate the publishers for the heavy expenses incurred.

The fourth edition of this standard work will no doubt be as fully appreciated as the three former editions. It is unnecessary to say a word in its praise, for the verdict has already been passed upon it by the most competent judges, and "Mackenzie on the Eye" has justly obtained a reputation which it is no figure of speech to call world-wide.—*British and Foreign Medico-Chirurgical Review*.

This new edition of Dr. Mackenzie's celebrated treatise on diseases of the eye, is truly a miracle of industry and learning. We need scarcely say that he has entirely exhausted the subject of his speciality.—*Dublin Quarterly Journal*.

MILLER (JAMES), F. R. S. E.,
Professor of Surgery in the University of Edinburgh, &c.

PRINCIPLES OF SURGERY. Fourth American, from the third and revised Edinburgh edition. In one large and very beautiful volume, leather, of 700 pages, with two hundred and forty exquisite illustrations on wood. (*Now Ready*, 1856.) \$3 75.

The extended reputation enjoyed by this work will be fully maintained by the present edition. Thoroughly revised by the author, it will be found a clear and compendious exposition of surgical science in its most advanced condition.

In connection with the recently issued third edition of the author's "Practice of Surgery," it forms a very complete system of Surgery in all its branches.

The work of Mr. Miller is too well and too favorably known among us, as one of our best text-books, to render any further notice of it necessary than the announcement of a new edition, the *fourth* in our country, a proof of its extensive circulation among us. As a concise and reliable exposition of the science of modern surgery, it stands deservedly high—we know not its superior.—*Boston Med. and Surg. Journal*.

The works of Professor Miller are so well known to the profession, that it is unnecessary for us say anything in relation to their general merits. The present edition of his "Principles," however, deserves a special notice, from the number, variety, and faithfulness of its illustrations. The wood-cuts are beautifully executed, and many of them are new and exceedingly instructive, particularly those illustrating mortification, diseased and fractured bones, and the varieties of aneurism.—*Western Lancet*.

This edition is far superior, both in the abundance and quality of its material, to any of the preceding. We hope it will be extensively read, and the sound principles which are herein taught treasured up for future application. The work takes rank with Watson's Practice of Physic; it certainly does not

fall behind that great work in soundness of principle or depth of reasoning and research. No physician who values his reputation, or seeks the interests of his clients, can acquit himself before his God and the world without making himself familiar with the sound and philosophical views developed in the foregoing book.—*New Orleans Med. and Surg. Journal*.

Without doubt the ablest exposition of the principles of that branch of the healing art in any language. This opinion, deliberately formed after a careful study of the first edition, we have had no cause to change on examining the second. This edition has undergone thorough revision by the author; many expressions have been modified, and a mass of new matter introduced. The book is got up in the finest style, and is an evidence of the progress of typography in our country.—*Charleston Medical Journal and Review*.

We recommend it to both student and practitioner, feeling assured that as it now comes to us, it presents the most satisfactory exposition of the modern doctrines of the principles of surgery to be found in any volume in any language.—*N. Y. Journal of Medicine*.

BY THE SAME AUTHOR. (*Lately Published*.)

THE PRACTICE OF SURGERY. Third American from the second Edinburgh edition. Edited, with Additions, by F. W. SARGENT, M. D., one of the Surgeons to Will's Hospital, &c. Illustrated by three hundred and nineteen engravings on wood. In one large octavo volume, leather, of over 700 pages. \$3 75.

No eulogium of ours could add to the popularity of Miller's Surgery. Its reputation in this country is unsurpassed by that of any other work, and, when taken in connection with the author's *Principles of Surgery*, constitutes a whole, without reference to which no conscientious surgeon would be willing to practice his art. The additions, by Dr. Sargent, have materially enhanced the value of the work.—*Southern Medical and Surgical Journal*.

It is seldom that two volumes have ever made so profound an impression in so short a time as the "Principles" and the "Practice" of Surgery by Mr. Miller—or so richly merited the reputation they have acquired. The author is an eminently sensible, practical, and well-informed man, who knows exactly what he is talking about and exactly how to talk it.—*Kentucky Medical Recorder*.

By the almost unanimous voice of the profession,

his works, both on the principles and practice of surgery have been assigned the highest rank. If we were limited to but one work on surgery, that one should be Miller's, as we regard it as superior to all others.—*St. Louis Med. and Surg. Journal*.

The author, distinguished alike as a practitioner and writer, has in this and his "Principles," presented to the profession one of the most complete and reliable systems of Surgery extant. His style of writing is original, impressive, and engaging, energetic, concise, and lucid. Few have the faculty of condensing so much in small space, and at the same time so persistently holding the attention; indeed, he appears to make the very process of condensation a means of eliminating attractions. Whether as a text-book for students or a book of reference for practitioners, it cannot be too strongly recommended.—*Southern Journal of Med. and Phys. Sciences*.

MONTGOMERY (W. F.), M. D., M. R. I. A., &c.,
Professor of Midwifery in the King and Queen's College of Physicians in Ireland, &c.

AN EXPOSITION OF THE SIGNS AND SYMPTOMS OF PREGNANCY.

With some other Papers on Subjects connected with Midwifery. From the second and enlarged English edition. With two exquisite colored plates, and numerous wood-cuts. In one very handsome octavo volume, extra cloth, of nearly 600 pages. (*Just Ready*.) \$3 75.

The present edition of this classical volume is fairly entitled to be regarded as a new work, every sentence having been carefully rewritten, and the whole increased to more than double the original size. Translated into various languages, and universally regarded as the highest authority on the numerous interesting questions discussed, it will be found in its present form to fully maintain its high character, the author having exhausted the subject in all its bearings, examining every point with all the aid furnished by the most recent investigations, assisted by his own long and enlarged experience. The title of the work scarcely does justice to the extent and importance of the topics brought under consideration, embracing, with the exception of the operative procedures of midwifery, almost everything connected with obstetrics, either directly or incidentally; and there are few physicians who will not find in its pages much that will prove of great interest and value in their daily practice. The special Essays on the Period of Human Gestation, the Signs of Delivery, and the Spontaneous Amputation and other Lesions of the Fetus in Utero present topics of the highest interest fully treated and beautifully illustrated.

In every point of mechanical execution the work will be found one of the handsomest yet issued from the American press.

NEILL (JOHN), M. D.,

Surgeon to the Pennsylvania Hospital, &c.; and

FRANCIS GURNEY SMITH, M. D.,

Professor of Institutes of Medicine in the Pennsylvania Medical College.

AN ANALYTICAL COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE; for the Use and Examination of Students. A new edition, revised and improved. In one very large and handsomely printed royal 12mo. volume, of about one thousand pages, with three hundred and seventy-four illustrations on wood. Strongly bound in leather, with raised bands. (*Now Ready, 1856.*) \$3 00.

The very flattering reception which has been accorded to this work, and the high estimate placed upon it by the profession, as evinced by the constant and increasing demand which has rapidly exhausted two large editions, have stimulated the authors to render the volume in its present revision more worthy of the success which has attended it. It has accordingly been thoroughly examined, and such errors as had on former occasions escaped observation have been corrected, and whatever additions were necessary to maintain it on a level with the advance of science have been introduced. The extended series of illustrations has been still further increased and much improved, while, by a slight enlargement of the page, these various additions have been incorporated without increasing the bulk of the volume.

The work is, therefore, again presented as eminently worthy of the favor with which it has hitherto been received. As a book for daily reference by the student requiring a guide to his more elaborate text-books, as a manual for preceptors desiring to stimulate their students by frequent and accurate examination, or as a source from which the practitioners of older date may easily and cheaply acquire a knowledge of the changes and improvement in professional science, its reputation is permanently established.

In the rapid course of lectures, where work for the students is heavy, and review necessary for an examination, a compend is not only valuable, but it is almost a *sine qua non*. The one before us, in most of the divisions, the most unexceptionable of all books of the kind that we know of. The newest and soundest doctrines and the latest improvements and discoveries are explicitly, though concisely, laid before the student. Of course it is useless for us to recommend it to all last course students, but there is a class to whom we very sincerely commend this cheap book as worth its weight in silver—that class is the graduates in medicine of more than ten years' standing, who have not studied medicine since. They will perhaps find out from it that the science is not exactly now

what it was when they left it off.—*The Stethoscope.*

We recommend it to our readers as the best work of the kind with which we are acquainted.—*Med. Examiner*, April, 1856.

Having made free use of this volume in our examinations of pupils, we can speak from experience in recommending it as an admirable compend for students, and as especially useful to preceptors who examine their pupils. It will save the teacher much labor by enabling him readily to recall all of the points upon which his pupils should be examined. A work of this sort should be in the hands of every one who takes pupils into his office with a view of examining them; and this is unquestionably the best of its class.—*Transylvania Med. Journal.*

NEILL (JOHN), M. D.,

Professor of Surgery in the Pennsylvania Medical College, &c.

OUTLINES OF THE VEINS AND LYMPHATICS. With handsome colored plates. 1 vol., cloth. \$1 25.

OUTLINES OF THE NERVES. With handsome plates. 1 vol., cloth. \$1 25.

NELIGAN (J. MOORE), M. D., M. R. I. A., &c.

(*A splendid work. Now Ready.*)

ATLAS OF CUTANEOUS DISEASES. In one beautiful quarto volume, extra cloth, with splendid colored plates, presenting nearly one hundred elaborate representations of disease. \$4 50.

This beautiful volume is intended as a complete and accurate representation of all the varieties of Diseases of the Skin. While it can be consulted in conjunction with any work on Practice, it has especial reference to the author's "Treatise on Diseases of the Skin," so favorably received by the profession some years since. The publishers feel justified in saying that no more beautifully executed plates have ever been presented to the profession of this country.


The diagnosis of eruptive disease, however, under all circumstances, is very difficult. Nevertheless Dr. Neligan has certainly, "as far as possible," given a faithful and accurate representation of this class of diseases, and there can be no doubt that these plates will be of great use to the student and practitioner in drawing a diagnosis as to the class, order, and species to which the particular case may belong. While looking over the "Atlas" we have been induced to examine also the "Practical Treatise," and we are inclined to consider it a very superior work, combining accurate verbal description, with sound views of the pathology and treatment of eruptive diseases.—*Glasgow Med. Journal.*

The profession owes its thanks to the publishers of Neligan's Atlas of Cutaneous Diseases, for they have

placed within its reach and at a moderate cost a most accurate and well delineated series of plates illustrating the eruptive disorders. These plates are all drawn from the life, and in many of them the daguerreotype has been employed with great success. Such works as these are especially useful to country practitioners, who have not an opportunity of seeing the rarer forms of cutaneous disease, and hence need the aid of illustrations to give them the requisite information on the subject. With these plates at hand, the inexperienced practitioner is enabled to discriminate with much accuracy, and he is thus, comparatively speaking, put on an equal footing with those who have had the opportunity of visiting the large hospitals of Europe and America.—*Va. Med. Journal*, June, 1856.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. In one neat royal 12mo. volume, extra cloth, of 334 pages. \$1 00.

 The two volumes will be sent by mail on receipt of *Five Dollars*.

OWEN ON THE DIFFERENT FORMS OF THE SKELETON, AND OF THE TEETH.

One vol. royal 12mo., extra cloth, with numerous illustrations. (*Just Issued.*) \$1 25.

(Now Complete.)

PEREIRA (JONATHAN), M. D., F. R. S., AND L. S.

THE ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS.

Third American edition, enlarged and improved by the author; including Notices of most of the Medicinal Substances in use in the civilized world, and forming an Encyclopædia of Materia Medica. Edited, with Additions, by JOSEPH CARSON, M. D., Professor of Materia Medica and Pharmacy in the University of Pennsylvania. In two very large octavo volumes of 2100 pages, on small type, with about 500 illustrations on stone and wood, strongly bound in leather, with raised bands. \$9 00.

Gentlemen who have the first volume are recommended to complete their copies without delay. The first volume will no longer be sold separate. Price of Vol. II. \$5 00.

When we remember that Philology, Natural History, Botany, Chemistry, Physics, and the Microscope, are all brought forward to elucidate the subject, one cannot fail to see that the reader has here a work worthy of the name of an encyclopædia of Materia Medica. Our own opinion of its merits is that of its editors, and also that of the whole profession, both of this and foreign countries—namely, "that in copiousness of details, in extent, variety, and accuracy of information, and in lucid explanation of difficult and recondite subjects, it surpasses all other works on Materia Medica hitherto published." We cannot close this notice without alluding to the special additions of the American editor, which pertain to the prominent vegetable productions of this country, and to the directions of the United States Pharmacopœia, in connection with all the articles contained in the volume which are referred to by it. The illustrations have been increased, and this edition by Dr. Carson cannot well be regarded in any other light than that of a treasure which should be found in the library of every physician.—*New York Journal of Medical and Collateral Science.*

The third edition of his "Elements of Materia Medica, although completed under the supervision of others, is by far the most elaborate treatise in the English language, and will, while medical literature is cherished, continue a monument alike honorable to his genius, as to his learning and industry.—*American Journal of Pharmacy.*

The work, in its present shape, forms the most comprehensive and complete treatise on materia medica extant in the English language.—Dr. Pereira has been at great pains to introduce into his work, not only all the information on the natural, chemical, and commercial history of medicines, which might be serviceable to the physician and surgeon, but whatever might enable his readers to understand thoroughly the mode of preparing and manufacturing various articles employed either for preparing medicines, or for certain purposes in the arts connected with materia medica and the practice of medicine. The accounts of the physiological and therapeutic effects of remedies are given with great clearness and accuracy, and in a manner calculated to interest as well as instruct the reader.—*Edinburgh Medical and Surgical Journal.*

PEASELEE (E. R.), M. D.,

Professor of Anatomy and Physiology in Dartmouth College, &c.

HUMAN HISTOLOGY, in its applications to Physiology and General Pathology; designed as a Text-Book for Medical Students. With numerous illustrations. In one handsome royal 12mo. volume. (Preparing.)

The subject of this work is one, the growing importance of which, as the basis of Anatomy and Physiology, demands for it a separate volume. The book will therefore supply an acknowledged deficiency in medical text-books, while the name of the author, and his experience as a teacher for the last thirteen years, is a guarantee that it will be thoroughly adapted to the use of the student.

PIRRIE (WILLIAM), F. R. S. E.,

Professor of Surgery in the University of Aberdeen.

THE PRINCIPLES AND PRACTICE OF SURGERY. Edited by JOHN NEILL, M. D., Professor of Surgery in the Penna. Medical College, Surgeon to the Pennsylvania Hospital, &c. In one very handsome octavo volume, leather, of 780 pages, with 316 illustrations. \$3 75.

We know of no other surgical work of a reasonable size, wherein there is so much theory and practice, or where subjects are more soundly or clearly taught.—*The Stethoscope.*

There is scarcely a disease of the bones or soft parts, fracture, or dislocation, that is not illustrated by accurate wood-engravings. Then, again, every instrument employed by the surgeon is thus represented. These engravings are not only correct, but really beautiful, showing the astonishing degree of perfection to which the art of wood-engraving has

arrived. Prof. Pirrie, in the work before us, has elaborately discussed the principles of surgery, and a safe and effectual practice predicated upon them. Perhaps no work upon this subject heretofore issued is so full upon the science of the art of surgery.—*Nashville Journal of Medicine and Surgery.*

One of the best treatises on surgery in the English language.—*Canada Med. Journal.*

Our impression is, that, as a manual for students, Pirrie's is the best work extant.—*Western Med. and Surg. Journal.*

PANCOAST (J.), M. D.,

Professor of Anatomy in the Jefferson Medical College, Philadelphia, &c.

OPERATIVE SURGERY; or, A Description and Demonstration of the various Processes of the Art; including all the New Operations, and exhibiting the State of Surgical Science in its present advanced condition. Complete in one royal 4to. volume, extra cloth. of 380 pages of letter-press description and eighty large 4to. plates, comprising 486 illustrations. Second edition, improved. \$10 00.

This excellent work is constructed on the model of the French Surgical Works by Velpéau and Malgaigne; and, so far as the English language is con-

cerned, we are proud as an American to say that, OF ITS KIND IT HAS NO SUPERIOR.—*N. Y. Journal of Medicine.*

PARKER (LANGSTON),

Surgeon to the Queen's Hospital, Birmingham.

THE MODERN TREATMENT OF SYPHILITIC DISEASES, BOTH PRIMARY AND SECONDARY; comprising the Treatment of Constitutional and Confirmed Syphilis, by a safe and successful method. With numerous Cases, Formulæ, and Clinical Observations. From the Third and entirely rewritten London edition. In one neat octavo volume, extra cloth, of 316 pages. (Just Issued.) \$1 75.

PARRISH (EDWARD),

Lecturer on Practical Pharmacy and Materia Medica in the Pennsylvania Academy of Medicine, &c.

AN INTRODUCTION TO PRACTICAL PHARMACY. Designed as a Text-Book for the Student, and as a Guide for the Physician and Pharmacist. With many Formulæ and Prescriptions. In one handsome octavo volume, extra cloth, of 550 pages, with 243 Illustrations. (*Just Issued.*) \$2 75.

A careful examination of this work enables us to speak of it in the highest terms, as being the best treatise on practical pharmacy with which we are acquainted, and an invaluable *vade-mecum*, not only to the apothecary and to those practitioners who are accustomed to prepare their own medicines, but to every medical man and medical student. Throughout the work are interspersed valuable tables, useful formulæ, and practical hints, and the whole is illustrated by a large number of excellent wood-engravings.—*Boston Med. and Surg. Journal*.

This is altogether one of the most useful books we have seen. It is just what we have long felt to be needed by apothecaries, students, and practitioners of medicine, most of whom in this country have to put up their own prescriptions. It bears, upon every page, the impress of practical knowledge, conveyed in a plain common sense manner, and adapted to the comprehension of all who may read it. No detail has been omitted, however trivial it may seem, although really important to the dispenser of medicine.—*Southern Med. and Surg. Journal*.

To both the country practitioner and the city apothecary this work of Mr. Parrish is a godsend. A careful study of its contents will give the young graduate a familiarity with the value and mode of administering his prescriptions, which will be of as much use to his patient as to himself.—*Va. Med. Journal*.

Mr. Parrish has rendered a very acceptable service to the practitioner and student, by furnishing this book, which contains the leading facts and principles of the science of Pharmacy, conveniently arranged for study, and with special reference to those features of the subject which possess an especial practical interest to the physician. It furnishes the student, at the commencement of his studies, with that information which is of the greatest importance in initiating him into the domain of Chemistry and Materia

Medica; it familiarizes him with the compounding of drugs, and supplies those minutiae which but few practitioners can impart. The junior practitioner will, also, find this volume replete with instruction.—*Charleston Med. Journal and Review*, Mar. 1856.

There is no useful information in the details of the apothecary's or country physician's office conducted according to science that is omitted. The young physician will find it an encyclopedia of indispensable medical knowledge, from the purchase of a spatula to the compounding of the most learned prescriptions. The work is by the ablest pharmacist in the United States, and must meet with an immense sale.—*Nashville Journal of Medicine*, April, 1856.

We are glad to receive this excellent work. It will supply a want long felt by the profession, and especially by the student of Pharmacy. A large majority of physicians are obliged to compound their own medicines, and to them a work of this kind is indispensable.—*N. O. Medical and Surgical Journal*.

We cannot say but that this volume is one of the most welcome and appropriate which has for a long time been issued from the press. It is a work which we doubt not will at once secure an extensive circulation, as it is designed not only for the druggist and pharmacist, but also for the great body of practitioners throughout the country, who not only have to prescribe medicines, but in the majority of instances have to rely upon their own resources—whatever these may be—not only to compound, but also to manufacture the remedies they are called upon to administer. The author has not mistaken the idea in writing this volume, as it is alike useful and invaluable to those engaged in the active pursuits of the profession, and to those preparing to enter upon the field of professional labors.—*American Lancet*, March 24, 1856.

RICORD (P.), M. D.,

A TREATISE ON THE VENEREAL DISEASE. By JOHN HUNTER, F. R. S.

With copious Additions, by PH. RICORD, M. D. Edited, with Notes, by FREEMAN J. BUMSTEAD, M. D. In one handsome octavo volume, extra cloth, of 520 pages, with plates. \$3 25.

Every one will recognize the attractiveness and value which this work derives from this presenting the opinions of these two masters side by side. But it must be admitted, what has made the fortune of the book, is the fact that it contains the "most complete embodiment of the veritable doctrines of the Hôpital du Midi," which has ever been made public. The doctrinal ideas of M. Ricord, ideas which, if not universally adopted, are incontestably dominant, have heretofore only been interpreted by more or less skillful

secretaries, sometimes accredited and sometimes not.

In the notes to Hunter, the master substitutes himself for his interpreters, and gives his original thoughts to the world in a lucid and perfectly intelligible manner. In conclusion we can say that this is incontestably the best treatise on syphilis with which we are acquainted, and, as we do not often employ the phrase, we may be excused for expressing the hope that it may find a place in the library of every physician.—*Virginia Med. and Surg. Journal*.

BY THE SAME AUTHOR.

ILLUSTRATIONS OF SYPHILITIC DISEASE.

Translated by THOMAS F. BETTON, M. D. With fifty large quarto colored plates. In one large quarto volume, extra cloth. \$15 00.

LETTERS ON SYPHILIS, addressed to the Chief Editor of the Union Médicale. Translated by W. P. LATTIMORE, M. D. In one neat octavo volume, of 270 pages, extra cloth. \$2 00.

RIGBY (EDWARD), M. D.,

Senior Physician to the General Lying-in Hospital, &c.

A SYSTEM OF MIDWIFERY. With Notes and Additional Illustrations.

Second American Edition. One volume octavo, extra cloth, 422 pages. \$2 50.

BY THE SAME AUTHOR. (*Now Ready.*)

ON THE CONSTITUTIONAL TREATMENT OF FEMALE DISEASES.

In one neat royal 12mo. volume, extra cloth, of about 250 pages. \$1 00. *Now Ready.*

The following condensed summary of the contents will show the topics treated in this little volume. The aim of the author has been throughout to present sound practical views of the important subjects under consideration; and without entering into theoretical disputations and disquisitions to embody the results of his long and extended experience in such a condensed form as would be easily accessible to the practitioner.

CHAPTER I. Amenorrhœa.—II. Dysmenorrhœa.—III. Menorrhagia.—IV. Uterine and Vaginal Discharges.—V. Inflammation of the Os and Cervix.—VI. Ulceration of the Os and Cervix.—VII. Displacement of the Uterus.—VIII. Retroversion.—IX. Anteversion.—X. Prolapsus Vesicæ.—XI. Prolapsus Uteri.—XII. Fibrous Tumor of the Uterus.—XIII. Malignant Disease of the Uterus.—XIV.—Cauliflower Excrescence of the Os.—XV. Corroding Ulcer of the Os.—XVI. Pruritus Pudendi.—XVII. Vascular Tumor of Orifice of Urethra.—XVIII. Ovarian Affections.—XIX. Displacement of the Ovary.—XX. Ovarian Tumors.

RAMSBOTHAM (FRANCIS H.), M.D.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND

SURGERY, in reference to the Process of Parturition. A new and enlarged edition, thoroughly revised by the Author. With Additions by W. V. KEATING, M. D. In one large and handsome imperial octavo volume, of 650 pages, strongly bound in leather, with raised bands; with sixty-four beautiful Plates, and numerous Wood-cuts in the text, containing in all nearly two hundred large and beautiful figures. (*Lately Issued, 1856.*) \$5 00.

In calling the attention of the profession to the new edition of this standard work, the publishers would remark that no efforts have been spared to secure for it a continuance and extension of the remarkable favor with which it has been received. The last London issue, which was considerably enlarged, has received a further revision from the author, especially for this country. Its passage through the press here has been supervised by Dr. Keating, who has made numerous additions with a view of presenting more fully whatever was necessary to adapt it thoroughly to American modes of practice. In its mechanical execution, a like superiority over former editions will be found.

From Prof. Hodge, of the University of Pa.

To the American public, it is most valuable, from its intrinsic undoubted excellence, and as being the best authorized exponent of British Midwifery. Its circulation will, I trust, be extensive throughout our country.

The publishers have shown their appreciation of the merits of this work and secured its success by the truly elegant style in which they have brought it out, excelling themselves in its production, especially in its plates. It is dedicated to Prof. Meigs, and has the emphatic endorsement of Prof. Hodge, as the best exponent of British Midwifery. We know of no text-book which deserves in all respects to be more highly recommended to students, and we could wish to see it in the hands of every practitioner, for they will find it invaluable for reference.—*Med. Gazette.*

But once in a long time some brilliant genius rears his head above the horizon of science, and illuminates and purifies every department that he investigates; and his works become types, by which innumerable imitators model their feeble productions. Such a genius we find in the younger Ramsbotham, and such a type we find in the work now before us. The binding, paper, type, the engravings and woodcuts are all so excellent as to make this book one of the finest specimens of the art of printing that have given such a world-wide reputation to its enterprising and liberal publishers. We welcome Ramsbotham's Principles and Practice of Obstetric Medi-

cine and Surgery to our library, and confidently recommend it to our readers, with the assurance that it will not disappoint their most sanguine expectations.—*Western Lancet.*

It is unnecessary to say anything in regard to the utility of this work. It is already appreciated in our country for the value of the matter, the clearness of its style, and the fulness of its illustrations. To the physician's library it is indispensable, while to the student as a text-book, from which to extract the material for laying the foundation of an education on obstetrical science, it has no superior.—*Ohio Med. and Surg. Journal.*

We will only add that the student will learn from it all he need to know, and the practitioner will find it, as a book of reference, surpassed by none other.—*Stethoscope.*

The character and merits of Dr. Ramsbotham's work are so well known and thoroughly established, that comment is unnecessary and praise superfluous. The illustrations, which are numerous and accurate, are executed in the highest style of art. We cannot too highly recommend the work to our readers.—*St. Louis Med. and Surg. Journal.*

ROKITANSKY (CARL), M.D.,

Curator of the Imperial Pathological Museum, and Professor at the University of Vienna, &c.

A MANUAL OF PATHOLOGICAL ANATOMY. Four volumes, octavo, bound in two, extra cloth, of about 1200 pages. (*Now Ready.*) \$5 50

Vol. I.—Manual of General Pathological Anatomy. Translated by W. E. SWAINE.

Vol. II.—Pathological Anatomy of the Abdominal Viscera. Translated by EDWARD SIEVEKING, M. D.

Vol. III.—Pathological Anatomy of the Bones, Cartilages, Muscles, and Skin, Cellular and Fibrous Tissue, Serous and Mucous Membrane, and Nervous System. Translated by C. H. MOORE.

Vol. IV.—Pathological Anatomy of the Organs of Respiration and Circulation. Translated by G. E. DAY.

To render this large and important work more easy of reference, and at the same time less cumbersome and costly, the four volumes have been arranged in two, retaining, however, the separate paging, &c.

The publishers feel much pleasure in presenting to the profession of the United States the great work of Prof. Rokitansky, which is universally referred to as the standard of authority by the pathologists of all nations. Under the auspices of the Sydenham Society of London, the combined labor of four translators has at length overcome the almost insuperable difficulties which have so long prevented the appearance of the work in an English dress, while the additions made from various papers and essays of the author present his views on all the topics embraced, in their latest published form. To a work so widely known, eulogy is unnecessary, and the publishers would merely state that it is said to contain the results of not less than THIRTY THOUSAND *post-mortem* examinations made by the author, diligently compared, generalized, and wrought into one complete and harmonious system.

The profession is too well acquainted with the reputation of Rokitansky's work to need our assurance that this is one of the most profound, thorough, and valuable books ever issued from the medical press. It is *sui generis*, and has no standard of comparison. It is only necessary to announce that it is issued in a form as cheap as is compatible with its size and preservation, and its sale follows as a matter of course. No library can be called complete without it.—*Buffalo Med. Journal.*

An attempt to give our readers any adequate idea of the vast amount of instruction accumulated in these volumes, would be feeble and hopeless. The effort of the distinguished author to concentrate in a small space his great fund of knowledge, has

so charged his text with valuable truths, that any attempt of a reviewer to epitomize is at once paralyzed, and must end in a failure.—*Western Lancet.*

As this is the highest source of knowledge upon the important subject of which it treats, no real student can afford to be without it. The American publishers have entitled themselves to the thanks of the profession of their country, for this timely and beautiful edition.—*Nashville Journal of Medicine.*

As a book of reference, therefore, this work must prove of inestimable value, and we cannot too highly recommend it to the profession.—*Charleston Med. Journal and Review*, Jan. 1856.

This book is a necessity to every practitioner.—*Am. Med. Monthly.*

ROYLE'S MATERIA MEDICA AND THERAPEUTICS; including the Preparations of the Pharmacopœias of London, Edinburgh, Dublin, and of the United States. With many new medicines. Edited by JOSEPH CARSON, M. D. With ninety-eight illustrations. In one large octavo volume, extra cloth, of about 700 pages. \$3 00.

SMITH (HENRY H.), M. D.,

Professor of Surgery in the University of Pennsylvania, &c.

MINOR SURGERY; or, Hints on the Every-day Duties of the Surgeon. Illustrated by two hundred and forty-seven illustrations. Third and enlarged edition. In one handsome royal 12mo. volume. pp. 456. In leather, \$2 25; extra cloth, \$2 00.

And a capital little book it is. . . Minor Surgery, we repeat, is really Major Surgery, and anything which teaches it is worth having. So we cordially recommend this little book of Dr. Smith's.—*Med.-Chir. Review*.

This beautiful little work has been compiled with a view to the wants of the profession in the matter of bandaging, &c., and well and ably has the author performed his labors. Well adapted to give the requisite information on the subjects of which it treats.—*Medical Examiner*.

The directions are plain, and illustrated throughout with clear engravings.—*London Lancet*.

One of the best works they can consult on the subject of which it treats.—*Southern Journal of Medicine and Pharmacy*.

A work such as the present is therefore highly useful to the student, and we commend this one to their attention.—*American Journal of Medical Sciences*.

No operator, however eminent, need hesitate to consult this unpretending yet excellent book. Those who are young in the business would find Dr. Smith's treatise a necessary companion, after once understanding its true character.—*Boston Med. and Surg. Journal*.

No young practitioner should be without this little volume; and we venture to assert, that it may be consulted by the senior members of the profession with more real benefit, than the more voluminous works.—*Western Lancet*.

BY THE SAME AUTHOR, AND

HORNER (WILLIAM E.), M. D.,

Late Professor of Anatomy in the University of Pennsylvania.

AN ANATOMICAL ATLAS, illustrative of the Structure of the Human Body. In one volume, large imperial octavo, extra cloth, with about six hundred and fifty beautiful figures. \$3 00.

These figures are well selected, and present a complete and accurate representation of that wonderful fabric, the human body. The plan of this Atlas, which renders it so peculiarly convenient for the student, and its superb artistic execution, have been already pointed out. We must congratulate

late the student upon the completion of this Atlas, as it is the most convenient work of the kind that has yet appeared; and we must add, the very beautiful manner in which it is "got up" is so creditable to the country as to be flattering to our national pride.—*American Medical Journal*.

SARGENT (F. W.), M. D.

ON BANDAGING AND OTHER OPERATIONS OF MINOR SURGERY.

Second edition, enlarged. One handsome royal 12mo. vol., of nearly 400 pages, with 182 woodcuts. (*Now Ready*, 1856.) Extra cloth, \$1 40; leather, \$1 50.

This very useful little work has long been a favorite with practitioners and students. The recent call for a new edition has induced its author to make numerous important additions. A slight alteration in the size of the page has enabled him to introduce the new matter, to the extent of some fifty pages of the former edition, at the same time that his volume is rendered still more compact than its less comprehensive predecessor. A double gain in thus effected, which, in a *vade-mecum* of this kind, is a material improvement.—*Am. Medical Journal*.

Sargent's Minor Surgery has always been popular, and deservedly so. It furnishes that knowledge of the most frequently requisite performances of surgical art which cannot be entirely understood by attending clinical lectures. The art of bandaging, which is regularly taught in Europe, is very frequently overlooked by teachers in this country; the student and junior practitioner, therefore, may often require that knowledge which this little volume so tersely and happily supplies. It is neatly printed and cop-

iously illustrated by the enterprising publishers, and should be possessed by all who desire to be thoroughly conversant with the details of this branch of our art.—*Charleston Med. Journ. and Review*, March, 1856.

A work that has been so long and favorably known to the profession as Dr. Sargent's Minor Surgery, needs no commendation from us. We would remark, however, in this connection, that minor surgery seldom gets that attention in our schools that its importance deserves. Our larger works are also very defective in their teaching on these small practical points. This little book will supply the void which all must feel who have not studied its pages.—*Western Lancet*, March, 1856.

We confess our indebtedness to this little volume on many occasions, and can warmly recommend it to our readers, as it is not above the consideration of the oldest and most experienced.—*American Lancet*, March, 1856.

SKEY'S OPERATIVE SURGERY. In one very handsome octavo volume, extra cloth, of over 650 pages, with about one hundred woodcuts. \$3 25.

STANLEY'S TREATISE ON DISEASES OF THE BONES. In one volume, octavo, extra cloth, 286 pages. \$1 50.

SOPLY ON THE HUMAN BRAIN; its Structure, Physiology, and Diseases. From the Second and

much enlarged London edition. In one octavo volume, extra cloth, of 500 pages, with 120 woodcuts. \$2 00.

SIMON'S GENERAL PATHOLOGY, as conducive to the Establishment of Rational Principles for the prevention and Cure of Disease. In one neat octavo volume, extra cloth, of 212 pages. \$1 25.

STILLÉ (ALFRED), M. D.

PRINCIPLES OF GENERAL AND SPECIAL THERAPEUTICS In handsome octavo. (*Preparing*.)

**SHARPEY (WILLIAM), M. D., JONES QUAIN, M. D., AND
RICHARD QUAIN, F. R. S., &c.**

HUMAN ANATOMY. Revised, with Notes and Additions, by **JOSEPH LEIDY**, M. D., Professor of Anatomy in the University of Pennsylvania. Complete in two large octavo volumes, leather, of about thirteen hundred pages. Beautifully illustrated with over five hundred engravings on wood. \$6 00.

It is indeed a work calculated to make an era in anatomical study, by placing before the student every department of his science, with a view to the relative importance of each; and so skilfully have the different parts been interwoven, that no one who makes this work the basis of his studies, will hereafter have any excuse for neglecting or undervaluing any important particulars connected with the structure of the human frame; and whether the bias of his mind lead him in a more especial manner to surgery, physic, or physiology,

he will find here a work at once so comprehensive and practical as to defend him from exclusiveness on the one hand, and pedantry on the other.—*Journal and Retrospect of the Medical Sciences.*

We have no hesitation in recommending this treatise on anatomy as the most complete on that subject in the English language; and the only one, perhaps, in any language, which brings the state of knowledge forward to the most recent discoveries.—*The Edinburgh Med. and Surg. Journal.*

SMITH (W. TYLER), M. D.,
Physician Accoucheur to St. Mary's Hospital, &c.

**ON PARTURITION, AND THE PRINCIPLES AND PRACTICE OF
OBSTETRICS.** In one royal 12mo. volume, extra cloth, of 400 pages. \$1 25.

BY THE SAME AUTHOR.—(Just Issued.)

**A PRACTICAL TREATISE ON THE PATHOLOGY AND TREATMENT
OF LEUCORRHOEA.** With numerous illustrations. In one very handsome octavo volume, extra cloth, of about 250 pages. \$1 50.

We decide this book to be one of the most useful monographs which has appeared in this country. What was before unutterable confusion in regard to its subject has now the order, regularity, and harmony of a most beautiful science. Dr. Smith has placed the whole profession directly, and mankind

indirectly, under abiding obligations.—*Nashville's Journ. of Medicine.*

We hail the appearance of this practical and invaluable work, therefore, as a real acquisition to our medical literature.—*Medical Gazette.*

SIBSON (FRANCIS), M. D.,
Physician to St. Mary's Hospital.

MEDICAL ANATOMY. Illustrating the Form, Structure, and Position of the Internal Organs in Health and Disease. In large imperial quarto, with splendid colored plates. To match "Maclise's Surgical Anatomy." Part I. (Preparing.)

SCHOEDLER (FRIEDRICH), PH. D.,
Professor of the Natural Sciences at Worms, &c.

THE BOOK OF NATURE; an Elementary Introduction to the Sciences of Physics, Astronomy, Chemistry, Mineralogy, Geology, Botany, Zoology, and Physiology. First American edition, with a Glossary and other Additions and Improvements; from the second English edition. Translated from the sixth German edition, by **HENRY MEDLOCK, F. C. S., &c.** In one thick volume, small octavo, extra cloth, of about seven hundred pages, with 679 illustrations on wood. Suitable for the higher Schools and private students. (Now Ready.) \$1 80.

TANNER (T. H.), M. D.,
Physician to the Hospital for Women, &c.

A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS.

To which is added The Code of Ethics of the American Medical Association. Second American Edition. In one neat volume, small 12mo. Price in extra cloth, 87½ cents; flexible style, for the pocket, 80 cents. (Lately Published.)

In this admirable little work the author's object has been to give the young practitioner that kind of information which enables him to make practical application of the knowledge acquired by his studies, and which is not to be found in the text-books. Such a manual has been much wanted, as it fills a void which has long been felt, but which there has hitherto been no attempt to supply. That the author has succeeded in his endeavor, is sufficiently shown by the unusually favorable reception which the work has already received, although only just published.

Dr. Tanner has, in a happy and successful manner, indicated the leading particulars to which, in the clinical study of a case of disease, the attention of the physician is to be directed, the value and import of the various abnormal phenomena detected, and the several instrumental and accessory means which may be called into requisition to facilitate diagnosis and increase its certainty.—*Am. Journal of Med. Sciences.*

In this small work is collected a fund of such information as the student at the commencement, and even during the continuance of his studies, is often sadly troubled to know where to look for.—*Montreal Med. Chronicle.*

The work is an honor to its writer, and must obtain a wide circulation by its intrinsic merit alone. It seems to us that but slight effort on the part of the publishers will be requisite to exhaust even a large edition. Suited alike to the wants of students and

practitioners, it has only to be seen, to win for itself a place upon the shelves of every medical library. Nor will it be "shelved" long at a time; if we mistake not, it will be found, in the best sense of the homely but expressive word, "handy." The style is admirably clear, while it is so sententious as not to burden the memory. The arrangement is, to our mind, unexceptionable. The work, in short, deserves the heartiest commendation.—*Boston Med. and Surg. Journal*

We cordially recommend every young practitioner who wishes to reap the greatest possible benefit from his observation of disease to make this book his daily companion.—*New Hampshire Journal of Medicine.*

As a convenient and suggestive book of reference, we accord it our hearty praise.—*Va. Med. and Surg. Journal.*

TAYLOR (ALFRED S.), M. D., F. R. S.,

Lecturer on Medical Jurisprudence and Chemistry in Guy's Hospital.

MEDICAL JURISPRUDENCE. Fourth American, from the fifth improved and enlarged English Edition. With Notes and References to American Decisions, by EDWARD HARTSHORNE, M. D. In one large octavo volume, leather, of over seven hundred pages. (*Just Ready*, June, 1856.) \$3 00.

This standard work has lately received a very thorough revision at the hands of the author, who has introduced whatever was necessary to render it complete and satisfactory in carrying out the objects in view. The editor has likewise used every exertion to make it equally thorough with regard to all matters relating to the practice of this country. In doing this, he has carefully examined all that has appeared on the subject since the publication of the last edition, and has incorporated all the new information thus presented. The work has thus been considerably increased in size, notwithstanding which, it has been kept at its former very moderate price, and in every respect it will be found worthy of a continuance of the remarkable favor which has carried it through so many editions on both sides of the Atlantic. A few notices of the former editions are appended.

We know of no work on Medical Jurisprudence which contains in the same space anything like the same amount of valuable matter.—*N. Y. Journal of Medicine*.

No work upon the subject can be put into the hands of students either of law or medicine which will engage them more closely or profitably; and none could be offered to the busy practitioner of either calling, for the purpose of casual or hasty reference, that would be more likely to afford the aid desired. We therefore recommend it as the best and safest manual for daily use.—*American Journal of Medical Sciences*.

So well is this work known to the members both of the medical and legal professions, and so highly is it appreciated by them, that it cannot be necessary for us to say a word in its commendation; its having already reached a fourth edition being the best possible testimony in its favor. The author has obviously subjected the entire work to a very careful revision.—*Brit. and Foreign Med. Chirurg. Review*.

This work of Dr. Taylor's is generally acknowledged to be one of the ablest extant on the subject of medical jurisprudence. It is certainly one of the

most attractive books that we have met with; supplying so much both to interest and instruct, that we do not hesitate to affirm that after having once commenced its perusal, few could be prevailed upon to desist before completing it. In the last London edition, all the newly observed and accurately recorded facts have been inserted, including much that is recent of Chemical, Microscopical, and Pathological research, besides papers on numerous subjects never before published.—*Charleston Medical Journal and Review*.

It is not excess of praise to say that the volume before us is the very best treatise extant on Medical Jurisprudence. In saying this, we do not wish to be understood as detracting from the merits of the excellent works of Beck, Ryan, Traill, Guy, and others; but in interest and value we think it must be conceded that Taylor is superior to anything that has preceded it. The author is already well known to the profession by his valuable treatise on Poisons; and the present volume will add materially to his high reputation for accurate and extensive knowledge and discriminating judgment.—*N. W. Medical and Surgical Journal*.

BY THE SAME AUTHOR.

ON POISONS, IN RELATION TO MEDICAL JURISPRUDENCE AND MEDICINE. Edited, with Notes and Additions, by R. E. GRIFFITH, M. D. In one large octavo volume, leather, of 688 pages. \$3 00

Now Complete (March, 1857).

TODD (ROBERT BENTLEY), M. D., F. R. S.,

Professor of Physiology in King's College, London; and

WILLIAM BOWMAN, F. R. S.,

Demonstrator of Anatomy in King's College, London.

THE PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN. With about three hundred large and beautiful illustrations on wood. Complete in one large octavo volume, of 950 pages, leather. Price \$4 50.

The very great delay which has occurred in the completion of this work has arisen from the desire of the authors to verify by their own examination the various questions and statements presented, thus rendering the work one of peculiar value and authority. By the wideness of its scope and the accuracy of its facts it thus occupies a position of its own, and becomes necessary to all physiological students.

Gentlemen who have received portions of this work, as published in the "MEDICAL NEWS AND LIBRARY," can now complete their copies, if immediate application be made. It will be furnished as follows, free by mail, in paper covers, with cloth backs.

PARTS I., II., III. (pp. 25 to 552), \$2 50.

PART IV. (pp. 553 to end, with Title, Preface, Contents, &c.), \$2 00.

Or, PART IV., SECTION II. (pp. 725 to end, with Title, Preface, Contents, &c.), \$1 25.

In the present part (third) some of the most difficult subjects in Anatomy and Physiology are handled in the most masterly manner. Its authors have stated that this work was intended "for the use of the student and practitioner in medicine and surgery," and we can recommend it to both, confident that it is the most perfect work of its kind. We cannot conclude without strongly recommending the present work to all classes of our readers, recognizing talent and depth of research in every page, and believing, as we do, that the diffusion of such knowledge will certainly tend to elevate the sciences of Medicine and Surgery.—*Dublin Quarterly Journal of Medical Sciences*.

TODD (R. B.), M. D., F. R. S., &c.

CLINICAL LECTURES ON CERTAIN DISEASES OF THE URINARY ORGANS AND ON DROPSIES. In one octavo volume. (*Just Ready*.) \$1 50.

The valuable practical nature of Dr. Todd's writings have deservedly rendered them favorites with the profession, and the present volume, embodying the medical aspects of a class of diseases not elsewhere to be found similarly treated, can hardly fail to supply a want long felt by the practitioner.

WATSON (THOMAS), M. D., &c.

LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC.

Third American edition, revised, with Additions, by D. FRANCIS CONDIE, M. D., author of a "Treatise on the Diseases of Children," &c. In one octavo volume, of nearly eleven hundred large pages, strongly bound with raised bands. \$3 25.

To say that it is the very best work on the subject now extant, is but to echo the sentiment of the medical press throughout the country.—*N. O. Medical Journal*.

Of the text-books recently republished Watson is very justly the principal favorite.—*Holmes's Rep. to Nat. Med. Assoc.*

By universal consent the work ranks among the very best text-books in our language.—*Illinois and Indiana Med. Journal*.

Regarded on all hands as one of the very best, if not the very best, systematic treatise on practical medicine extant.—*St. Louis Med. Journal*.

Confessedly one of the very best works on the principles and practice of physic in the English or any other language.—*Med. Examiner*.

As a text-book it has no equal; as a compendium of pathology and practice no superior.—*New York Annalist*.

We know of no work better calculated for being placed in the hands of the student, and for a text-book; on every important point the author seems to have posted up his knowledge to the day.—*Amer. Med. Journal*.

One of the most practically useful books that ever was presented to the student.—*N. Y. Med. Journal*.

WILSON (ERASMUS), M. D., F. R. S.,

Lecturer on Anatomy, London.

A SYSTEM OF HUMAN ANATOMY, General and Special. Fourth American,

from the last English edition. Edited by PAUL B. GODDARD, A. M., M. D. With two hundred and fifty illustrations. Beautifully printed, in one large octavo volume, leather, of nearly six hundred pages. \$3 00.

In many, if not all the Colleges of the Union, it has become a standard text-book. This, of itself, is sufficiently expressive of its value. A work very desirable to the student; one, the possession of which will greatly facilitate his progress in the study of Practical Anatomy.—*New York Journal of Medicine*.

Its author ranks with the highest on Anatomy.—*Southern Medical and Surgical Journal*.

It offers to the student all the assistance that can be expected from such a work.—*Medical Examiner*.

The most complete and convenient manual for the student we possess.—*American Journal of Medical Science*.

In every respect, this work as an anatomical guide for the student and practitioner, merits our warmest and most decided praise.—*London Medical Gazette*.

BY THE SAME AUTHOR. (Now Ready.)

THE DISSECTOR'S MANUAL; or, Practical and Surgical Anatomy. Third

American, from the last revised and enlarged English edition. Modified and rearranged, by WILLIAM HUNT, M. D., Demonstrator of Anatomy in the University of Pennsylvania. In one large and handsome royal 12mo. volume, leather, of 582 pages, with 154 illustrations. \$2 00.

The modifications and additions which this work has received in passing recently through the author's hands, is sufficiently indicated by the fact that it is enlarged by more than one hundred pages, notwithstanding that it is printed in smaller type, and with a greatly enlarged page.

It remains only to add, that after a careful examination, we have no hesitation in recommending this work to the notice of those for whom it has been expressly written—the students—as a guide possess-

ing very superior claims, well calculated to facilitate their studies, and render their labor less irksome, by constantly keeping before them definite objects of interest.—*The Lancet*.

BY THE SAME AUTHOR.

ON DISEASES OF THE SKIN. Third American, from the third London edition.

In one neat octavo volume, of about five hundred pages, extra cloth. \$1 75.

The "Diseases of the Skin," by Mr. Erasmus Wilson, may now be regarded as the standard work in that department of medical literature.—*Medico-Chirurgical Review*.

BY THE SAME AUTHOR.

ON CONSTITUTIONAL AND HEREDITARY SYPHILIS, AND ON SYPHILITIC ERUPTIONS.

In one small octavo volume, extra cloth, beautifully printed, with four exquisite colored plates, presenting more than thirty varieties of syphilitic eruptions. \$2 25.

BY THE SAME AUTHOR. (Just Issued.)

HEALTHY SKIN; A Popular Treatise on the Skin and Hair, their Preservation and Management. Second American, from the fourth London edition.

One neat volume, royal 12mo., extra cloth, of about 300 pages, with numerous illustrations. \$1 00; paper cover, 75 cents.

WHITEHEAD ON THE CAUSES AND TREATMENT OF ABORTION AND STERILITY.

Second American Edition. In one volume, octavo, extra cloth, pp. 308. \$1 75.

WALSHE ON DISEASES OF THE HEART, LUNGS, AND APPENDAGES; their Symptoms and Treatment.

In one handsome volume, extra cloth, large royal 12mo., 512 pages. \$1 50.

WILDE (W. R.),

Surgeon to St. Mark's Ophthalmic and Aural Hospital, Dublin.

AURAL SURGERY, AND THE NATURE AND TREATMENT OF DISEASES OF THE EAR.

In one handsome octavo volume, extra cloth, of 476 pages, with illustrations. \$2 80.

This work certainly contains more information on the subject to which it is devoted than any other with which we are acquainted. We feel grateful to

the author for his manful effort to rescue this department of surgery from the hands of the empirics who nearly monopolize it.

WEST (CHARLES), M. D.,

Accoucheur to and Lecturer on Midwifery at St. Bartholomew's Hospital, Physician to the Hospital for Sick Children, &c.

LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD.

Second American, from the Second and Enlarged London edition. In one volume, octavo, extra cloth, of nearly five hundred pages. (*Just Issued.*) \$2 00.

We take leave of Dr. West with great respect for his attainments, a due appreciation of his acute powers of observation, and a deep sense of obligation for this valuable contribution to our professional literature. His book is undoubtedly in many respects the best we possess on diseases of children. *Dublin Quarterly Journal of Medical Science.*

Dr. West has placed the profession under deep ob-

ligation by this able, thorough, and finished work upon a subject which almost daily taxes to the utmost the skill of the general practitioner. He has with singular felicity threaded his way through all the tortuous labyrinths of the difficult subject he has undertaken to elucidate, and has in many of the darkest corners left a light, which will never be extinguished.—*Nashville Medical Journal.*

BY THE SAME AUTHOR. (*Nearly Ready.*)

Publishing in the "Medical News and Library," for 1856.

LECTURES ON THE DISEASES OF WOMEN. In two parts.

PART I. 8vo. of about 300 pages, comprising the DISEASES OF THE UTERUS.

PART II. (*Preparing*), will contain DISEASES OF THE OVARIES, and of all the parts connected with the UTERUS; of the BLADDER, VAGINA, AND EXTERNAL ORGANS.

The object of the author in this work is to present a complete but succinct treatise on Female Diseases, embodying the results of his experience during the last ten years at St. Bartholomew's and the Midwifery Hospitals, as well as in private practice. The characteristics which have secured to his former works so favorable a reception, cannot fail to render the present volume a standard authority on its important subject. To show the general scope of the work, an outline of the Contents of Part I. is subjoined.

Lectures I, II.—INTRODUCTORY—Symptoms—Examination of Symptoms—Modes of Examination. *Lectures III, IV, V*—DISORDERS OF MENSTRUATION, Amenorrhœa, Menorrhagia, Dysmenorrhœa. *Lectures VI, VII, VIII*—INFLAMMATION OF THE UTERUS, Hypertrophy, Acute Inflammation, Chronic Inflammation, Ulceration of the Os Uteri, Cervical Leucorrhœa. *Lectures IX, X, XI, XII, XIII*—MISPLACEMENT OF THE UTERUS, Prolapsus, Anteversion, Retroversion, Inversion. *Lectures XIV, XV, XVI, XVII*—UTERINE TUMORS AND OUTGROWTHS, Mucous, Fibro-cellular, and Glandular Polypi, Mucous Cysts, Fibrinous Polypi, Fibrous Tumors, Fibrous Polypi, Fatty Tumors, Tubercular Diseases. *Lectures XVIII, XIX, XX*—CANCER OF THE UTERUS.

PART II. will receive an equally extended treatment, rendering the whole an admirable text-book for the student, and a reliable work for reference by the practitioner.

BY THE SAME AUTHOR. (*Just Issued*)

AN ENQUIRY INTO THE PATHOLOGICAL IMPORTANCE OF ULCERATION OF THE OS UTERI. In one neat octavo volume, extra cloth. \$1 00.**WILLIAMS (C. J. B.), M. D., F. R. S.,**

Professor of Clinical Medicine in University College, London, &c.

PRINCIPLES OF MEDICINE; comprising General Pathology and Therapeutics, and a brief general view of Etiology, Nosology, Semeiology, Diagnosis, Prognosis, and Hygienics. Edited, with Additions, by MEREDITH CLYMER, M. D. Fourth American, from the last and enlarged London edition. In one octavo volume, leather, of 476 pages. \$2 50.

The best exposition in our language, or, we believe, in any language, of rational medicine, in its present improved and rapidly improving state.—*British and Foreign Medico-Chirurg. Review.*

Few books have proved more useful, or met with a more ready sale than this, and no practitioner should regard his library as complete without it.—*Ohio Med. and Surg. Journal.*

WHAT TO OBSERVE**AT THE BEDSIDE AND AFTER DEATH, IN MEDICAL CASES.**

Published under the authority of the London Society for Medical Observation. A new American, from the second and revised London edition. In one very handsome volume, royal 12mo., extra cloth. (*Just Issued.*) \$1 00.

To the observer who prefers accuracy to blunders and precision to carelessness, this little book is invaluable.—*N. H. Journal of Medicine.*

One of the finest aids to a young practitioner we have ever seen.—*Peninsular Journal of Medicine.*

YOUATT (WILLIAM), V. S.

THE HORSE. A new edition, with numerous illustrations; together with a general history of the Horse; a Dissertation on the American Trotting Horse; how Trained and Jockeyed; an Account of his Remarkable Performances; and an Essay on the Ass and the Mule. By J. S. SKINNER, formerly Assistant Postmaster-General, and Editor of the Turf Register. One large octavo volume, extra cloth. \$1 50.

The attention of all who keep horses is requested to this handsome and complete edition of a work which is recognized as the standard authority on all matters connected with veterinary medicine. The very low price at which it is now offered, FREE BY MAIL, places it within the reach of every one.

BY THE SAME AUTHOR.

THE DOG. Edited by E. J. LEWIS, M. D. With numerous and beautiful illustrations. In one very handsome volume, crown 8vo., crimson cloth, gilt. \$1 25.





NATIONAL LIBRARY OF MEDICINE



NLM 03274535 1